

# Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle

*by* Widya Paramita

---

**Submission date:** 27-Nov-2018 03:40PM (UTC+0800)

**Submission ID:** 1045552684

**File name:** 11-MKH\_vol\_26\_No\_2\_th\_2010.pdf (522.55K)

**Word count:** 2978


**Character count:** 16158

ISSN 0215-8930

# MEDIA

## Kedokteran Hewan

*Veterinary Medicine Journal*



MKH (Vet.Med.J.) Vol. 26 No. 2 Pp. 76-138 Surabaya, May 2010 ISSN 0215-8930

Accreditation by Director General of Higher Education, No. 108/Dikti/Kept/2007, August 23, 2007

## Table of Contents

No.	Title	Page
1	Antiproliferation Effect of the n-Hexanal Extract of Kesum (Polygonum minus) at the Cells of the Rat Lung Cancer Effort Exploiting of Kesum as Drug of Lung Cancer	128 - 133
2	Antiproliferation Effect of the n-Hexanal Extract of Kesum (Polygonum minus) at the Cells of the Rat Lung Cancer Effort Exploiting of Kesum as Drug of Lung Cancer	128 - 133
3	Invitro Maturation of Domestic Cat (Felis catus Ovarian Oocytes fater being stored at 4o C	76 - 79
4	Concentrations of Liver and Plasma Malondealdehyde (MDA) of sprague Dawley Rats Injected by Ketamine-Xylazine HCL Anesthetics	80 - 83
5	Protein Profile of Sporozoite of Leucocytozoon sp. from Culicoides sp.	84 - 87
6	Expression of CD4+ T Cell, Interleukin-2 and Interleukin-4 on Splenectomy Balb/c Mice Post-Exposure Salmonela typhi	88 - 91
7	The Profile of Antioxidant Superoxide Dismutase (SOD) in Liver of Isoflavon, Zn, and Vitamins E-treated Rats	98 - 105
8	Species Identification of Coagulase Positive Staphylococci (CPS) by Multiplex Polymerase Chain Reaction (PCR)	106 - 111
9	Species Identification of coagulase positive staphylococci (CPS) by multiplex Polymerase Chain Reaction (PCR)	108 - 111
10	Mechanisms Underlying Abortion in Dexamethatione Treatment at Ewe in Midgestation	112 - 117
11	Antibody Against Prolactin: as a Teraphy for Molting Duck	118 - 123
12	<b>Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle</b>	124 - 127
13	Antiproliferation Effect of the n-Heksana Extract of Kesum (Polygomon minus) at the Cells of the Rat Lung Cancer: Effort Exploiting of Kesum as drug of Lung Cancer	128 - 133
14	Effect of Supplementation of Ccows or Direcly to Calves on Calf Performance in Bali Cattle Grazing Communal Pastures	134 - 138

## **Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle**

## **Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle**

1. Mirni Lamid --> Dosen Fakultas Kedokteran Hewan / mirnilamid@yahoo.com
2. Ni Nyoman Tri Puspaningsih --> Dosen Fakultas Sains dan Teknologi
3. Widya Paramita --> Dosen Fakultas Kedokteran Hewan

### **Abstract**

Biotechnology in animal feed using biological treatment such as cellulase is mainly aimed to increase degrade the cellulose; the second most abundant polysaccharide in nature. Information of using cellulase as biocatalyst for increasing animal feed quality as ruminants feed is limited. Cellulose is the major component of cell wall agroindustry waste after hemicellulose. This cellulase comes from *Actinobacillus* sp. that were isolated from rumen fluid of beef cattle. The aim of this research was to determine activity of cellulase and characterization of optimum temperature and pH. In this research identification with carboxyl methylcellulose (CMC) has been done. Methods for cellulase activity was 3,5-dinitrosalicylic acid (DNS). The results showed that *Actinobacillus* which have positive activity of cellulase. Characterization of *Actinobacillus* sp. had optimum temperature 45°C with activity 1.42 U/ml and optimum pH 6 with activity 0.57 U/ml.

Keyword : cellulase, carboxyl, methyl, cellulose, temperature, beef, cattle,

### **Daftar Pustaka :**

1. **Bhat MK and Hazlewood GP., (2001).** Enzymology and Other Characteristics of Cellulases and Xylanases, In: *Enzymes in Farm Animal Nutrition* M.R. Bedford and G.G. Partridge (Eds).. UK : CABI Publ
2. **Beauchemin KA, Colombatto D, Morgavi DP and Yang WZ., (2003).** Use of exogenous fibrolytic enzymes to improve feed utilization by ruminant. - : *J Anim Sci*; 81: E37 - E47
3. **Bondi AA., (1987).** Animal Nutrition. London : Wiley-Interscience Publication
4. **Campbell Mary K., (1995).** Biochemistry. 2nd Ed. Mount Holyoke College. USA : Saunders College Publ
5. **Lowe SE, Michael K, Theodorou and Anthony PJT., (1987).** Cellulases and Xylanases of an Anaerobic Rumen Fungus Grown on Wheat Straw, Wheat Straw Hemicellulose, Cellulose, and Xylan. - : *Appl Environ Microbiol.*
6. **Mc. Donald P, Edwards RA and Greenhalgh JFD., (1995).** Animal Nutrition.. London : Logman
7. **Miller Gail Lorenz., (1960).** Use of Dinitrosalicylic acid reagent for determination of reducing sugar.. - : *Anal Chem*
8. **Orskov ER and Ryle., (1990).** Energy Nutrition in Ruminant. Elsevier Application.. New York : Jhon Willey and Sons
9. **Owens FN and Goestch AL., (1988).** Ruminant Fermentation In: D.C. Church (Ed0, *The Ruminant Animal Digestive Physiology and Nutrition.* New Jersey : A Reston Book Prentice Hall. Englewood Cliffs
10. **Ottaway JH and Apps DK., (1984).** Biochemistry. ELBS : Cambridge
11. **Puspaningsih NNT., (2004).** Isolation and characterization of  $\beta$ -Xylosidase and  $\alpha$ -L-Arabinofurosidase from *Bacillus thermoleovorans* IT-08. biotechnology of lignocellulose degradataion and biomass utiliozation.. Publ Co, Ltd. : Mie University
12. **Spolestra SF, Steg A and Beuvink JMW., (1990).** Application of cell wall degrading enzymes to grass silage.. *Pudoc; Agricultural biotechnology in focus in The Netherlands.* J.J. Dekkers, H.C. van der Plas and D.H. Vuijk (Eds)
13. **Sunna A and Antranikian G., (1997).** Xylanolytic enzymes from fungi and bacteria. *Ctit Rev Biotech* : Graham G. Stewart and Inge Russel (Eds)

## Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle

Mirni Lamid<sup>1</sup>, Ni Nyoman Tri Puspaningsih<sup>2</sup>, and Widya Paramita L<sup>1</sup>

<sup>1</sup> Department of Husbandry, Faculty of Veterinary Medicine, Airlangga University; <sup>2</sup> Department of Chemistry, Faculty of Science and Technology, Airlangga University, C Campus, Jl Mulyorejo, Surabaya, Indonesia; e-mail: mirnylamid@yahoo.com

### Abstract

Biotechnology in animal feed using biological treatment such as cellulase is mainly aimed to increase degrade the cellulose; the second most abundant polysaccharide in nature. Information of using cellulase as biocatalyst for increasing animal feed quality as ruminants feed is limited. Cellulose is the major component of cell wall agroindustry waste after hemicellulose. This cellulase comes from *Actinobacillus* sp. that were isolated from rumen fluid of beef cattle. The aim of this research was to determine activity of cellulase and characterization of optimum temperature and pH. In this research identification with carboxyl methylcellulose (CMC) has been done. Methods for cellulase activity was 3,5-dinitrosalicylic acid (DNS). The results showed that *Actinobacillus* which have positive activity of cellulase. Characterization of *Actinobacillus* sp. had optimum temperature 45 °C with activity 1.42 U/ml and optimum pH 6 with activity 0.57 U/ml.

**Key words:** cellulase, carboxyl methyl cellulose, temperature, beef cattle

### Introduction

The potential of agro industry waste product is great, but the problem occurred was the low value of nutrition signed the high content of crude fiber (cellulose, hemicelluloses, and lignin) and the low value of crude protein. Cellulose and hemicelluloses provided used a little part only and the most become waste product. It is caused cell wall of agro industry waste product realized lignifications, make a complex bound with lignin. Cellulose as an organic material compound abundant in this nature has a sharp filament like; it is content of cell wall of plant. Cellulose is the highest component of the plant. Cellulose is the glucose polymer connected with glycoside no branch. Most of animal can not digest cellulose because they don't have cellulase enzyme in their digestive tract. Ruminant can use cellulose as their nutrient source because in their rumen contain cellulolytic bacteria produced cellulase enzyme to hydrolyze cellulose (Campbell, 1994).

Cellulose molecule is the polysaccharide with bound  $\beta$ -1-4 glycoside that is difficult to digested by rumen bacteria therefore digestibility of cellulose was very low (Mc. Donald *et al.*, 1995). Rumen microbes are main source of cellulase enzyme producer but digestibility of agro industry waste still low so need a

process before. Utilizing of cellulase enzyme group will increase nutrition value of agro industry waste product. Cellulase enzyme contains 3 enzyme components, that is: component C1 ( $\beta$ -1, 4-glucan cellobiohydrolase or exo- $\beta$ -1, 4-glucanase), component Cc (endo- $\beta$ -1, 4-glucanase) and component cellobyose ( $\beta$ -glucosidase). The centre point of cellulose degradation provide in the break of bound 1.4  $\beta$ -glycoside. The breaking of bound 1.4  $\beta$ -glycoside caused cellulose hydrolyzed become simpler component that is oligosaccharide (mainly cellobyose). Then it will hydrolyze polysaccharide (mainly glucose). The breaking of 1.4  $\beta$ -glycoside bound done by cellulase enzyme group (Bondy, 1987).

Improvement of hydrolyze enzymatically was a new prospect to handle ligno cellulose become the feed for ruminant. Biodegradation of cellulase enzymatically accomplished by cellulase enzyme produced by much kind of microbes, such as: fungi, bacteria or *actinomyces*, *alga*, *oomycetes*, ruminant (Campbell, 1995).

Cellulase enzyme has very big industry application that is textile industry, pharmacy, paper and pulp, food and animal feed, and chemistry industry. Application of cellulase enzyme becomes one of component in chicken rations (Sunna and

Antranikian, 1997). Cellulose enzyme gather with xylanase can break glucose reduce the fermentation result of structural polysaccharide of the plant cell wall at the grass cellulase process, thus utilizing of this enzyme can increase consumption and digestibility of ruminant (Spoelstra *et al.*, 1990). The giving of feed additive enzyme mainly cellulase and xylanase can increase feed efficiency for beef cattle and dairy cattle (Beauchemin *et al.*, 2003).

The protein structure of enzyme can influence catalytic activities. Catalytic activities of enzyme will be loss if protein denaturizing occurred. The occurring cause of protein denaturizing is extreme pH, high temperature, and influencing of other component such as detergent, ionic, organic solution, concentrate urea, big anion of strong acid and chaotropic ion (I-,SCN-)(Ottoway and Apps, 1984). Considered from the biochemistry structural, there are some difference between enzyme thermophilic and mesophilic. Beside from disulfide content, hydrophobic interaction level at the molecule has enzyme stability. The optimum temperature of thermophilic enzyme activities is about 50°-80° C, and for thermophilic enzyme is at the room temperature, about 25°-50° C. if the mesophilic enzyme placed at higher temperature will realize denaturizing.

This research used bacteria isolation from beef cattle rumen, *Actinobacillus* sp. The research was to investigate cellulase activities then accomplished characterization to determine optimal temperature and pH of cellulose enzyme from the bacteria isolation.

## Materials and Methods

### Sample

The sample used was bacteria isolation, *Actinobacillus* sp. produced from beef cattle rumen.

### Production of fluid media

Fluid media contains of 0.2 g trypton; 0.1 yeast extract; 0.2 g NaCl. All materials dissolved at the Erlenmeyer tube 100 ml with 20 ml aquadest, and then closed by cotton and aluminum foil. After that it sterilized by autoclave for 45 minutes at 121°C, 1 atm.

### Production of solid media

Fluid media contains of 0.2 g trypton; 0.1 yeast extract; 0.2 g NaCl; o, 3 g bacto agar. All materials dissolved at the Erlenmeyer tube 100 ml with 20 ml aquadest, and then closed by cotton and aluminum foil. After that it sterilized by autoclave for 45 minutes at 121°C, 1 tam. Then waited until lukewarm and poured into Petri disk then let it until cold and hard or solid.

### Halo test

Bacteria isolation, *Actinobacillus* sp. planted in the solid media contained substrate Carboxyl Methyl Cellulose (CMC) by streak ose to the bacteria colony then streak again to the surface of solid media. Incubated at 37°C for 16 hours. Observation done to find Halo (transparent area) around the colony then choose the isolation had the biggest Halo and also shown the highest cellulase activities.

Then Halo gotten was examined by Congo-red 0.1% poured in to the isolation. Positive result signed with Halo area transparent still and the color became red (Puspaningsih, 2004).

### Inoculums production

Isolation renewed was moved to the growth media 20 ml in the Erlenmeyer tube 100 ml, and then incubated with shaking 150 rpm at 39°C during optimal cultivation time received from growth curve. Cultivation result was cell suspension or inoculums

### Enzyme production

Enzyme production accomplished by inoculated 1% inoculums in to 50 ml fluid media contained carboxyl methyl cellulose (CMC) in the Erlenmeyer tube 250 ml and incubated at 39°C for 16 hours with shaking 170 rpm. Incubation result taken the supernatant for tested the cellulase enzyme activities.

### The cellulase activities tested by method 3,5-dinitro-salicylic (DNS) acid

A number 100µl enzyme and xilan substrate entered to the each Eppendorf tube then incubated at 39°C in the water heater for an hour. For control, the formula was same with sample without incubation. Then sample and control added 600µl DNS, after that heated in the boiled water for 15 minutes, then incubated in the ice bath for 20 minutes. Absorbance read in the wave length 550nm. 1 unit enzyme activities defined as a total enzyme produced 1µmol xylosa per minute for every ml enzyme, and the specific activities was enzyme unit divided protein content (Miller, 19960).

### Characterization of Cellulose Enzyme

Determination of optimum temperature for cellulase done by determined activities of cellulase at many kind temperatures variant, that was at temperature: 35, 40, 45, 50, 55 and 60°C.

Determination of optimum temperature of cellulase determined by activity assay to cellulase at pH 5-9 used buffer phosphate citrate at optimum temperature of cellulase.

### Results and Discussion

The research result of enzymes cellulase activities from bacteria *Actinobacillus* sp. planted in the medium contained MCM at the many kind of temperature presented at Table 1 and Fig. 1. The influence of temperature on the cellulase activities at this result investigated at 35-60°C. Optimization temperature result of cellulase received at 45°C with activities 1.42 U/ml, but at 50°C and 55°C cellulase still have good activities that reached 92.96% (1.32 U/ml) and 78.17% (1.11 U/ml). At the temperature above 55°C decreasing of cellulase activities occurred, because atoms in the enzyme molecule had big enough energy to move, it caused by the presence form changing of enzyme structure because increasing thermal tremble of atoms component, therefore protein as enzyme material denaturized.

The research of optimum temperature was lower then the research result reported by Lowe *et al.*, (1997) that rumen fungi *Neocallimastix frontalis* plant at the wheat straw substrate produced cellulase had optimum activities at 50°C.

Cellulase activities realized increasing from pH 5 reached the highest activities about 0.57 U/ml at pH 6, at pH 7 cellulase activities decreased reached 73.68% (0.42 U/ml) from activities at the optimum pH. It's probably caused by conformation changing between enzyme and substrate because pH decreasing (Fig. 2).

Generally, in the cattle rumen the optimum pH was 6-7. According to the research result, gotten optimum pH from cellulase enzyme was suitable with condition of rumen pH. It was one of cellulase characteristic that was total ions at that pH made confirmation of cellulase formed a pair with the


substrate. It caused also by enzyme received from rumen bacteria, thus pH gotten was suitable with bacteria's habitat. Optimum condition at pH 6-7 was suitable to growth cellulitic and hemicellulitic bacteria in the rumen to degraded fibrous feed produced VFA. This result synchronizes with the opinion of Owens and Goetsch (1988) that pH condition very influenced by microbes environmental in the feed degradation. That cellulolytic and hemicellulolytic bacteria need pH about 6.2-7.0 to develop fast (Orskov and Ryle, 1990).

**Table 1.** The Data of Optimum Temperature of Cellulase of Bacteria *Actinobacillus* sp.

Temperature (°C)	Cellulase activities (U/ml)
35	0.86
40	0.71
45	1.42
50	1.32
55	1.11
60	0.96

**Table 2.** The Data of Cellulase Optimum pH of Bacteria *Actinobacillus* sp.

pH	Cellulase Activities (U/ml)
5	0.53
6	0.57
7	0.42
8	0.16
9	0.15



**Figure 1.** The pattern of relationship between temperature and the cellulase enzyme activities.




Figure 2. The pattern of relationship between pH and the cellulase enzyme activities.

### Conclusion

The characterization of cellulase enzyme has optimum temperature 45°C about 1.42 U/ml and optimum pH about 0.56 U/ml

### References

- Bhat MK and Hazlewood GP. 2001. Enzymology and Other Characteristics of Cellulases and Xylanases, In: *Enzymes in Farm Animal Nutrition* M.R. Bedford and G.G. Partridge (Eds). CABI Publ, UK.
- Beauchemin KA, Colombatto D, Morgavi DP and Yang WZ. 2003. Use of exogenous fibrolytic enzymes to improve feed utilization by ruminant J Anim Sci; 81: E37 – E47
- Bondi AA. 1987, Animal Nutrition, Wiley-Interscience Publication, London.
- Campbell Mary K. 1995. Biochemistry. 2<sup>nd</sup> Ed. Mount Holyoke College, Saunders College Publ, United States of America. Pp. 330-332
- Lowe SE, Michael K, Theodorou and Anthony PJT. 1987. Cellulases and Xylanases of an Anaerobic Rumen Fungus Grown on Wheat Straw, Wheat Straw Holocellulose, Cellulose, and Xylan. Appl Environ Microbiol.
- Mc. Donald P, Edwards RA and Greenhalgh JFD. 1995. Animal Nutrition. 3<sup>rd</sup> Ed. Logman, London and New York.
- Miller Gail Lorenz. 1960. Use of Dinitrosalicylic acid reagent for determination of reducing sugar. Anal Chem; 31: 426-428
- Orskov ER and Ryle. 1990. Energy Nutrition in Ruminant. Elsevier Application. Jhon Willey and Sons, New York.
- Owens FN and Goetch AL. 1988. Ruminant Fermentation In: D.C. Church (Ed), The Ruminant Animal Digestive Physiology and Nutrition. A Reston Book Prentice Hall. Englewood Cliffs, New Jersey.
- Ottaway JH and Apps DK. 1984. Biochemistry. 4<sup>th</sup> Ed. Cambridge, ELBS.
- Puspaningsih NNT. 2004. Isolation and characterization of  $\beta$ -Xylosidase and  $\alpha$ -L-Arabinofurosidase from *Bacillus thermoleovorans* IT-08. biotechnology of lignocellulose degradataion and biomass utiliozation. Mie University. Publ Co, Ltd.
- Spolestra SF, Steg A and Beuvink JMW. 1990. Application of cell wall degrading enzymes to grass silage. In: Agricultural biotechnology in focus in The Netherlands. J.J. Dekkers, H.C. van der Plas and D.H. Vuijk (Eds). Pudoc Wageningen.
- Sunna A and Antranikian G. 1997. Xylanolytic enzymes from fungi and bacteria. Graham G. Stewart and Inge Russel (Eds). Ctit Rev Biotech; 17(1): 39-67



# Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle

## ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

7%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.journal.unair.ac.id">www.journal.unair.ac.id</a> Internet Source	11%
2	<a href="http://lib.dr.iastate.edu">lib.dr.iastate.edu</a> Internet Source	1%
3	<a href="http://www.imedpub.com">www.imedpub.com</a> Internet Source	1%
4	<a href="http://ufdcimages.uflib.ufl.edu">ufdcimages.uflib.ufl.edu</a> Internet Source	1%
5	<a href="http://fspublishers.org">fspublishers.org</a> Internet Source	1%
6	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	1%
7	<a href="http://jddtonline.info">jddtonline.info</a> Internet Source	1%
8	M. Vatanparast, V. Hosseiniveh, M. Ghadamyari, S. Minoo Sajjadian. "Plant cell wall degrading enzymes, pectinase and cellulase, in	1%

the digestive system of the red palm weevil,  
Rhynchophorus ferrugineus (Coleoptera:  
Curculionidae)", Plant Protection Science, 2014

Publication

- 
- |   |   |    |
|---|---|----|
| 9 | <a href="http://ijlr.org">ijlr.org</a><br>Internet Source | 1% |
|---|---|----|
- 
- |    |   |    |
|----|---|----|
| 10 | <a href="http://unair.ac.id">unair.ac.id</a><br>Internet Source | 1% |
|----|---|----|
- 
- |    |  |     |
|----|--|-----|
| 11 | Yunus, Muchammad, and Endang Suprihati. "Potency of Attenuated Eimeria tenella in Protective Immunity Induction on Homologous and Heterologous Challenges", Procedia Chemistry, 2016.<br>Publication | <1% |
|----|--|-----|
- 
- |    |  |     |
|----|--|-----|
| 12 | Routes to Cellulosic Ethanol, 2011.<br>Publication | <1% |
|----|--|-----|
- 
- |    |                                      |     |
|----|--------------------------------------|-----|
| 13 | Submitted to iGroup<br>Student Paper | <1% |
|----|--------------------------------------|-----|
- 
- |    |   |     |
|----|---|-----|
| 14 | A.M.F. Milagres, P.O. Magalhães, A. Ferraz. "Purification and properties of a xylanase from cultivated on ", FEMS Microbiology Letters, 2005<br>Publication | <1% |
|----|---|-----|
- 
- |    |  |     |
|----|--|-----|
| 15 | Anuradha Jabasingh, S.. "Utilization of pretreated bagasse for the sustainable bioproduction of cellulase by Aspergillus | <1% |
|----|--|-----|

nidulans MTCC344 using response surface methodology", Industrial Crops & Products, 201111

Publication

16

[www.e-sciencecentral.org](http://www.e-sciencecentral.org)

Internet Source

<1%

17

[link.springer.com](http://link.springer.com)

Internet Source

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Optimization of Temperature and Cellulase pH from Rumen Bacteria Isolation of Beef Cattle

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---