

Effect of earthworm (*Lumbricus rubellus*) in feed formulation to improve fatty acids profile in eel (*Anguilla bicolor*) meat

by Widya Paramita

Submission date: 27-Nov-2019 12:12PM (UTC+0800)

Submission ID: 1222615128

File name: Bukti_C_44_Effect_of_earthworm_lumbricus....pdf (213.74K)

Word count: 2554

Character count: 11910

Effect of earthworm (*Lumbricus rubellus*) in feed formulation to improve fatty acids profile in eel (*Anguilla bicolor*) meat

K Farah¹, I R Gunawan¹, G B Putra¹, Agustono², W P Lokapirnasari⁴, M Lamid², E D Masithah³, T Nurhajati¹ and Rozi²

¹ Faculty of Fisheries and Marine, Airlangga University, Jl. Airlangga 60115, Surabaya, East Java, Indonesia.

² Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine Airlangga University, Jl. Airlangga 60115, Surabaya, East Java, Indonesia.

³ Departement of Marine, Faculty of Fisheries and Marine, Airlangga University, Jl. Airlangga 60115, Surabaya, East Java, Indonesia.

⁴ Department of Animal Science, Faculty of Veterinary Medicine Airlangga University, Jl. Airlangga 60115, Surabaya, East Java, Indonesia.

E-mail: dwiyp@fpk.unair.ac.id

Abstract. Eel requires unsaturated fatty acids of linolenic acid for growth. Which can be supplied from earthworms. In this study, addition of earthworm in formulation feed aimed to improve the fatty acid profile eel meat. This research used experimental method and randomized complete design method with five treatments. Each treatment was repeated four times. The use of earthworms in feeding treatment formulation was done for 21 days with different level i.e: 0 % (P0), 25 % (P1), 50 % (P2), 75 % (P3) and 100 % (P4). The result showed that the addition of earthworm significantly influenced the omega 3 contents (EPA & DHA) of eel meat.

1. Introduction

Eel (*Anguilla bicolor*) is a freshwater fish that has a large potential to be developed. According to Hameed *et al.* [1], eel contains 48.430 % saturated fatty acids, 50.639% unsaturated fatty acids. In addition, there is a 0.461 % EPA, 1.294 % DHA, 9.134 % linoleic acid and 0.472 % arachidonic acid. Eel contains 1.337 mg/100 g DHA and 742mg/100g EPA. Consumers' demand for eels has increased due to its high nutrient content. Demand for eels in international markets has reached 300,000 tons/year. The market demand for eels is increasing because people consider the eel is savory and rich beneficial for the body [2]. Eels are known as a fishery commodity that is rich in protein, fat, minerals, and vitamins compared to other fish species [3].

Polyunsaturated fatty acids functions as a nutrient in the body, such as EPA and DHA that have benefits to human health. EPA and DHA contained in fatty fish and they cannot be synthesized in the human body [4]. The ratio between omega-3 and omega-6 fatty acids is a good indicator for comparing the relative nutritional value of different species of fish. The ratio of omega-3 and omega-6 fatty acids helps to prevent coronary heart disease by reducing the levels of plasma lipids and risk of cancers [5]. Omega-3 and omega-6 fatty acids are polyunsaturated and are important components of cell membranes and are precursors to many other substances in the body such as those involved with regulating blood

pressure and inflammatory responses, thus they must be obtained through food [6]. Therefore, this research was aimed to know effect of addition of earthworms (*Lumbricus rubellus*) in feed formula on fatty acid profile of eels.

2. Methodology

2.1 Preparation of earthworm (*Lumbricus rubellus*)

Earthworms were obtained from Malang, East Java. They were mixed with feed formulation (commercial fishmeal) in doses of 0, 25, 50, 75, and 100 % and shaped like pasta. The feed then went through proximate analysis as shown in table 1.

Table 1. Nutritional feed ingredients.

Feed Ingredients	Dry Ingredients	Crude Protein (%)	Crude Fat (%)	Crude Fiber (%)	Ash (%)	NFE (%)
Formulation Feed	80.029	42.067	13.011	6.806	14.228	3.916
Earthworms	22.896	13.634	5.749	0.548	1.589	1.376

2.2 Experimental design

Eels used were from fingerling stage weighing 20-25 g with the total of 100 g and they were supplied from Malang, East Java. Eels were selected and then acclimatized for 30 min. Furthermore, eels were adapted for a week. For 21 days, the eels were given feed containing 0 %, 25 %, 50 %, 75 %, and 100 % earthworm.

2.3 Fish composition

The levels of fat, protein, fiber, and energy ingredients without nitrogen (BETN) in eel meat were analyzed based on AOAL methods [7].

2.4 Fatty acid derivatisation

Meat was cut, chopped, weighed for 1 g, and put into test tubes. Sodium chloride (0.5 g) and 4 mL hexane were added and the mixture was vortexed for 2 min until it was clear. Clear hexane layer was taken and transferred into the next derivatisation tube and drained with stream of nitrogen. NaOH 2 % (2 mL) was added to methanol then close to temperate at 90 °C for 5 min. The result was left to cool before added with 2 mL methanol in BF3 further heated again for 30 min. Samples were then extracted with 3 mL of hexane to final stage. Extract was analyzed by GC-MS [8].

2.5 Gas chromatography (GC)

Samples were analyzed using gas chromatography Shimadzu GC-2014 with helium as a carrier gas and SGE forte BPX 70 column (film thickness of 30 m x 0.25 mm ID x 0.25 µm) (SGE Europe Ltd. Milton Keynes, UK) as the analytical column. The peaks were identified using standard mix of 38 external FAME (FAME Mix C4-C24, Supelco; Sigma – Aldrich). Initial column temperature was set at 50 °C for 1 min. Temperature was raised at 2 °C/min until it reached 188 °C and maintained for 10 min. next, the temperature was increased further to 240 °C and maintained for 4 min before it was returned to the initial temperature [9].

2.6 Statistical analysis

The data were expressed as mean ± standard deviation. The data were analyzed using SPSS version 16.0 (SPSS Inc., Chicago, IL). Differences between means were analyzed by Analysis of Variance followed by Duncan's multiple comparison test. Significant difference was set at $p < 0.05$.

3. Results

3.1 Proximate analysis of eel meat

ANOVA showed that there were no significant effect of different feed formulations ($p > 0.05$) (Figure 1) on nutritional content of eel meat. The use of earthworms led to increasing levels of fat on eel meat with 1.612 % in P0, 1.242 % in P1, 1.256 % in P2, 1.690 % in P3, and 1.505 % in P4. Protein levels in eel meat were 17.073 % in P0, 16.957 % in P1, 16.599 % in P2, 17.413 % in P3, and 17.846 % in P4. Meanwhile, ash content ranged from 0.083 % to 1.148 % and the energy content was within the range of 0.95 to 1.014 %.




Figure 1. Proximate composition of eel meat as affected by different feed formula added with earthworms on the feed formulation against fish eel meat content.

3.2 Fatty acid profile

There were no significant effect given by the different feed formula on saturated fatty acids, unsaturated fatty acids, and omega 6 fatty acids of eel meat. Meanwhile, the treatments gave significant differences in EPA and DHA content of eel meat, where P4 had significantly higher EPA and DHA contents than P0 (table 1).

Table 2. Fatty acid profile of eel meat.

4	FAMES	Treatment (%)				
		P0	P1	P2	P3	P4
	C12:0 Lauric acid	0.347	1.003	0.624	0.397	0.830
	C14:0 Myristic acid	5.291	5.417	5.507	4.942	5.465
	C13:0 Pentadecanoic acid	0.665	0.696	0.629	0.701	0.739
	C16:0 Palmitic acid	25.347	25.057	25.410	24.529	25.164
	C18:0 Stearic acid	5.317	5.471	5.010	5.834	5.552
	Σ Saturated Fatty acid (SFA)	36.967	37.644	37.180	36.403	37.750
	C16:1 Palmitoleic acid	6.219	6.217	6.445	5.699	6.070
	C18:1 Oleic acid (ω9)	32.478	31.880	32.177	32.193	31.859
	C18:1 Elaidic acid	3.894	4.179	3.978	3.898	4.099
	C20:1 cis 11 Eicosenoic acid	2.268	2.648	2.521	3.403	2.638

C22:1 Erucic acid	0.293	0.256	0.372	0.458	0.273
∑Mono Unsaturated Fatty Acid (MUFA)	45.152	45.180	45.493	45.651	44.939
16 8:2 Linoleic acid (LA) (ω6)	8.139	8.946	8.593	9.1635	8.661
C20:4 Arachidonic acid (ω6)	1.432	1.481	1.314	1.488	1.392
C20:5 Eicosapentaenoic acid (EPA) (ω3)	1.243 ^a	1.014 ^c	1.194 ^{ab}	1.101 ^{abc}	1.073 ^{bc}
C20: 3 Cis 8, 11, 14 Eicosatrienoic acid (hGLA) (ω6)	0.712	0.771	0.673	0.792	0.749
C18:3 α Linolenic (ω3)	0.600	0.541	0.577	0.600	0.543
C22:6 Docosahexaenoic (DHA) (ω3)	4.599 ^a	3.257 ^b	4.045 ^{ab}	3.655 ^{ab}	3.790 ^{ab}
∑Poly Unsaturated Fatty Acid (PUFA)	16.725	16.010	16.396	16.799	16.208
∑Unsaturated Fatty Acid (UFA)	61.877	61.190	61.889	62.450	61.147
SFA/UFA	0.597	0.615	0.600	0.582	0.617
UFA/SFA	1.673	1.625	1.664	1.715	1.619
∑n3	6.442	4.812	5.816	5.356	5.406
∑n6	10.283	11.198	10.580	11.443	10.802
n3/n6	0.626	0.429	0.549	0.468	0.500
n6/n3	1.596	2.327	1.819	2.136	1.998
EPA	1.243 ^a	1.014 ^c	1.194 ^{ab}	1.101 ^{abc}	1.073 ^{bc}
DHA	4.599 ^a	3.257 ^b	4.045 ^{ab}	3.655 ^{ab}	3.790 ^{ab}
EPA/DHA	0.270	0.311	0.295	0.301	0.283

P0 = commercial feed and earthworm (100%: 0%). P1 = commercial feed and earthworm (75%: 25%). P2: commercial feed and earthworm (50%: 50%). P3 = commercial feed and earthworms (25%: 75%). P4 = commercial feed and earthworm (0%: 100%)

4. Discussion

Figure 1 is in accordance with the results of Litzow *et al.* [10] who stated that the fat content in fish feed should be about 15%. Fat content in eel meat is highly correlated with the content of essential fatty acids. Moreover, Kandemir and Polat [11] stated that the content of fatty acids in aquatic organisms can be influenced by the living condition, either wild in nature or in captivity. There was lack of linolenic acid found in feed formula although the fatty acid was found in the earthworms.

The fatty acid profile of eel meat as shown Table 2 is in accordance with Oku *et al.* [12] who reported fatty acid content of Japanese eel (*Anguilla japonica*) fresh meat consisted mostly of monounsaturated fatty acids (MUFA), while unsaturated fatty acids (PUFA) appeared in low amount. Variation of fatty acids in aquatic organisms can be influenced by seasons, geographical location, and environment salinity [13].

Different doses of earthworms in eel feed formula could increase EPA and DHA contents in eel meat. The content of the omega-3 fatty acids EPA and DHA was affected by the presence or absence of earthworms in the feed formulation. According to Robin *et al.* [14] stated that when feed is rich in omega-3 fatty, then the fish meat composition would be influenced. This is in accordance with the results of Huang *et al.* [15] stating that fatty acids contained in fish meat is derived from the fatty acids consumed by the fish.

Omega 3 and omega 6 fatty acids are polyunsaturated fatty acids (PUFA). Omega 6 in eel meat showed a higher percentage compared with omega 3 [1]. Extremely high Omega 6 can negatively affect the body. The number and ratio of omega 3 and omega 6 fatty acids are important to be considered in formulating fish feed. A good ratio of omega 3: omega 6 is 10:1, which means there should be higher omega 3 content compared with omega 6 [3]. The best ratio was found in treatment P1 (75 %: 25 %) that was 1: 1.8. The composition of fatty acids in feed formulation can affect the ratio of omega 3 and omega 6 [16].

Fatty acids n-3 and n-6 are required in fat biosynthesis, so that in the event of a shortage or excess of one of the fatty acids, it will inhibit the rate of biosynthesis of other fatty acids and eventually it will

24 affect the composition of fatty acids in fish. an imbalance 5 ratio of omega 3 and omega 6 can lead to competition in utilizing enzymes in fat metabolism, which can affect growth. As a conclusion, the use of earthworms in eel feed did not significantly affect fatty acid profile in eel meat.

5. Conclusions

Additions of earthworm in eel feed formula increase EPA and DHA contents in eel meats. The best EPA and DHA ratio of commercial feed and earthworm was in treatment P1 (75%: 25%).

6. References

- [1]. Hameed A S, Hussain, Shabbir, Pasha and Song 2017 *Am. J. of Biochem. & Biotech.* **13** 15-26
- [2]. Ozogul F and Ozogul Y 2006 *Food Chem.* **99** 574-578
- [3]. Seo J S, Choi J H, Seo J H, Ahn T H, Chong W S, Kim S H, Cho H S and Ahn J C 2013 *Fish Aquat. Sci.* **16** 85-92
- [4]. Kolanowski W and Laufenberg G 2006 *Eur Food Res Technol.* **222** 472-477
- [5]. Simopoulos A. P 2002 *Biomed. Pharmacother.* **8** 36-79
- [6]. Escott-Stump S and Mahan L K 2005 Krause, alimentos, nutricao and dietoterapia *Editoraroca* **9**
- [7]. AOAC 2005 Official methods of analysis (18th edition) association of official analytical *Chemists International* maryland USA
- [8]. Amer B, Caroline N, Hanne C B, Grith M, Kjeld H and Trine K D 2013 *Elsevier* **32** 199-203
- [9]. Farhat J and Shakoor C 2011 *J. Food Chem.* **125** 991-996
- [10]. Litzow M A, Bailey K M, Fredrick G and Prahl H R 2006 *Mar. Ecol. Prog. Ser.* **315** 1-11
- [11]. Kandemir S and Polat 2007 *Turkish J. Fish Aquatic Sci.* **7** 27-31
- [12]. Oku T, Sugawara, Choundhury, Komatsu, Yamada and Ando 2009 *Food Chem.* **115** 436-440.
- [13]. Ozyurt G O, Duysak, Akameca and Tureli 2006 *Food Chem.* **95** 382-385
- [14]. Robin J H, Regost, Arzel and Kaulshik 2003 Fatty acid profile following a change in dietary fatty acid source model of fatty acid composition with a dilution hypothesis **225** 10-13.
- [15]. Huang W Y, Wu J T, Chiang Y R and Jane W N 2006 *Aquat. Toxicol.* **80** 38-45
- [16]. Bae J Y, Han, Park and Bai 2004 *J. Aquacult.* **17** 275-281

Effect of earthworm (*Lumbricus rubellus*) in feed formulation to improve fatty acids profile in eel (*Anguilla bicolor*) meat

ORIGINALITY REPORT

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

12%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 www.er-c.org 2%
Internet Source
- 2 Farhat Jabeen, Abdul Shakoor Chaudhry. "Chemical compositions and fatty acid profiles of three freshwater fish species", Food Chemistry, 2011 2%
Publication
- 3 mulpix.com 1%
Internet Source
- 4 res.mdpi.com 1%
Internet Source
- 5 www.mypetshop.co.nz 1%
Internet Source
- 6 Jabeen, F.. "Chemical compositions and fatty acid profiles of three freshwater fish species", Food Chemistry, 20110401 1%
Publication
- 7 William J Aronson, John A Glaspy, Srinivasa T

Reddy, David Reese, David Heber, Dilprit Bagga. "Modulation of omega-3/omega-6 polyunsaturated ratios with dietary fish oils in men with prostate cancer", Urology, 2001

Publication

1%

8

Feriel Ghribi, Dhouha Boussoufa, Fatma Aouini, Safa Bejaoui, Imene Chetoui, Imen Rabeh, M'hamed El Cafsi. " Seasonal variation of biochemical composition of Noah's ark shells (L. 1758) in a Tunisian coastal lagoon in relation to its reproductive cycle and environmental conditions ", Aquatic Living Resources, 2018

Publication

1%

9

www.tandfonline.com

Internet Source

1%

10

www.ajas.info

Internet Source

1%

11

"Lipids in Aquatic Ecosystems", Springer Nature, 2009

Publication

1%

12

Jae-Sung Seo, Jae-Hee Choi, Ji-Hun Seo, Tae-Ho Ahn, Won-Seog Chong, Seung-Hoon Kim, Hye-Sun Cho, Jun-Cheul Ahn. "Comparison of Major Nutrients in Eels *Anguilla japonica* Cultured with Different Formula Feeds or at Different Farms", Fisheries and aquatic sciences, 2013

1%

13	pdfs.semanticscholar.org Internet Source	<1 %
14	springerplus.springeropen.com Internet Source	<1 %
15	www.pubmedcentral.nih.gov Internet Source	<1 %
16	Wiesenfeld, P.. "Flaxseed increased @a-linolenic and eicosapentaenoic acid and decreased arachidonic acid in serum and tissues of rat dams and offspring", Food and Chemical Toxicology, 200306 Publication	<1 %
17	www.naturecures.co.uk Internet Source	<1 %
18	albertahorseindustry.ca Internet Source	<1 %
19	www.chekyang.com Internet Source	<1 %
20	journals.plos.org Internet Source	<1 %
21	www.agilent.com Internet Source	<1 %
22	content.sciendo.com Internet Source	<1 %

23

www.sciencedev.net

Internet Source

<1%

24

Emi Suzuki. "Alteration in the contents of unsaturated fatty acids in dnaA mutants of Escherichia coli : Fatty acid composition in dnaA mutants", Molecular Microbiology, 05/01/2002

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On