

PROCEEDING

The International Conference on
**Pharmacy and Advanced
Pharmaceutical
Sciences**



Faculty of Pharmacy UGM
Yogyakarta Indonesia
October 2009



Pharmacy and Advanced Pharmaceutical Sciences
Faculty of Pharmacy UGM



Organizing Committee

Steering committee

Prof. Dr. Marchaban, DESS, Apt (Pharmacy, UGM, INDONESIA)

Prof. Dr. Umar Anggara Jenie, M.Sc., Apt (LIPI, INDONESIA)

Prof. Dr. Retno S. Sudiby, M.Sc., Apt. (Pharmacy, UGM, INDONESIA)

Prof. Dr. Syed Azhar S. Sulaiman (USM, MALAYSIA)

Prof. Masashi Kawaichi, Ph.D (NAIST JAPAN)

Dr. Edy Meiyanto, M.Si., Apt (CCRC, UGM INDONESIA)

Prof. Dr. Zullies Ikawati, Apt (Pharmacy, UGM, INDONESIA)

Chairman

Dr. Hilda Ismail, M.Si., Apt.

Secretary

Dr. rer. nat. Ronny Martien, M.Si

Treasurer

Dr. Ratna Asmah S., M.Si., Apt

Scientific Committee

Prof. Dr. Sismindari, SU., Apt.

Prof. Dr. Lukman Hakim, M.Sc., Apt

Prof. Dr. Suwidjijo Pramono, Apt.

Prof. Dr. Suwaldi, M.Sc., Apt

CONTENTS

Preface from the Editor	i
Organizing Committee	ii
Welcome Message Proceeding International Conference on Pharmacy and Advanced Pharmaceutical Sciences	
From the committee	iii
Remark of the Dean Faculty	v
Senior Vice Rector For Education	v
CONTENT	vi
Pharmacogenetics : in case of cytochrome P450 oxidases (CYPS) related to adverse drug reactions Arum Pratiwi, Harianto Lim and Ronny Martien	1 – 4
Interaction of turmeric and garlic extract combination against free radical scavenging activity Patonah, Daryono H. Tjahjono, Elin Yulinah S. and I Ketut Adnyana	5 – 6
Influenced of Kojic Acid and B-Cyclodextrin on SPF Value Sunscreen Product Contained Oxybenzone and Octyl Dimetyl Paba (3:7) (In vanishing cream base formulation) Diana, Tristiana Erawati, Widji Soeratri and Noorma Rosita	7 – 14
Isolation and Antimicrobial activity of endophytic fungi <i>Kabatiella caulivora</i> var B isolated from <i>Alyxia reinwardtii</i> BL Noor Erma Sugijanto, Dian Anggraeny and Noor Cholies Zaini	15 – 17
Rapid and Simple Luciferase Reporter Gene Assays for the Discovery of Peroxisome Proliferator-Activated Receptor α and γ Agonists and Nuclear Factor- κ B Inhibitors from Medicinal Plants. N. Fakhruddin, S. Vogl, P. Picker, E. H. Heiss, J. Saukel, G. Reznicek, B. Kopp, A. G. Atanasov and V. M. Dirsch	18 – 24
Identification of components of essential oil from <i>Cananga odorata</i> which penetrated into the rat skin / (wistar strain) in the practice of <i>Timung</i> (development of <i>Timung</i> as alternative healing) Mangestuti Agil, Esti Hendradi and Budiastuti	25 – 29
In Vivo Antihyperglycemic Test of Albedo Durian (<i>Durio zibethinus</i> M) Extract on Aloxan- Induced Diabetic White Rat (<i>Rattus norvegicus</i>) F. M. Cahyani, I. Susanti, R. Ratna, Y. D. Panggi and Y. Pravitasari	30 – 33
Effect of Pasak Bumi's Root (<i>Eurycoma longifolia</i> , Jack) on Sperm Output in Rats Farida Hayati and Mustofa	34 – 37
<i>Proceeding International Conference on Pharmacy and Advanced Pharmaceutical Sciences Yogyakarta, Indonesia, 2009</i>	xi

The Influence of Arbutin 3% and Sesame Oil (3,5,7 % w/w) on SPF Values of Oxybenzon and Padimate O (3:7% w/w) in carbomer Gel Base Noorma Rosita, Tristiana Erawati and Rafi Jikrona	38 – 43
Sulochrin as α -glucosidase inhibitor <i>lead compound</i> Rizna Triana Dewi, Ahmad Darmawan, Sofna D.S Banjarnahor, Hani Mulyani, Marisa Angelina and Minarti	44 – 48
The Practice of Complementary Indigenous Malay Therapies In Rural Areas: Do Users' Attitudes, Beliefs And Perceptions Significantly Differ From Non-Users? Pei Lin Lua, Rohayu Izanwati Mohd Rawi, Suffian Mohamad Tajudin, Norlida Mamat and Ahmad Zubaidi Abdul Latif	49 – 54
An Interventional Pilot Study: Effect Of Dark Chocolate Consumption On Anxiety Level Among Female Nursing Students Sok Yee Wong, Pei Lin Lua, Rohayu Izanwati Mohd Rawi, Rokiah Awang, Ahmad Zubaidi and Abdul Latif	55 – 62
The Anti-proliferation Assay of Bioactive Fraction from <i>Curcuma zedoaria</i> Rhizome Ros Sumarny, Priyosoeryanto B. P., Ietje W., Latifah K. D. and Chairul	63 – 67
Studies of Sub-acute Toxicity Assay from <i>Acorus calamus</i> L. in Experimental Animal Models Banjarnahor S.D.S, Sri Hartati and Megawati	68 – 71
Antioxidant Properties and Phenolics Content of <i>Mikania scandens</i> L.(Wild) Sumi Wijaya, Ting Kang Nee, Khoo Teng Jin and Christophe Wiert	72 – 77
The Influence of Olive Oil Addition on Increasing of Arbutin Penetration in the Carbomer-940's Gel (Observation on Inhibition of Enzyme Tyrosinase Activity) Widji Soeratri, Tristiana Erawati, Noorma Rosita and Fahriyatul Wahyuni	78 – 81
The difference of antioxidant activity of various tea (<i>Camellia sinensis</i> L.) methanol extract Wahyu Widowati, Tati Herlina and Hana Ratnawati	82 – 88
Chemical Stability of Cisplatin and Ondansetron During Simulation of hemotherapy Administration Yahdiana Harahap, Rizka Andalusia and Armon Fernando	89 – 94
The Effects of Cassava Starch (<i>Manihot utilissima</i> , Pohl.) as a Binder on Physicochemical Characteristics of Acetaminophen Tablet Formulation Yandi Syukri, Tri Rahayu Ningsih and M. Hatta Prabawa	95 – 98
Drug Interaction Study in Hospitalized Hepatic Cirrhosis Patient in Dr. Ramelan Navy Hospital Amelia Lorensia, Aziz Hubeis, Widyati and Hary Bagijo	99 – 102
The Effect of Cold Storage in Krebs-Henseleit Buffer in the Viability and Metabolic Activities of Precision Cut Intestinal Slices Dewi Setyaningsih, AA Khan and GMM Groothuis	103 – 110

Influence of Tween 80 Concentration in Carbomer/ Tween 80 Aggregate on Kojic Acid Penetration (Observed on Inhibiting Tyrosinase Activity in Vanishing Cream) Siti Evi Jayanti, Tristiana Erawati and Noorma Rosita	186 – 191
Validation for Result Degradation of Nifedipine Residue with Thin Layer Chromatography-Densitometry and Thin Layer Chromatography-Spectrophotometry Sitti faika and Sudibyo Martono	192 – 195
Synthesis and Biological Activity Test of Antibiotic UK-3 Analogues, 2-Hydroxynicotinyl-Butyl-Serine-Ester and Its Derivatives Ade Arsianti, Kiyomi Kakiuchi, Tsumoru Morimoto, M.Hanafi and Endang Saefudin	196 – 198
Vitamin e content in the dragon fruit Established by high performance thin layer chromatography–densitometry Any Guntarti and Warsi	199 – 204
Drug interaction study in hospitalized hepatic cirrhosis patient in Dr. Ramelan navy hospital Amelia Lorensia, Aziz Hubeis, Widyati and Hary Bagijo	205 – 208
PGV-1 inhibits G2M phase progression in WIDr colon cancer cell Endah Puji Septisetyani, Edy Meiyanto, Masashi Kawaichi and Muthi' Ikawati	209 – 212
Proceeding International Conference on Pharmacy and Advanced Pharmaceutical Sciences The influence of oleic acid pre-treatment on transport of epigallocatechin gallat in green tea (<i>Camellia sinensis</i> , L) extract Across mice skin in vitro Nining Sugihartini, Achmad Fudholi, Suwidjiyo Pramono and Sismindari	213 – 215
Development and Production of Anti Tuberculosis Fixed Dose Combinations (FDCs) Barokah Sri Utami, Syamsul Huda, Nurliya Irfiani and Badrus S.	216 – 218
The Characteristics and Release of Diclofenac Sodium of Niosome System in Carbomer 940 Gel Base Preparation (Niosome System of Diclofenac Sodium-Span 60-Cholesterol with Molar Ratio 1:5 :5) Esti Hendradi, Tutiek Purwanti, Bety Nurfia Puspitarini and Bianda Ida Kurnia	219 – 224
The Characteristics and Release of Diclofenac Sodium of Niosome System in Carbomer 940 Gel Base Preparation (Niosome System of Diclofenac Sodium-Span 20-Cholesterol with Molar Ratio 1:5 :5) Esti Hendradi, Tutiek Purwanti, Anditasari and Srimaryati	225 – 231
An Interventional Pilot Study: Effect Of Dark Chocolate Consumption On Anxiety Level Among Female Nursing Students Sok Yee Wong, Pei Lin Lua, Rohayu IZanwati Mohd Rawi, Rokiah Awang and Ahmad Zubaidi Abdul Latif	232 – 239
Antiemetics utilization in cancer patients with high emetogenic cytotoxic drugs in two govermental hospital in indonesia Dyah Aryani Perwitasari and Ana Hidayati	240 – 243
KEY WORDS INDEX	244
DISCUSSION	246

The Effect Of β -Cyclodextrin And Oxybenzone-Octyl Dimethyl Paba (3:7% W/W) Addition On The Penetration Of Kojic Acid In Vanishing Cream (Based on Activity Inhibition of Tyrosinase) Diana Winarita, Tristiana Erawati, Noorma Rosita and Widji Soeratri	111 – 116
The profile of knowledge and self-medication in handling cough symptoms by students of pharmacy at Airlangga university Elida Zairina, Liza Pristianty and Lestriana Kusumasari	117 – 120
The Characteristics and Release of Diclofenac Sodium of Niosome System in Carbomer 940 Gel Base Preparation (Niosome System of Diclofenac Sodium-Span 60-Cholesterol with Molar Ratio 1:5 :5) Esti Hendradi, Tutiek Purwanti, Bety Nurfia Puspitarini and Bianda Ida Kurnia	121 – 128
Red Betel Vine (<i>Piper Crocatum</i>) Essential Oil as Antituberculosis Farida Juliantina Rachmawaty	128 – 133
Effect of Pasak Bumi's Root (<i>Eurycoma longifolia</i> , Jack) on Sperm Output in Rats Farida Hayati and Mustofa	134 – 137
The Influence of Sesame Oil Addition on Arbutin Release from Carbomer-940 Gel Bases Hanifa Rahma, Tristiana Erawati and Noorma Rosita	138 – 141
Phytochemical Screening and Determination of Antioxidant Activity of Fractions from Ethyl Acetate Extract of <i>Phyllanthus acidus</i> (L.) Skeels Leaf Hindra Rahmawati, Hesty Utami and Moordiani	142 – 145
Study on Antihyperglycaemic Activity of Ethyl Acetate Extract of Sidaguri (<i>Sida rhombifolia</i> L.) Stem on Alloxan-Induced Diabetic Mice (<i>Mus musculus</i> L.) Irma Ratna K, Muktiningsih, Suhartono, Natalia Elisabeht and Muhammad Ali Zulfikar	146 – 152
The Influence of Arbutin and Olive Oil as an Enhancer in Characteristic and SPF Value of Sunscreen (Combination of Oxybenzone and Octyldimethyl Paba in <i>Carbomer</i> 940 Gel Base) Josephine Paramita Ayuningtyas, Tristiana Erawati, Noorma Rosita and Widji Soeratri	153 – 160
The Effect of Secondary Emollients Triethylhexanoate, Isopropyl myristate, and Propyleneglycol Isostearate on in-vitro skin penetration of tocopheryl acetate cream using Franz-diffusion cell Joshita Djajadisastra, Sutriyo and Fraida Aryani	161 – 165
Immunomodulatory activity of <i>Plantago major</i> L. on IgM titer of mice Kartini, A. Kirtishanti, Dessy, Fauziah and Isnaini	166 – 169
Antibacterial activities of <i>Aleurites moluccana</i> (Euphorbiaceae) Othman Abd Samah and Rasyidah Mohamad Razar	170 – 178
Total synthesis and revised structure of benzophenone glucopyranosides from <i>Phaleria macrocarpa</i> Phebe Hendra, Yukiharu Fukushi and Yasuyuki Hashidoko	179 – 185

The Influence of Arbutin 3% and Sesame Oil (3,5,7 % w/w) on SPF Values of Oxybenzone and Padimate O (3:7% w/w) in carbomer Gel Base

Noorma Rosita, Tristiana Erawati, Rafi Jikrona

Pharmaceutics Department, Faculty of Pharmacy Airlangga University, Surabaya Indonesia

Email: itanr@yahoo.com

Abstract

This research was intended to know the influence of addition 3% Arbutin and (3; 5; 7)% w/w sesame oil on SPF value of sunscreen combination Oxybenzone and Padimate - O (3:7 % w/w) was formulated in carbomer gel base. Characteristic test was done in 2nd day after formulated included organoleptic, pH, and spread-ability. The sunscreen effectiveness determination (SPF value) was performed in vitro by measuring the absorbencies of its 10 ppm solutions in isopropanol between 290 - 400 nm. Organoleptic characteristic all formulas were different with the base, so did the pH and spread ability result. From SPF value determination results conclude that addition of arbutin increased SPF value, so did the increased of sesame oil concentration. Formula with the higher concentration of sesame oil had a higher SPF values. From this research, there was suggested to evaluate in vivo acceptability test and affectivity in vivo trial test on human skin.

Key words: Arbutin, Sesame oil, sunscreens, carbomer gel, Oxybenzone, Padimate - O, SPF values

Introduction

Sunscreens are mostly used to protect the skin from harmful effects of solar radiation. The active ingredients of sunscreens are divided into physical and chemical sunscreen agents. Physical sunscreen agents reflect the UV rays from our skin. It is not transparent and has to be used in a high concentration to be effective, which made them not preferable to some people. The chemical sunscreen agents work by absorbing UV-A radiation like oxybenzone and absorbing UV-B radiation such as Padimate-O (octyldimethyl PABA). To obtain a higher protection and producing a broad spectrum sunscreen, a combination of anti UV-A and anti UV-B sunscreen agents has been used nowadays in many sunscreen product [Widianingsih and Lumintang, 2002].

Normally, our skin has its own protection mechanism against the harmful effect of UV rays such as the thickening of *stratum corneum*, sweat and skin pigmentation. Increasing abnormality of melanin as a result of our skins natural protection can result in a non homogenous skin color which is usually reduce likeliness amongst people. To overcome this problem, whitening agents are used to control the production and metabolism of melanin in the epidermis. One example of a frequently used whitening agent is arbutin, a hydroquinone derivatives which inhibits melanin production. Arbutin has lower toxicity than hydroquinone's and its depigmentation effect is higher than kojic acid and vitamin C [Mashhood, 2006]. This substance is used as whitening agent at a various concentration ranging from 0.5-3.0%. It has a low partition coefficient and penetration rate. Therefore, it needs addition of a penetration enhancer [Zulkarnain, 2003; Galilee, 2008; Mitsui, 1998].

One of enhancers that frequently used in cosmetic formula is sesame oil. Sesame oil is a vegetable oil that obtained from pressing *Sesamum Indicum* seed. Mechanism of sesame enhanced polar pathway in stratum corneum because it contains protein (25%) dan globulin (67,3%) as main substance. Another mechanism is to enhance the oclusiveness (Sarma, K., 1993) to facilitate arbutin penetration. Sesame oil has been widely used in sunscreen and other cosmoceutics preparation for its emollient activity; it is nonirritant, considered as a natural lipid that has the highest compatibility with human skin and has sunburn healing effect [O'Neil, 2006].

Among several variant sunscreen preparations, gel gives us a cool sensation, not sticky, elegant, and smooth and easy to be washed from the skin. A synthetic gelling agent

like carbomer 940 usually requires only a small amount of them to produce a gel with good consistency compared to other types of gelling agent.

The aims of this research wants to know whether addition of arbutin 3% w/w and various concentration of sesame oil (3, 5, and 7% w/w) as arbutin's enhancer affect SPF value of sunscreen combination of oxybenzone 3% w/w (as anti UV-A) and Padimate O 7% w/w (as anti UV-B) which is formulated in carbomer 940 gel base. SPF values are observed by using a spectrophotometric method.

Methodology

Preparation of sunscreen gel containing arbutin and sesame oil

Oxybenzone, Padimate-O and propylene glycol were mixed and put into the good emulsion system contains arbutin, water, BHT, sesame oil, and tween 80 as emulsifying agent. After that it poured into the carbomer gel base and stirred well to form the sunscreen gel. The sunscreen were then kept in a tight container and stored well for further analysis. Preparation of each formula is described as below table.

Table 1: Formula used in research

Composition	Function	Concentration in formula (% w/w)				
		S	S+A	S+A+O 3%	S+A+O 5%	S+A+O 7%
Oxybenzone	Sunscreen agent	3	3	3	3	3
Padimate O	Sunscreen agent	7	7	7	7	7
Arbutin	Whitening agent	0	3	3	3	3
Tween 80	Emulsifying agent	0.5	0.5	0.5	0.5	0.5
Sesame oil	Enhancer	0	0	3	5	7
Carbomer	Gelling agent	1	1	1	1	1
Distilled water up to	Solvent	100	100	100	100	100

* Formula I = S = Sunscreen

Formula II = S+A = Sunscreen + Arbutin

Formula III = S+A+O 3% = Sunscreen + Arbutin + Sesame oil 3%

Formula IV = S+A+O 5% = Sunscreen + Arbutin + Sesame oil 5%

Formula V = S+A+O 7% = Sunscreen + Arbutin + Sesame oil 7%

Characteristics determination of sunscreens gel

The characteristic of preparation were determined include: organoleptic test visually, while the determination of pH and spread ability are done in 2 days after the formula were made by using a digital pH meter Schott CG 842, and a spread-ability apparatus.

Determination of SPF value of sunscreens gel

SPF value was determined by Petro's method.

Firstly, 100.0 mg sunscreens were dissolved in 2.0 ml isopropanol, the solution was then centrifuged for 15 minutes at 50 rpm. 1.0 ml of the filtrate is taken and poured into a 5.0 ml metered flask and shake well until its homogenized (10000 ppm). This solution diluted it and then its shaken well until it reached a concentration of 100 ppm (contains 10 ppm sunscreen's active ingredients). An UV spectrum of this solution was measured at 290-400 nm using Double Beam UV-Vis Spectrophotometer (Perkin Elmer Lambda EZ 201) at an interval of 2 nm which has an absorption that is larger than 0.050.

The AUC of each formula from the shortest and longest wavelength are calculated using the following equation:

$$AUC_{\lambda_{p-\alpha}}^{\lambda_p} = \frac{A_{p-\alpha} + A_p}{2} (\lambda_p - \lambda_{p-\alpha})$$

Whereas:

AUC = Area under Curve
 Ap = Absorbtion on p wavelength
 Ap-α = Absorbtion on p-α wavelength

The SPF values of a formula were obtained by converting log SPF calculating from the total AUC into the equation below:

$$\text{Log SPF} = \frac{\text{Total area}}{\lambda_n - \lambda_1} \times 2$$

Whereas:

λn= longest wavelength above 290 nm that has an absorption value higher than 0.050
 λ1= shortest wavelength 290 nm

Statistical analysis

One-way ANOVA were used to assess the significant of differences only for preparation with the same sunscreen category.

Results and Discussions

The average data of organoleptic observation from all sunscreen formula were shown in table 2. From the data we could see that the addition either of arbutin or sesame oil to the formula affect the color and smell.

Table 2: Organoleptic analysis of the sunscreens formula

Formula	Colour	Smell	Consistency
Gel base	Transparent	Carbomer like	A bit viscous
Sunscreen (S)	Pale Yellow	Padimate-O like	Viscous
S+Arbutin (A)	Pale Yellow	Padimate-O like	Viscous
S+A+Olive oil (O) 3%	Yellowish	Sesame oil like	Viscous
S+A+O 5%	Yellowish	Sesame oil like	Viscous
S+A+O 7%	Yellowish	Sesame oil like	Viscous

Table 3: pH of sunscreen formula

Formula	pH (average)	% var. coefficient
Gel base	6.60 ± 0.07	1.05
Sunscreen (S)	6.31 ± 0.04	0.74
S+Arbutin (A)	6.27 ± 0.05	0.70
S+A+Sesame oil (O) 3%	6.26 ± 0.02	0.28
S+A+O 5%	6.30 ± 0.04	0.63
S+A+O 7%	6.30 ± 0.05	0.83

* The result were obtained from an average of 3 times replication

One of the important factors that influence sunscreen SPF value is pH, besides extinction coefficient and solvent polarity. Therefore it is important to know the mechanism

of SPF changes in the treatment formula. The pH of sunscreen gels also play a major rule in the sunscreen gels characteristic as the carbomer consistency were heavily affected by its acidity. Acid condition would lower the gels viscosity. From pH data, value of F calculation = 26.083 > F table = 5.42. Tukey HSD test of the pH data has shown decreasing pH by addition of sunscreen ingredient. However, addition of arbutin and/or sesame oil did not affect the pH even though sesame oil concentration increased. Overall, from the pH data we know that all sunscreen formula were in a range of skin pH.

The spread-ability of the formula was measured and the results were shown in table 4: From the data we could see that the spread-ability of the formula increased with the addition of arbutin but decreased with a rise in sesame oil concentration.

Table 4: Spread-ability of sunscreens formula

Formula	Average slope (mm/g)	% Var. coefficient
Gel base	0.2833 ± -	-
Sunscreen (S)	0.3359 ± 0.004	2.26
S+Arbutin (A)	0.3859 ± 0.004	3.09
S+A+Sesame oil (O) 3%	0.2713 ± 0.002	3.47
S+A+O 5%	0.2474 ± 0.003	2.91
S+A+O 7%	0.2238 ± 0.0007	1.25

* The result were obtained from an average of 3 times replication

The SPF analysis was done by extracting the sunscreen from its base by using isopropanol. In order to assure that the base gel *carbomer* 940 will not give any absorption in the UV spectrum, it is also extracted using isopropanol and observed at the wavelength of 290-400 nm. From the spectra observed it is known that the gel base did not give any absorption at all, Therefore it is assumed that any change within the observation of SPF value were caused by the active ingredient only.

The SPF value that resulted from the calculation (Table 5) was then categorized based on the American Society of Health System Pharmacist standard in order to classify its protection capability.

Table 6: SPF data from the treatment formula

Formula	Average SPF** ± SD	% KV	SPF Category (American Society of Health System Pharmacist Standard)	% increasing SPF value compare with control
I: S (Sunscreen preparation in gel)	6,658 ± 0,15	2,21	Moderate Sun Protection Product	0
II: S + arbutin	9,094 ± 0,24	2,63	High Sun Protection Product	36.58
III: S+A+O 3%	9,615 ± 0,50	5,18	High Sun Protection Product	44.42
IV: S+A+O 5%	11,119 ± 0,34	3,08	High Sun Protection Product	67.01
V: S+A+O 7%	16,435 ± 0,61	3,72	Very High Sun Protection Product	146.84

**The data were obtained from an average of 3 different products in 1 formula




Figure 2: Absorption spectrum of sunscreen

According to category as presented Table 5, we know that arbutin elevated the sunscreen protection category, from "Moderate Sun Protection Product" to "High Sun Protection Product". Increasing sesame oil up to 5% did not alter the category. However addition of 7% sesame provide the maximum protection resulted the sunscreen category enhanced to be "Very High Sun Protection Product"

From the screening spectra profile as shown in Figure 2, it is learned that the addition of arbutin and sesame oil up to 5% did not cause any shiftness on the maximum wavelength. Nevertheless, an increase in the intensity of absorption was observed. Thus it is predicted that an interaction occurred between the molecule of arbutin and sunscreen agent which intensify the effect of aucsochrome group. And it is clear that addition of 7% sesame oil not only increased the absorption intensity but also increased the protection sunscreen, by maximum wavelength shift. This shiftness cause decreasing polarity of sunscreen gels that affect the delocalization of the molecule and resulted in a rise of energy demand needed for excitation to happen and hence, increase the SPF value.

Table 9: HSD test of the sunscreens SPF

SPF

TukeyHSD^a

Kelompok	N	Subset for alpha = .05			
		1	2	3	4
Formula I	9	6.6580			
Formula II	9		9.0936		
Formula III	9			9.6153	
Formula IV	9				11.1192
Sig.		1.000	1.000	1.000	1.000

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 9.000.

Conclusion

The addition of arbutin and sesame oil affect the physical appearance (organoleptic and consistency) of sunscreen product as well as its spread ability and significant increase in SPF value of the sunscreen formula combination Oxybenzone and Padimate - O (3:7 % w/w) which was formulated in carbomer gel base.

Acknowledgement

This study was supported financially by Project Grant Faculty of Pharmacy, Airlangga University. We would also like to thank Surya Dermato Medica Factory for providing the materials needed for the sunscreens preparation.

References

- American Society of Health-System Pharmacists, 2002, AHFS Drug Information, USA: American Pharmaceutical Association.
- Galilee, 2008, Technical Data Sheet: Arbutin (Beta-Arbutin), Israel.
- Mashood, A.A., 2006, Treatment of Hyperpigmentation Disorders, Journal of Pakistan Association of Dermatologists, Vol. 16, p. 65-68.
- Mitsui, T., 1998, New Cosmetic Science, 2nd ed, Amsterdam: Elsevier
- O' Neil, M.J., 2006, The Merck Index, 14th ed, USA: Merck and Co., Inc.
- Petro, A.J., 1981, Correlation of Spectrophotometric Data with Sunscreen Protection Factors, International Journal of Cosmetic Science, Vol. 3, p. 185-196
- Pratiwi, V.K., 2006, Pengaruh Asam Glikolat Terhadap Stabilitas Fisik dan Peningkatan Efektivitas in vitro Sediaan Tabir Matahari Kombinasi Oksibenson-Oktilsimetil PABA (3:7) %b/b, Skripsi, Surabaya: Fakultas Farmasi Universitas Airlangga
- Sarma, K., 1993, Vegetable oil-based skin permeation enhancer compositions and associated method and systems, US Patent, patent no. 5,229,130
- Widianingsih, N.P.T., and Lumintang, H., 2002, Pemakaian Tabir Surya (Sunscreen), Berkala Ilmu Penyakit Kulit dan Kelamin, Vol. 14 No. 2, p. 155-167.