



ICPPS 2014

Proceeding

The 1st International Conference on Pharmaceutics & Pharmaceutical Sciences

Drug Delivery Systems:
From Drug-Discovery, Pre-formulation, Formulation and Technological Approaches for
Poorly Soluble Drugs and Protein



The 1st International Conference on Pharmaceutics & Pharmaceutical Sciences Proceedings

ISBN : 978-602-72333-0-0

(Letter of National ISBN Agency No. 4127/E.8/p/03.2015 Date 18 March 2015)

1st edition Proceeding

Published by:

Faculty of Pharmacy Universitas Airlangga
Surabaya, Indonesia

Address:

Kampus B Jl. Dharmawangsa Dalam
Surabaya 60286

Phone +62 31 5033710

Fax +62 31 5020514

Website: www.icpps2014.com or www.ff.unair.ac.id

Email: icppinfo@gmail.com

ISBN 978-602-72333-0-0



TABLE of CONTENT

Preface from Chairman

Committee	ii
Table of Contents	iii
Author Index	iii

AUTHOR INDEX

COMPARISON OF SODIUM STARCH GLYCOLATE AND CROSSCARMELLOSE SODIUM AS SUPERDISINTEGRANT IN MEFENAMIC ACID FAST DISINTEGRATING TABLET Adeltrudis Adelsa D, Oktavia Eka Puspita, Amalia Ayuningtyas, Marulita Isadora	1
STUDY EXPRESSION OF HUMAN ERYTHROPOIETIN EXPRESSION IN MAMMALIAN CELL Adi Santoso, Popi Hadiwisnuwardhani, Yana Rubiana, Yulaika Romadhani, Endah Puji Septisetyani, Dyanningtyas D.P. Putri	4
ANTIOXIDANT STABILITY ASSAY OF ALPHA TOCOPHERYL ACETATE IN SOLID LIPID NANOPARTICLE SYSTEM (LIPID BASE BEESWAX AND MONOSTEARIC GLISERYL 50:50) Anggie Widhi, Noorma Rosita, Widji Soeratri	8
A BIOACTIVE <i>BOVINE HYDROXYAPATITE</i> –GELATIN IMPLANT FOR IN VITRO GENTAMICIN RELEASE Aniek Setiya Budiadin, M. Zainuddin, Junaidi Khotib, Diah Himawati	13
EFFECT OF COMPARISON SURFACTANT AND COSURFACTANT IN WATER/OIL MICROEMULSION IN RELEASE OF OVALBUMIN Microemulsion Water/Oil with Surfactant (Span 80-Tween 80) : Cosurfactant (Ethanol) =5:1, 6:1, and 7:1) Anisa Rizki Amalia, Riesta Primaharinastiti, Esti Hendradi	18
ANALYSIS OF MYCOLIC ACIDS CLEAVAGE PRODUCT OF <i>Mycobacterium tuberculosis</i> BY GAS CHROMATOGRAPHY-FLAME IONIZATION DETECTOR Asri Darmawati, Deby Kusumaningrum, Isnaeni, Muhamad Zainuddin	21
PERIPLASMIC EXPRESSION OF GENE ENCODING ANTI-EGFRvIII SINGLE-CHAIN VARIABLE FRAGMENT ANTIBODY USING PeIB LEADER SEQUENCE IN <i>ESCHERICHIA COLI</i> Kartika Sari Dewi, Debbie Sofie Retnoningrum, Catur Riani, Asrul Muhamad Fuad	24
IN VIVO ANTIMALARIAL ACTIVITY OF ETHANOL EXTRACT AND ETHYL ACETATE FRACTION OF <i>Alectryon serratus</i> LEAVES ON <i>Plasmodium berghei</i> INFECTED MICE Aty Widyawaruyanti, Uswatun Khasanah, Lidya Tumewu, Hilkatul Ilmi, Achmad Fuad Hafid, Indah S Tantular	30
PROFILE OF COMMUNITY PHARMACISTS KNOWLEDGE IN PATIENT ASSESSMENT WITH INFLUENZA SYMPTOMS AND ITS PRODUCTS Azza Faturrohmah, Arie Sulistyarini, Ana Yuda	33

SOLUBILITY AND DISSOLUTION STUDY OF KETOPROFEN – HIDROXYPROPYL- β -CYCLODEXTRIN INCLUSION COMPLEX (Prepared by Kneading Method) Bambang Widjaja, Achmad Radjaram, Arafah Zulhana	37
FORMULATION AND STABILITY TESTING OF MELOXICAM SOLID DISPERSION GEL Budipratiwi Wisudyaningsih, Inka Dewi Nur Anggaraini, Fersiya Wardani	40
EFFECT OF MENTHOL AS PENETRATION ENHANCER TO DICLOFENAC SODIUM MEMBRANE-TYPED TRANSDERMAL PATCH CHARACTERIZATION Destria Indah Sari, Esti Hendradi, Junaidi Khotib	43
PHYSICAL CHARACTERISTICS AND RELEASE STUDY OF OVALBUMIN FROM ALGINATE MICROSPHERES PREPARED BY DIFFERENT CONCENTRATION OF ALGINATE AND BaCl ₂ USING AEROSOLIZATION TECHNIQUE Dewi Melani Hariyadi, Tristiana Erawati, Sisilia Ermawahyuningtyas	46
MUCOADHESIVE TABLET OF ETHANOLIC EXTRACT OF SAMBILOTO (<i>Andrographis paniculata</i>) AS ANTIDIABETIC USING CHITOSAN Dhadhang Wahyu Kurniawan, Hening Pratiwi, and Lingga Ikaditya	50
PHYSICAL INTERACTION STUDY OF IBUPROFEN-STEARIC ACID BINARY MIXTURE Diajeng Putri Paramita, Dwi Setyawan, Dewi Isadiartuti	59
MOLECULAR MODELING AND SYNTHESIS OF 1-(3,4-Dichlorobenzoyl)-1,3-dimethylurea Dian Agung Pangaribowo, Siswandono, Bambang Tri Purwanto	63
EXPRESSION OF RECOMBINANT HUMAN GRANULOCYTE-COLONY STIMULATING FACTOR WITHIN PERIPLASMIC COMPARTMENT OF <i>Escherichia coli</i> USING PeIB LEADER PEPTIDE Dian Fitria Agustiyanti, Asrul Muhammad Fuad	66
EVALUATION OF ANTIHYPERURICEMIC ACTIVITY FROM BULBS OF BAWANG TIWAI (<i>Eleutherine palmifolia</i> (L.) Merr.) BY IN VITRO AND IN VIVO STUDIES Dian Ratih Laksmiawati, Rininta Firdaus, Yulinda, Mediana Astika	72
ANTIOXIDANT ACTIVITY OF 96% ETHANOL EXTRACT OF COMBINATION OF STRAWBERRY FRUIT (<i>Fragaria x ananassa</i> Duch.) AND STARFRUIT (<i>Averrhoa carambola</i> L.) USING ABTS FREE RADICAL SCAVENGING METHOD Diana Serlahwaty, Indira Natalia Timang	76
ENHANCEMENT OF SOLUBILITY AND DISSOLUTION ATORVASTATIN BY MICROCRYSTALLIZATION METHOD Dolih Gozali, Yoga Windu Wardhana, Ronny Tandela	79
<i>IN VITRO</i> ANTIMALARIAL ACTIVITY OF DICHLOROMETHANE SUB-FRACTION OF <i>Eucalyptus globulus</i> L. Stem AGAINST <i>Plasmodium falciparum</i> Elis Suwarni, Achmad Fuad Hafid, Aty Widayawaruyanti	86
<i>Arcangelisia flava</i> INCREASES RATS' LEUKOCYTES BUT HAS BIPHASIC EFFECT ON RATS' LYMPHOCYTE Endah Puspitasari, Evi Umayah Ulfa, Vita Ariati, Mohammad SulthonHabibi	89

IN VITRO ANTIMALARIAL ACTIVITY OF CHLOROFORM SUBFRACTION OF SALAM BADAK LEAVES (<i>Acmena acuminatissima</i>) Erna Cahyaningsih, Achmad Fuad Hafid, Aty Widyawaruyanti	92
CHARACTERIZATION OF DOSAGE FORM AND PENETRATION DICLOFENAC SODIUM WITH MICROEMULSION SYSTEM IN HPMC 4000 GEL BASE (Microemulsion W/O with ratio use of surfactant Span 80 – Tween 80 : Cosurfactant Ethanol 96% = 6:1) Esti Hendradi, Tutiek Purwanti, Karina Wahyu Irawati	95
CONSTRUCTION AND VALIDATION OF THE STRUCTURE-BASED VIRTUAL SCREENING PROTOCOLS WITH PDB CODE OF 3LN1 TO DISCOVER CYCLOOXYGENASE-2 INHIBITORS Mumpuni E, Nurrochmad A, Pranowo HD, Jenie UA, Istyastono EP	99
VALIDATED UV SPECTROPHOTOMETRIC METHOD FOR THE DETERMINATION OF ASPIRIN IN RABBIT PLASMA : APPLICATION TO BIOAVAILABILITY STUDY OF ASPIRIN MICROCAPSULE IN RABBIT Faizatun, Novi yantih, Teguh Iman Saputra	102
EFFECT OF COMPARISON OF SURFACTANT AND COSURFACTANT W/O MICROEMULSION OVALBUMIN WITH SOYBEAN OIL TO PHYSICO-CHEMICAL CHARACTERIZATION (w/o Microemulsion with Surfactant Span 80- Tween 80 : Cosurfactant Ethanol 96% = 5:1; 6:1 and 7:1) Farida Mutiara Sari, Riesta Primaharinastiti, Esti Hendradi	105
pH INFLUENCE IN DESALTING PROCESS OF CRUDE PERTUSSIS TOXIN (PT) AND FILAMENTOUS HEMAGGLUTININ (FHA) PURIFICATION FROM <i>Bordetella pertussis</i> BY SEPHADEX G-25 COLUMN CHROMATOGRAPHY Faris Adrianto, Esti Hendradi, Neni Nurainy, Isnaeni	108
SEPARATION OF COSMETIC PRESERVATIVES USING SILICA-BASED MONOLITHIC COLUMN Febri Annuryanti, Riesta Primaharinastiti, Moch. Yuwono	111
PREPARATION AND CHARACTERIZATION OF TELMISARTAN-CITRIC ACID CO-CRYSTAL Fikri Alatas, Hestiary Ratih, Sundani Nurono Soewandhie	114
PATIENTS' AND CAREGIVERS' LIQUID MEDICATION ADMINISTRATION ERRORS Gusti Noorrizka Veronika Achmad, Gesnita Nugraheni	117
THE POTENCY OF CANARIUM OIL (<i>Canarium indicum</i>) AS A MATERIAL FOR STRUCTURED LIPID PRODUCTION Hamidah Rahman, Johnner P Sitompul, Kusnandar Anggadiredja, Tutus Gusdinar	121
EFFECT OF TREHALOSE ON THERMAL PROPERTIES OF PHOSPHOLIPID-DDA AND TPGS MIXTURES Helmy Yusuf	124
PREPARATION AND CHARACTERIZATION OF FLUKONAZOLE- β -CYCLODEXTRIN INCLUSION COMPLEXES Hestiary Ratih, Fikri Alatas, Erin Karlina	127
ISOLATION AND IDENTIFICATION OF ANTIOXIDANT COMPOUND BY BIOPRODUCTION OF ENDOPHYTIC FUNGI OF TURMERIC (<i>Curcuma longa</i> L.) ISOLATE CL.SMI.RF11 Hindra Rahmawati, Bustanussalam, Partomuan Simanjuntak	130

MODIFICATION PROCESS OF NATURAL CASSAVA STARCH : THE STUDY OF CHARACTERISTICS AND PHYSICAL PROPERTIES Prasetia, Jemmy A, C.I.S. Arisanti, N.P.P.A. Dewi, G.A.R. Astuti, N.W.N Yulianingsih, I M.A.G. Wirasuta	133
DRUG USE PROFILE OF DIABETIC PATIENTS IN EAST SURABAYA PRIMARY HEALTH CARE I Nyoman Wijaya, Azza Faturrohmah, Ana Yuda, Mufarriha, Tesa Geovani Santoso, Dina Kartika, Hikmah Prasasti N, Whanni Wido Agustin	136
GLYCINE MAX DETAM II VARIETY AS PREVENTIVE AND CURATIVE ORGAN DAMAGE DUE TO EXPOSURE TO ,LEAD (Pb) Rika Yulia, Sylvan Septian Ressandy, Gusti Ayu Putu Puspikaryani, I Putu Agus Yulyastrawan, Dewa Ayu Kusuma Dewi	142
AN ACTIVITY TEST OF MATOA LEAVES EXTRACT AS HEART RATE FREQUENCY REDUCTION WITH ADRENALINE INDUCTION Ika Purwidyaningrum, Elin Yulinah Sukandar, Irda Fidrianny	144
EFFORT TO REDUCE COMPRESSIBILITY OF RAMIPRIL THROUGH CRYSTAL ENGINEERING Indra, Sundani N Soewandhi	147
IN VITRO ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY OF ETHANOLIC LEAF EXTRACT AND FRACTIONS OF <i>Rauwolfia serpentina</i> (L.) Benth. ex Kurz Julie Anne D. Bolaños, Ivan L. Lawag	150
PERIPLASMIC EXPRESSION OF GENE ENCODING ANTI-EGFRvIII SINGLE-CHAIN VARIABLE FRAGMENT ANTIBODY USING PeIB LEADER SEQUENCE IN <i>Escherichia coli</i> Kartika Sari Dewi, Debbie Sofie Retnoningrum, Catur Riani, Asrul Muhamad Fuad	153
CHARACTERIZATION AND LD ₅₀ VALUE DETERMINATION OF 1,5-bis(3'-ethoxy-4'-hydroxyphenyl)-1,4-pentadiene-3-one (EHP) Lestari Rahayu, Septian, Esti Mumpuni	159
DEVELOPMENT OF MELOXICAM TRANSDERMAL MATRIX TYPE PATCH USING POLYVINYLPIRROLIDONE, HYDROXYPROPYL METHYLCELLULOSE, AND ETHYL CELLULOSE COMBINATION Lidya Ameliana, Monica Iwud, Selly Rio	162
ANTIHEPATITIS C VIRUS ACTIVITY SCREENING ON <i>Harpullia arborea</i> EXTRACTS AND ISOLATED COMPOUND Lidya Tumewu, Evhy Apryani, Mei Ria Santi, Tutik Sri Wahyuni, Adita Ayu Permanasari, Myrna Adianti, Chie Aoki, Aty Widyawaruyanti, Achmad Fuad Hafid, Maria Inge Lusida, Soetjipto, Hak Hotta.....	165
HPLC METHOD PRECISION TO ASSAY OF A-MANGOSTIN IN Mangosteen (<i>Garcinia mangostana</i> L.) FRUIT RIND EXTRACT FORMULATED IN ORAL SOLUTION Lilie Nurhidayati, Siti Sofiah, Ros Sumarny, Kevin Caesar	168

PREPARATION AND CHARACTERIZATION OF NARINGENIN-LOADED CHITOSAN NANOPARTICLES FOR CHEMOPREVENTION Lina Winarti, Lusia Oktora Ruma Kumala Sari	170
RELATIONSHIP OF KNOWLEDGE AND PATIENT BEHAVIOR ON SELF MEDICATION PIROXICAM (Studies of Pharmacy in Sukun District, Malang City) Liza Pristianty, Reshtia Eriana Putri, Hidayah Rachmawati	173
EFFECT OF CHRONIC USE OF ENERGY DRINK ON KIDNEY Mahardian Rahmadi, Zamrotul Izzah, Mareta Rindang A, Aniek Setya B, Suharjono	176
SCREENING OF SURFACE MODIFIERS TO PRODUCE STABLE NANOSUSPENSION : A GENERAL GUIDANCE Maria Lucia Ardhani Dwi Lestari	179
DEVELOPMENT OF SIMPLE POLYPHENOL SENSOR BASED ON SODIUM META PERIODATE AND 3-METHYL-2-BENZOTHAZOLINONE HYDRAZONE FOR COFFEE SAMPLES Moch. Amrun Hidayat, Nindya Puspitaningtyas, Agus Abdul Gani, Bambang Kuswandi	181
VALIDATION OF AN HPLC ANALYTICAL METHOD FOR DETERMINATION OF LEVOFLOXACIN IN OPHTHALMIC PREPARATIONS Mochamad Yuwono, Riesta Primaharinastiti, Ageng Teguh Wardoyo	184
VALIDATION OF SPECTROPHOTOMETRIC METHOD FOR ESTIMATION OF EPERISONE HYDROCHLORIDE IN TABLET DOSAGE FORM Nia Kristiningrum, Diah Yuli Pangesti	187
ANTIOSTEOPOROTIC ACTIVITY OF 96% ETHANOLIC EXTRACTS OF ABELMOSCHUS MANIHOT L.MEDIK LEAVES AND EXERCISE ON INCREASING BONE DENSITY OF FEMALE MICE'S FEMORAL TRABECULAR Niliestria Ayu Faramitha Sholikhah	190
EFFECT OF -CYCLODEXTRIN ON SPF VALUE AND INHIBITION OF KOJIC ACIDSTYROSINASE ACTIVITY IN VANISHING CREAM BASE FORMULATION (ON SUNSCREEN PRODUCT CONTAINED OXYBENZONE) Noorma Rosita, Diana, Diana Winarita, Tristiana Erawati, Widji Soeratri	193
ANTIMICROBIAL ACTIVITY OF LACTOBACILLI PROBIOTIC MILK AND GUAVA LEAF ETHANOLIC EXTRACT (Psidium guajava) COMBINATION AGAINST BACTERIAL CAUSE OF DIARRHEA Nur Putri Ranti, Isnaeni, Juniar Moechtar, Febri Annuryanti	197
THE INFLUENCES OF PARTICLE SIZE AND SHAPE ON ZETA POTENTIAL OF COENZYME Q10 NANOSUSPENSION Nuttakorn Baisaeng	200
SYNTHESIS, MOLECULAR DOCKING, AND ANTITUMOR ACTIVITY OF N,N'-Dibenzoyl-N,N'-Dimethylurea AGAINST HUMAN BREAST CANCER CELL LINE (MCF-7) Nuzul Wahyuning Diyah	203
EXPRESSION OF ANTI-EGFRVIII SINGLE CHAIN FRAGMENT ANTIBODY (SCFV) ON THE SURFACE OF PICHIA PASTORIS Pratika Viogenta, Asrul Muhamad Fuad, Suharsono	206

SUBCLONING OF <i>csf3syn</i> (COLONY STIMULATING FACTOR-3) GENE INTO pGAPZ? AND TRANSFORMATION OF RECOMBINANT VECTOR INTO PICHIA PASTORIS Prety Ihda Arfia, Asrul Muhamad Fuad	212
THE USE OF PERICARP MANGOSTEEN (<i>Garcinia mangostana</i> L.) EXTRACT IN FORMULATION OF CREAM-TYPE O / W Rahmah Elfiyani, Naniek Setiadi Radjab, Mia Sagita Sofyan	218
CHITOSAN BASED PARTICULATE CARRIER OF DITERPENE LACTON OF SAMBILOTO PREPARED BY IONIC GELATION-SPRAY DRYING :EFFECT OF STIRRING RATE AND NOZZLE DIAMETER Retno Sari, Titin Suhartanti, Dwi Setyawan, Esti Hendradi, Widji Soeratri	222
GAS CHROMATOGRAPHY-MASS SPECTROMETRY METHOD VALIDATION FOR PESTICIDES RESIDUES ANALYSIS IN FOOD USING QuEChERS KIT Riesta Primaharinastiti, Setyo Prihatiningtyas, Mochammad Yuwono	225
CHARACTERIZATION OF PARACETAMOL ORALLY DISINTEGRATING TABLET USING GELATIN 1% AND 2% AS BINDER AND POLYPLASDONE XL-10 10% AS DISINTEGRANT (Prepared by Freeze Drying Method) Roisah Nawatila, Dwi Setyawan. Bambang Widjaja	230
ANTIOXIDANT STUDY OF COSOLVENT SOLUTION OF MANGOSTEEN (<i>Garcinia mangostana</i> L.) RIND EXTRACT IN RATS BY USING MDA PARAMETER Ros Sumarny, Liliek Nurhidayati, Yati Sumiyati, Fransiska Diana Santi	233
SCREENING OF SELECTED PHILIPPINE ROOT CROPS FOR β -Glucosidase INHIBITION Sarah Jane S. Almazora, Ivan L. Lawag	236
NIOSOME EMULGEL FORMULATION AND STABILITY TEST OF CINNAMON (<i>Cinnamomum burmanii</i> Nees & Th. Nees) Bark ETHANOLIC EXTRACT AS ANTIOXIDANT Sasanti Darijanto, Fidriani Irda, Widhita P.A.S	239
THEOPHYLLINE RELEASE FROM SUSTAINED RELEASE TABLET USING LACTOSA AND PVP K30 AS A CHANELLING AGENT Sugiyartono, Retno Sari, Agus Syamsur Rijal, TriMulyani, Agustina Maharani	242
EFFORTS TO PRODUCE 1-(BENZOYLOXY)UREA AS ANTICANCER DRUG CANDIDATE Suko Hardjono	245
DEVELOPMENT OF PEGylated RAPAMYCIN LITHOCHOLIC ACID MICELLE FOR CANCER THERAPY Aran Tapsiri, Kanokwan Jaiprasert, Rungtip Nooma, Awadsada Sukgasem, Supang Khondee	248
FEASIBILITY OF ORAL IMMUNIZATION AGAINST JAPANESE ENCEPHALITIS VIRUS USING CHITOSAN PARTICLES Supavadee Boontha, Worawan Boonyo, Hans E. Junginger, Tasana Pitaksuteepong, Neti Waranuch, Assadang Polnok, Narong Nitatpattana, Sutee Yoksan	251

THE INFLUENCE OF HYDROXY GROUP AT ORTHO (o) AND PARA (p) POSITIONS ON METILBENZOAT AGAINST SYNTHESIS OF HIDROKSIBENZOAHIDRAZIDA DERIVATIVES Suzana, Adita R, Melanny Ika .S, Juni Ekowati, Marcellino Rudyanto, Hadi Poerwono, Tutuk Budiati	254
ANTIOXIDANT ACTIVITY TEST OF BEE PROPOLIS EXTRACT (Apis mellifera L.) USING DPPH (1,1-diphenyl-2-picrilhidrazy) FREE RADICALS SCAVENGING ACTIVITY Titiek Martati, Shahyawidya	258
CAPSULE FORMULATION and EVALUATION COMBINATION OF AQUEOUS EXTRACT OF Phaleria's (Phaleria macrocarpa (Scheff Boerf)) FRUITS and LEAVES as ANTIHIPERTENSIVE AGENT Titta H Sutarna, Sri Wahyuningsih, Julia Ratnawati, Fahrouk P, Suci Nar Vikasari, Ita Nur Anisa	260
MANUFACTURE AND CHARACTERIZATION OF SOLID DISPERSION GLIKLAZID- PVP K90 Titta H Sutarna, Fikri Alatas, Cich Ayu Ningsih	264
ANTI-INFLAMMATORY ACTIVITY OF PARA METHOXY CINNAMIC ACID (PMCA) IN NANOEMULSION USING SOYBEAN OIL Tristiana Erawati M, Anneke Indraswari P, Nanda Ghernasih N.C, Noorma Rosita, Suwaldi Martodihardjo, Widji Soeratri	266
PHYSICAL CHARACTERISTICS AND PENETRATION OF DICLOFENAC SODIUM NIOSOMAL SYSTEM USING SPAN 20 AND SPAN 60 Tutiek Purwanti, Esti Hendradi, Noverika A. Putri, Nurtya J. Devi	271
FORMULATION AND CHARACTERISZATION OF JUICE OF LIME GEL USING CMC-Na BASE Uswatun Chasanah, Esti Hendradi, Inayah	275
HEPATOPROTECTIVE ACTIVITY OF Bidens pilosa L. IN CARBON TETRACHLORIDE INDUCED HEPATOTOXICITY IN RATS Vina Alvionita Soesilo, C.J. Soegihardjo	278
CYTOTOXIC ACTIVITY ASSAY AGAINTS HELA CELL LINES OF NOVEL ANTICANCER DRUG : N-(PHENYL CARBAMOYL)ISOBUTIRAMIDE Wimzy Rizqy Prabhata, Tri Widiandani, Siswandono	284
SIMPLE STEPS PURIFICATION OF RECOMBINANT HUMAN ERYTHROPOIETIN PRODUCED IN CHINESE HAMSTER OVARY CELL CULTURE Yana Rubiyana, Endah Puji Septisetyani, Adi Santoso	287
KINETICS STUDY COCRYSTALS KETOCONAZOLE-SUCCINIC ACID PREPARED WITH SLURRY METHOD BASED ON POWDER X-RAY DIFFRACTION (PXRD) Yuli Ainun Najih, Dwi Setyawan, Achmad Radjaram	290
CONSTRUCTION OF RECOMBINANT IMMUNOTOXIN Anti-EGFRvIII scFv::HPR FUSION PROTEIN AND INDUCIBLE EXPRESSION IN Pichia pastoris AS A TARGETED DRUG CANDIDATE Yuliawati, Asrul Muhamad Fuad	297
ALTERED PHARMACOKINETIC OF LEVOFLOXACIN BY COADMINISTRATION OF ATTAPULGITE Zamrotul Izzah, Toetik Aryani, Amalia Illiyin, Budi Suprpti	303

EFFECT OF β -Cyclodextrin ON SPF VALUE AND INHIBITION OF KOJIC ACID TYROSINASE ACTIVITY IN VANISHING CREAM BASE FORMULATION (ON SUNSCREEN PRODUCT CONTAINED OXYBENZONE)

Noorma Rosita, Pharmaceutics Department, Faculty of Pharmacy Airlangga University, itanr@yahoo.com or noorma-r@ff.unair.ac.id, +6281331031346; **Diana**, Pharmaceutics Department, Faculty of Pharmacy Airlangga University, **Diana Winarita**, Pharmaceutics Department, Faculty of Pharmacy Airlangga University, **Tristiana Erawati**, Pharmaceutics Department, Faculty of Pharmacy Airlangga University, **Widji Soeratri**, Pharmaceutics Department, Faculty of Pharmacy Airlangga University

INTRODUCTION

Newly, we found many cosmetics combined of sunscreen substance and lightening agent in their ingredients.

Kojic acid is a skin lightening agent which has very small molecular size, so easily absorbed up through the basal membrane. Because of its small molecule, kojic acid hardly absorbed through the lipid membrane of its target sites, the melanocytes (Manosroi et al, 2005) and can penetrate into the systemic (Nakayama et al., 2005). It is likely absorbed through voids between cells on the skin. Kojic acid is produced mainly by microbial fermentation using *aspergillus* and *penicillium* spp. In vivo test, cream containing kojic acid compounds have been reported as effective in preventing pigmentation changes in human skin due to exposure to UVA and UVB. This inhibition has been shown to be due to chelation of Cu, a prosthetic group in tyrosinase (Barel et al, 2001). Cream contained 1 % of kojic acid showed steady but slow whitening effect. ¹⁴C-labeled kojic acid cream was observed to be quickly absorbed from the skin to the liver, intestines, and kidneys in mice.




Fig 1. Properties of Kojic Acid
5-hidroksi-2-hidroksimetil-4-pyrone

MW: 142.1
MP: 153-154oC
pKa: 7.90-8.
Log P (octanol-water): -0,64

When the absorption was thus quick, the depigmentation agent did not stay at the epidermis where it had its target organ, melanocytes, for a long enough time to inhibit melanogenesis. Therefore, kojic acid was mixed with β -cyclodextrin to slow the absorption into the dermis (Elsner and Maibach, 2005). Losses due to kojic acid to the systemic is not able to work effectively inhibits melanin formation took place in the basal membrane of skin that results are not optimal lightening effect. Addition of β -cyclodextrin reported can decrease penetration of kojic acid (Nakayama et al., 2005).

Cyclodextrins are cyclic oligosaccharides with a hydrophilic outer surface and a somewhat hydrophobic central cavity. Cyclodextrins are able to form inclusion complexes with many drugs by taking up the drug molecule, or a lipophilic moiety of the molecule, into the cavity (Loftsson and O'Fee, 2003).




Fig 2. β -cyclodextrin (Sweetman, 2007)



Sunscreen agents, such as: oxybenzone and octyl dimethyl PABA, are commonly present in whitening products to compensate for the photosensitivity effect caused by the whitening agent. This combination is more efficient but may have a profound effect on the efficacy of the sunscreen agent.

Therefore, this study was aimed to investigate the SPF value of sunscreen product contained oxybenzone and octyl dimethyl PABA (3:7) with the addition of kojic acid 1% b/b as whitening agent and their complex with β-cyclodextrin in 1:1 molar equivalent. In this research, we also studied the effect of β-cyclodextrin (BCD) on penetration of Kojic Acid, as lightening substance In Vanishing Cream Base Formulation (on Sunscreen Product Contained Oxybenzone), by determination of tyrosinase activity inhibition.

MATERIAL AND METHODE

Material

Oxybenzone, Octyldimethyl PABA and Kojic Acid were from Surya Dermato, β-cyclodextrin p.a (Sigma Aldrich), Stearic Acid, cetyl alcohol, span 80, tween 80), Methyl paraben Propyl paraben, sorbitol 70%, Isopropanol p.a., (Brataco Chemicals). All of the active ingredients used here were in a pharmaceutical grade except for the chemical reagent isopropanol and β- cyclodextrin which were in pro analytical grade. While material for inhibition of tyrosinase activity test is the Mushroom tyrosinase, L-Tyrosine, NaH2PO4. 2H2O (p.a) and Na2HPO4 (p.a).

The SPF assay were carried out by using a Double Beam Spectrophotometer UV- Vis Perkin Elmer Lambda EZ 201, Ultrasonic Branson 3510, Hettich zentrifugen EBA 20, Mettler Toledo AL 204 analytical balance. Digital pH meter Schott CG 842 and spreading capacity measurer are also used for the organoleptic analysis.

Methode

The sunscreen product contained oxybenzone and octyl dimethyl PABA (3:7) with the addition of kojic acid 1% w/w as whitening agent and their complex with β-cyclodextrin in 1:1 molar equivalent. Vanishing cream as base was made with inversion technique.

The characteristics of the finished product evaluated were emulsion type, pH, spreading-ability, and organoleptic such as colors, odors.

Table 1. Formula composition

Component	Percentage		
	F1	F2	F3
Kojic acid	1	1	1
β cyclodextrin	4	-	4
oxybenzone	-	3	3
Octyl dimethyl PABA	-	7	7
Vanishing cream base	Ad 100	Ad 100	Ad 100

Determination of SPF value

The SPF value was determined by Petro correlation for in vitro method. 2 mg/cm2 or 2 μL/ cm2 sunscreen agent for in vivo test was equivalent with 10 ppm sunscreen agent dissolved in isopropanol. An UV spectrum of this solution was then measured at 290-400 nm by using Double Beam UV-Vis Spectrophotometer Perkin Elmer Lambda EZ 201 at an interval of 2 nm which has absorbance for 0.05 or more. According to Petro, the absorbance was then converted into the absorbance for 10 ppm solution concentration for each wavelength. Then it was proceed in this following equation:

$$AUC_{\lambda_{p-\alpha}}^{\lambda_p} = \frac{A_{p-\alpha} + A_p}{2} (\lambda_p - \lambda_{p-\alpha})$$

Whereas:

AUC = Area under curve

Ap = Absorption on p wavelength

Ap-α = Absorption on p-α wavelength



The total AUC were obtained by totaling each AUC between 2 wavelengths in series from 290 nm till 400 nm which has an absorbance value above 0.050 and the SPF value of a formula were obtained by inserting the total AUC into the equation below:

$$\text{Log SPF} = \frac{\text{Total area}}{\lambda_n - \lambda_1} \times 2$$

Whereas:

λ_n = longest wavelength above 290 nm that has an absorbance higher than 0.050

λ_1 = shortest wavelength 290 nm

Determination of tyrosinase activity inhibition:

A. Preparation of a solution of the reaction components. The solution should be prepared for the implementation of this study was 0.1 M phosphate buffer pH 6.5, solution of tyrosinase (5370 units / mg solid in 0.1 M phosphate buffer pH 6.5 to 100.0 ml volume), a solution of 5.52 mM L-tyrosine, and 30% TCA.

B. Preparation of test sample solution. The cream (around 3 grams) was put in the diffusion cell then covers with the Millipore membrane which was impregnated with isopropyl-myristate as modified lipid membrane. Then the preparation of cream in diffusion cell was put into the penetration chamber contain 500 ml of phosphate buffer pH 6.5 ± 0.05 at 37 ± 0.5°C as diffusion medium, and then the paddle was stirred 100 rpm. The sample solution around 3 ml was collected at 6 hours after it penetrated.

Determination of the dopakrom maximum wavelength. L-tyrosine solution of 5.52 mM has taken a number of prepared and then added 0.5 ml 3 ml of buffer solution pH 6.5 ± 0.05. Then the mixture add with 1.0 ml of tyrosinase and oxygenated for 5 minutes. The mixture was incubated for 15 minutes at 26 ±

0.5 °C then added 0.5 ml of 30% TCA. The solution is then inserted into cuvet and placed on the sample position in the spectrophotometer. Used as a blank solution 0.1 M phosphate buffer pH 6.5. Then do the reading of absorbance values from a wavelength of 400 nm to 500 nm, and the selected wavelength that gives the greatest absorption.

Inhibition Tyrosinase Activity Determination
 L-tyrosine solution 5.52 mM added with 0.5 ml 3 ml of sample solution. Then the mixture added with 1.0 ml of tyrosinase and oxygenated for 5 minutes. The mixture was incubated for 15 minutes at 26 ± 0.5 °C. Then add 0.5 ml of TCA 30% and observed the absorbance at dopakrom maximum wavelength. Inhibition tyrosinase activity is the percent inhibition values obtained from the later absorbance value calculated by the equation:

$$A \times 100$$

$$\% \text{ Inhibition} = 100 -$$

B

A = Absorbance at the maximum with the skin lightening

B = Absorbance at maximum λ without the skin lightening

RESULT AND DISCUSSION

All the formulas had either pH in skins pH range or similar spreadability.

The SPF value increased in formula 3. The complex formation of the sunscreen agent with BCD was suspected to be responsible for this phenomenon. It might be caused of complex formation of the sunscreen agent with β - cyclodextrin increased sunscreen agent solubility.

Based on one way ANOVA test, known that the addition of BCD decreased inhibition percent compared to the control constraints. Nevertheless the addition of sunscreen does not affect the inhibition percent of tyrosinase activity. Addition of BCD on combination kojic acid-sunscreen preparations showed the



same phenomena such as decreased its inhibition percent of tyrosinase activity. Decreasing of inhibition percent of tyrosinase activity after addition of BCD or combination BCD-sunscreen was might be caused by decreasing of kojic acid penetration. It was probably caused by some of the things that the release process of kojic acid from base and in the process of penetration through the membrane. Before reach basal membrane where the kojic acid site of action, kojic acid molecule should released from base further more penetrate through the skin. Increasing viscosity will decrease the molecules mobility of active ingredient that caused barriers against the release of kojic acid.

Table 2. Average of pH

FORMULA	pH ± SD
1	4,51 ± 0,08
2	4,29 ± 0,03
3	4,04 ± 0,03




Figure 3. Profile of Spreading-ability




Figure 4. Profile of sunscreens absorbance measurement

Table 3. Average of SPF

Formula	SPF (average)
1	10,607 ± 0,432
2	11,741 ± 0,479
3	19,113 ± 0,295

Table 5. Average percent of tyrosinase inhibition

Formula	% tyrosinase activity inhibition
1	28,39 ± 0,51
2	60,50 ± 0,98
3	23,69 ± 0,61

CONCLUSION

1. The complex formation of the sunscreen agent with BCD decreased the SPF value of sunscreen product.
2. BCD decrease penetration kojic acid based on inhibition of tyrosinase enzyme activity in invitro.
3. The existence of BCD and combination sunscreen oxybenzone and octyl dimethyl PABA (3:7)% w/w decrease penetration kojic acid based on inhibition of tyrosinase enzyme activity in invitro.

REFERENCE

Barel, A.O., Paye, M., Maibach, H.I., 2001. Handbook of Cosmetic Science and Technology. New York: Marcel Dekker Inc.

Elsner, P., Maibach, H.I., 2005. Cosmeceuticals and Active Cosmetics: Drug Versus Cosmetics, 2nded, Boca Raton: Taylor and Francis Group.

Loftsson, T., O'Fee, R., 2003. Cyclodextrins in Solid Dosage Forms. Business Briefing Pharmatech.

Manosroi., Wongtrakul., Midorikawa., Hanyu., Yuasa., Sugawara., Sakai., Abe, 2005. The Entrapment of kojic oleate in bilayer vesicles, International Journal of Pharmaceutics.

Nakayama, H., Ebihara, T., Satoh, N., dan Jinnai, T., 2000. Depigmentation Agents. In: Elsner, P., dan Maibach, H.I. (eds.). Cosmeceuticals and Active Cosmetics: Drug vs Cosmetics, Boca Raton: Taylor & Francis, p. 185-204.

Petro, A.J., 1981. Correlation of Spectrophotometric Data with Sunscreen Protection Factors. International Journal of Cosmetic Science, Vol. 3, p. 185-196.