

DAFTAR PUSTAKA

- Adhiatma, K, Dony, Y 2015, *Anemia pada Penyakit Kronik*, Hematologi Eritrosit dan Kelainannya.Ed.1, Lembar S, Tjahyadi C dkk, Universitas Katolik Indonesia Atma Jaya, 293-305.
- Angeles Vazquez, Lopez, M, Molonis, FL, Carmona, ML, Morales, AC, Munoz Fico, Fj 2006, Serum Transferin Receptor in Children: Usefulness for Determining the Nature of Anemia in Children, *Journal Pediatric Hematologica Oncology*; 28:809-15.
- Agarwal, MB, Swatipai 2017, Reticulocyte Hb Content (CHR): The Gold Standart for Diagnosing Iron Deficiency, *Journal of The Association of Physicians of India*, 65
- Babit JL, Lin HY 2012, Mechanisms of anemia in CKD. *Journal American Society Nephrology*. 23(10), 1631-4.
- Braga, F, Infusino 2014, Soluble Transferin Receptor in Complicated Anemia. *Clinical Chemistry Acta*, 431:143-147.
- Brugnara, Schiller 2006, Reticulocyte Hemoglobin Equivalent (RET-He) and Assessment of Iron Deficient States, *Clinical Laboratory Haematology* 28, 303-308
- Buttarello, M, Temporin, V, *et al* 2004. The New Retyculocyte Parameter RET-Y of the Sysmex XE 2100, Its Use in the Diagnosis and Monitoring of Post Treatment Sideropenic Anemia, *American Journal of Clinical Pathology*, 130, 104-116
- Chung M *et al.*, 2012, Biomarkers for Assessing and Managing IDA in Late- Stage CKD, Evaluating Iron Status in HD patients, Review Article, *Nephrology Nursing Journal*.
- Dalimunthe, NN, Lubis, AR 2016, Usefulness of Reticulocyte Hemoglobin Equivalent in Management of Regular Hemodialysis Patients with Iron Deficiency Anemia, *Roman Journal International Medicine*. 54, 31-36
- Das Gupta, S, Uppal, B, Pawar, B 2003, High serum Transferin Receptor level in Anemia of Chronic Disorders indicates coexistent iron deficiency, *American Journal Hematology*,72:158-161.
- Dimitriou, H, Stiakaki,E, Markaki, EA 2000, Soluble transferin Receptor levels and soluble transferrin receptor/log ferritin index in the Evaluation of Erythropoietic Status in Childhood Infections and Malignancy, *Acta Paediatrica*, 89: 1169-73

- Fishbane, S, Imbriano L.J, *et al.*,1997, Reticulocyte Hemoglobin content in the Evaluation of Iron Status of Hemodialysis Patient, *Kidney International* 52. 217-222.
- Fishbane, S, Shapiro, W, Dutka, P, Valenzuela OF, Faubert J 2001, A Randomized Trial of Iron Deficiency Testing Strategies in Hemodialysis Patients, *Kidney International*, 60, 2406-11.
- Goyal Richa, Das Reena 2008, Serum transferrin Receptor-Ferritin index shows concomitant iron deficiency anemia, anemia of chronic disease is common in Patients of Rhematoid Arthritis, *Journal Pathologis Microbiology*, 51, 102-104
- Heil, W, Ehrhard, V 2008, *Clinical Chemistry and Immunological Tests, serum/plasma*, Reference Ranges for Adults and Children, 9thedition. Mannheim Roche Diagnostics GmbH, 68-69.
- Hoffbrand R, Edward, J.Benz, Jr 2018, 'Pathophysiology of Iron Homeostasis', *Hematology Basic Principles and Practice*, 7th edition, 468-70.
- Indonesian Renal Registry 2013, 6th Report of Indonesian Renal Registry, Jakarta,,: 21-24.
- Jankowska Ewa, A 2013. Iron Deficiency in Patients with Chronic Kidney Disease Requiring Hemodialysis: do Inflammation and Overexpressed Hcpidin contribute Significantly, *Polskie Archwum Medycyny Wewnfrznej* 123, 1-2.
- KDIGO 2012, KDIGO, Clinical Practice Guideline for Anemia in Chronic Kidney Disease. *Kidney International Supplement*, 2, 292-8.
- Lankhorst, C.E, Wish J.B 2010, Anemia in Renal Disease: Diagnosis and Management, *Blood Reviews*, 24.39-47.
- Latif, A, Alam, MR, Khanam, A, Hoque, F, Rahim, MA, Islam, RN 2017, Role of Serum Transferrin Receptor in Diagnosing and Differentiating Iron Deficiency Anemia from Anemia of Chronic Disease in Patients with Chronic Kidney Disease, *BIRDEM Medical Journal*, 7(2), 132-137.
- Lopez Ruzafa E, *et al.* 2016, Reference Values of Reticulocyte Hemoglobin Content and their relation with other indicators of Iron Status in Healthy Children, *J.Paediatric Hematology Oncology*.
- Malope, BL, Machpail, AP, Albert, M, Hiss, DC 2001, The Ratio of serum Transferrin Receptor and Serum Ferritin in Diagnosis of Iron Status. *Bitish Journal Haematologia*, 115: 84-89

- Malyszko, J, Koc.Zorawska, *et al.* 2012, New Parameters in Iron Metabolism and Functional Iron Deficiency in Patients on Maintenance Hemodialysis. *Pol Arch Medicine Wewn*, 122:537-542.
- Markovic, M, Majkic, S 2005, Usefulness of sTfR and Ferritin in Iron Deficiency Anemia and Anemia Chronic Disease. *Scandinavian Journal Clinical Laboratory Investment*, 65(7), 571-6.
- Mast, AE, Blinder, MA, *et al.* 2002, Cincial Utility of the Reticulocyte Hemoglobin Content in the Diagnosis of Iron Deficiency, *Blood* , 99: 1489-91
- Matsuda, A, Beesho, M 2002, Diagnostic Significance of Serum soluble Fransferin receptor in Various Anemic Disesease the First Multi-Institusional Joint Study in Japan. *Journal Haematologia*, 32(3). 225-38
- Mica, N, *et al.* 2010, Usefulness of Measuring Reticulocyte Hemoglobin Equivalent in the Management of Hemodialysis Patients with Iron Deficiency, *International Journal Laboratory Hematology*, 32, 248-55.
- Mittman, N, *et al.* 1997, Reticulocyte Hemoglobin Content Predicts Functional Iron Deficiency in HD Patient Receive rHuEPO, *American Journal Kidney Disease*.
- NKF-KDOQI 2006, KDOQI Clinical Practice Guideline and Clinical Practition Recommendations for Anemia in Chronic Kidney Disease, *American Journal Kidney Disease*, 47 (Suppl 3), S11-145.
- Wirawan, R, Arlinda, T 2017, *Peran parameter Retikulosit Tambahan Pada Tatalaksana Penyakit Ginjal Kronik*, Semiloka Mutu.Badan Khusus PKEL PDS PATKLIN, 59-66.
- Skikne, BS, Punnonen, K, Caldron, PH, Bennett, MT, Rehu, M, Gasior GH, Chamberlin, JS, Sullivan, LA, Bray, KR, Southwick, PC 2011. Improved Differential Diagnosis of Anemia of Chronic Disease and Iron Deficiency Anemia: A Prospective Multicenter Evaluation of soluble Transferrin receptor and the sTfR/log ferritin index, *American Journal Hematology*. 86, 923-7.
- Somvanshi, S, Khan, NZ, Ahmad, M 2012, Anemia in chronic kidney disease patients, *Nephrology*, 1(3), 198-204.
- Sudir, M, Laxmi, K 2016, Reticulocyte Hemoglobin vis-a-vis Serum Ferritin as a Marker of Bone Marrow Iron Store in Iron Deficiency Anemia, *Journal of the Association of Physician of India*, 64
- Suhardjono, Lubis, HR 2011, Penyakit Ginjal Kronik dan Anemia: Definisi dan Klasifikasi. Konsensus Manajemen Anemia Pada Penyakit Ginjal Kronik, Ed.2, Lydia A, PERNEFRI, 1-4

- Sysmex Corporation. XN series Instruction for Use. Kobe: Sysmex 2011, Sysmex Educational Enhancement and Development, Clinical Utility of the Reticulocyte Content in *Blood Journal*.
- Sysmex XE-2100 and Sysmex XN 1000 Operator's manual 2010 and 2016
- Thomas, C, Thomas, L 2002, Biochemical Markers and Hematologic Indices in the Diagnosis of Functional Iron Deficiency, *Clinical Chemistry*, 48: 1066-1076.
- Toki, Y, Ikuta, K 2016, Usefulness of Reticulocyte Hemoglobin Equivalent for Diagnosis of Iron Deficiency, *Blood*, 128, 361.
- Torino Ana Beatriz Barbosa, Maria de Fatima Pererirera *et al.* 2015, Evaluation of Erythrocyte and Reticulocyte Parameters as Indicative of Iron Deficiency in Patients with Anemia of Chronic Disease, *Rev Bras Hematology Hemoter*, 37(2), 77-81.
- Xue, Z, Zhi, Z. 2002. The Diagnostic Role of sTfR in Patient with Various Anemia, *Taipei Journal Medicine*, 65(2), 55-60.