Research Journal Chemistry, ChemTech, Journal PharmTech, Inter...




>>
About Us
Objective
Consultancy
Int. Journal ChemTech
Int. Journal PharmTech
Contact Us
Site Map

Site is Under Reconstruction in Latest Technology....



Home | About Us | Objectives | Cosultancy | Int. Jornal ChemTech Research | Int. Journal PharmTec

0 0 1 2 3 0 7 7 0 All Privacy & Copyrights reserved with Sai Scientific Communication



Click on following links for issues : Site is Under Reconstruction in Latest Techno

Template for writing a Paper

International Editorial Board

Managing Executive Editors[Dev, A & R]

Editor-in-Chief:

Dr. Tsygankova Victoria Anatolyivna

Outstanding Intellectuals of the 21st Century.

Who's Who in the World.

Google Scholar Profile= https://scholar.google.ru/citations?u ResearchGate Profile= https://www.researchgate.net/profile/\

Dr. Biol. Sci., Sci. Dr., Principal researcher, Senior Staff Scier Head of group of screening of synthetic compounds of Depa Nitrogen-Containing Heterocyclic Compounds, Institute of Bioorganic Chemistry and Petrochemistry of Nati Ukraine, 1, Murmanskaya str, Kyiv-94, 02660, Ukraine. Tel. +38(044) 558-53-88; Fax +38(044) 573-25-52; E-mail: vtsygankova@ukr.net; victoria_biotex@i.ua

Associate Editors

1

2

Dr.S.Ravichandran,

Associate Professor in Chemistry, School of Chemical Engineering and Physical Sciences, Lovely Professional University, Jalandhar - Delhi GT Road, Phagwara (Punjab)- 144 411.

Mobile. 09840735064. E-mail: ravichandran.23324@lpu.co.in Google Scholar Profile= https://scholar.google.co.in/citatio ResearchGate Profile= https://www.researchgate.net/profi 1

	Dr. (Ms.) Himani N. Chopde
3	Department of Applied Chemistry, G. H. Raisoni Academy Nagpur, MH, India
	E-mail: himani.chopde@raisoni.net, himanichopade@gmail Tel: 91 9604950439, +91 9405703635
	Google Scholar Profile= https://scholar.google.co.in/citation ResearchGate Profile= https://www.researchgate.net/profile=
	Dr. Siamak Hosseinzadeh.
4	Young Researchers and Elite Club, West Tehran Branch, Is Email: hoseinzadeh.siamak@gmail.com Tel: +989375361451
	Google Scholar Profile= https://scholar.google.com/citation ResearchGate Profile= https://www.researchgate.net/prof
	Karel Diéguez Santana
5	Ceslao Marin, Puyo, Pastaza, Ecuador/ Lecturer. Amazonia Email: karel.dieguez.santana@gmail.com, <u>kdieguez@uea.e</u> Tel: 0995041600
	<u>Google Scholar Profile</u> = https://scholar.google.com.ec/citat <u>ResearchGate Profile</u> = https://www.researchgate.net/profile
	Dr. Dhinesh Balasubramanian
	Mepco Schlenk Engineering College, Sivakasi, Virudhunagar, Tamil Nadu – 626005, India
6	Email: dhineshbala91@gmail.com, dhineshbala91@mepco Tel: +91 9791625490,04562 235603
	<u>Google Scholar Profile</u> = https://scholar.google.com/citation <u>ResearchGate Profile</u> = https://www.researchgate.net/profile
	Dr. Arumuga Perumal D
	7/20 C.M.S.South Street, INFORMATION Karungulam (Post),
7	Srivaikundam (Taluk), Tuticorin-628615 Tamilnadu, INDIA/ NIT Karnataka, Surathkal, Mangalore, Ka
	Email: perumal.iit@gmail.com, d.perumal@iitg.ernet.in Tel: + 91 91598 60535
	<u>Google Scholar Profile</u> = https://scholar.google.co.in/citation <u>ResearchGate Profile</u> = https://www.researchgate.net/profile
E>	ecutive Editors[A & R]
1	Dr. Fazlul Huq, Head Cancer Research Group, School of Medical Sciences, Fac
2	Dr.Pavel Mokrejs, Thomas Bata University, The Czech Republic, Europe.
3	Dr. Adli Abdallah Hanna, National Research Centre, Egypt.
4	Dr. P.Y.Shirodkar, University of Mumbai,India.
41	


.

5	Dr. M. Al-Anber,
+	Mu'tah University, Jordan.
6	Dr.Saeid Amani,
-	Arak University,Iran .
7	Dr.Mitu Liviu,
	University of Pitesti, Romania.
	Dr. Nurfina Aznam,
9	University Negeri Yogyakarta, Indonesia.
	Dr. Tomasz Borowski, Centre Lab of Chem. Tech., Poland.
+	Dr. Mas Rosemal Hakim bin Mas Haris,
10	School of Chemical Sciences Universiti Sains Penang, Malaysia
+	Dr.Alamdar Ashnagar,
11	Pasteur Institute of Iran.
-	Dr. Abdulkareem Ismaiel Abdulraheem,
12	College of Health Sc., PAAET, Kuwait.
-	Dr. Ji-Tai Li,
13	Hebei University, China.
+	Dr.Abdalla Shalaby,
14	Zigzag University, Egypt
+	Dr. Georgia Koliopoulou,
15	University of Athens, Athens, Greece.
-	Dr. A. A. Mottahedi ,
16	IROST,Iran
	Dr. Gwendoline Ee Cheng Lian,
17	University PutraMalaysia, Malaysia.
	Dr. J.V.Manwar,
18	GSPS.IOP, M.S.India.
	Dr.Miss Swastika Ganguly,
19	Birla Insti.of Tech. India.
20	Prof. Dr. Sbirna Simona-Liana.
20	University of Craiova, Romania.
	Dr Carolyn Coulter,
21	University of Otago, New Zealand.
22	Dr. Konstantinos M. Kasiotis,
22	B P Institute, Laboratory of Pesticides Toxicology, Athens, 1456
22	Dr. Arun HS Kumar.
23	University College Cork, Ireland.
24	Associate editors: Dr. M.Z. Kumbhare, M.S.India. Dr. S.R.Nir

Home | About Us | Objectives | Consultancy | Int J ChemTech Res | Int J PharmTech Res | Site Map | Contact Us

All Privacy & Copyrights reserved with Sai Scientific Communications.

Designed & Developed by :ISC email : easifinatim(et)gmail(dot)com



	Association between Left Ventricular Global Longitudinal Strain and Six Minute Walk Test Before an After Chemotherapy with Anthracycline in Breast Cancer Patients					
(38-43)	Theresia W. Siagian, Andre P. Ketaren, Kamal B Siregar, Harris Hasan, Abdullah A. Siregar, Andik Sitepu					
	Abstract	Full Text PDF	Full Text XML			
	DOI : http://dx.do	.org/10.20902/IJCTR.2018.1103	<u>06</u>			
	The influence of collagen membranes and tetracycline on the PDGF-BB expression and osteoblas amount in bone defects healing: Experimental study in mice					
(44-51)	Agus Susanto, Susi Susanah, B.P. Priosoeryanto, Mieke Hemiawati Satari					
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.doi.org/10.20902/IJCTR.2018.110307					
	The Performance of Asphalt Concrete Mixture and Fiber Reinforcement Using Seawater Immersio Model to Indirect Tensile Strength					
(52-63)	Andi Alifuddin, A	ntariksa, Pitojo Tri Juwono, La	ambang Basri Said			
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.do	i.org/10.20902/IJCTR.2018.1103	<u>08</u>			
	-	Profile and Histopathological I n- Induced Diabetic Rat	Evaluation of Phaseoleus vulga	ris L beans Ethanoli		
(64-74)	Dwi Rita Anggrai	ni, Tri Widyawati, Henny Sri V	Vahyuni, Ade Putra Fratama Si	naga		
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.do	i.org/10.20902/IJCTR.2018.1103	<u>09</u>			
		Harris Hip Score functional o I fixation (ORIF) and hemiarth	utcome proximal femur fractu roplasty	are fixation with ope		
(75-78)	Yohanes Augustin	nus, Chairiandi Siregar				
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.do	i.org/10.20902/IJCTR.2018.1103	10			
	Enhancement of Water Infiltration Character of Hydrophobic Soil by Crude Cell Enzyme Mediate Bio-Alleviation					
(79-85)	Sridhar S, Hariswaran S, Nandhini					
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.do	i.org/10.20902/IJCTR.2018.1103	<u>35</u>			
,	To improve performance of Internal combustion engine with an aid of neodymium-doped yttrium aluminium garnet Laser Ignition System					
(86-93)	G.Mahesh, P.Vignesh, S.Manikandan, D.Monesh Kumar					
	Abstract	Full Text PDF	Full Text XML			
	DOI: http://dx.do	i.org/10.20902/IJCTR.2018.1103	<u>36</u>			
(94-100)	Properties of Concrete using Manufactured Sand as Fine Aggregate					

	Abstract	Full Text PDF	Full Text XML	
	DOI: http://dx.doi.org	/10.20902/IJCTR.2018.11033	7	
	Cadmium sulphide in cell	corporated reduced Graph	ene oxide as counter electrode	for dye sensitized
(101-110)	Sandhya Murali, F. M	lichael Raj, S.Vigneswari, A	.Jeya Rajendran	
			E H E AND	
	Abstract	Full Text PDF	Full Text XML	
		<u>Full Text PDF</u> /10.20902/IJCTR.2018.11033		
	DOI : http://dx.doi.org	/10.20902/IJCTR.2018.11033		Commercially Acti
(111-122)	DOI : http://dx.doi.org Adsorption of Cd(II) Carbon from the aque	/10.20902/IJCTR.2018.11033	8 Pistachios Seed Shell and C	Commercially Acti
(111-122)	DOI : http://dx.doi.org Adsorption of Cd(II) Carbon from the aque	by Chemically Activated solutions	8 Pistachios Seed Shell and C	Commercially Acti
(111-122)	DOI : http://dx.doi.org Adsorption of Cd(II) Carbon from the aque Malar Selvi M.S., S.A <u>Abstract</u>	/10.20902/IJCTR.2018.11033 by Chemically Activated cous solutions mala Fatima Rani , R.J. Sav	8 Pistachios Seed Shell and C ritha <u>Full Text XML</u>	Commercially Acti
(111-122)	DOI : http://dx.doi.org Adsorption of Cd(II) Carbon from the aque Malar Selvi M.S., S.A <u>Abstract</u>	/10.20902/IJCTR.2018.11033 by Chemically Activated cous solutions mala Fatima Rani , R.J. Sav Full Text PDF	8 Pistachios Seed Shell and C ritha <u>Full Text XML</u>	Commercially Acti





International Journal of ChemTech Research CODEN (USA): IJCRGG, ISSN: 0974-4290, ISSN(Online):2455-9555 Vol.11 No.03, pp 13-18, 2018

The Addition of L-Arginine in Capacitation Media to Motility, Viability, and Spermatozoa Capacity of Goats

Tri Wahyu Suprayogi¹, Rimayanti¹, Tatik Hernawati¹, Suharsono², Suherni Susilowati^{1*}

¹Department of Veterinary Reproduction, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

²Department of Anatomy, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia

Abstract : The purpose of this study is improving the provision of animal origin food of in vitro fertilization technology. Specific targets to be achieved in this study is to increase the incidence of capacitation spermatozoan and success in in vitro fertilization process . This research utilized four treatment groups, namely controlled group (P0) is the addition of LArginine in spermatozoa (HEPES) capacitance medium , group (P1) is the addition of LArginine in spermatozoa (HEPES) capacitation medium by 0,002 M / ml ,group (P1I) is the addition of L-Arginine in spermatozoa (HEPES) capacitation medium by 0,004 M / ml. and group (P1II) is the addition of L-Arginine in spermatozoa (HEPES) capacitation reaction medium by 0,006 M / ml Further evaluations included motility, viability, capacitation reaction of spermatozoa. Data obtained in the form of percentage of motility, viability and capacitation reaction is analyzed by F Test. The conclusion of this study, group (P1II) is the addition of LArginine in spermatozoa (HEPES) capacitation medium by 0,006 M / ml increase of motility, viability and capacitation reaction is analyzed by F Test. The conclusion of this study, group (P1II) is the addition of LArginine in spermatozoa (HEPES) capacitation medium by 0,006 M / ml increase of motility, viability and capacitation reaction.

Keywords : L-Arginine, capacitation medium, motility, viability and capacitation reaction.

1.Introduction

Various technologies have been created and have been used to improve the efficiency of livestock reproduction. Artificial Insemination is the initial use of Reproductive Biotechnology in male animals and then followed by research on in vitro manipulation of spermatozoa. How to manipulate spermatozoa in vitro include the method of centrifugation. In the process of in vitro fertilization requires the presence of a mature ovum and spermatozoa who have experienced maturity or capacitation. The maturation process of spermatozoa in invitro can be achieved by adding the appropriate medium⁸.

The adverse effects of seminal plasma splitting with centrifugation techniques are the increased formation of Reactive Oxygen Species (ROS) by spermatozoa¹¹. The increased accumulation of ROS production in centrifuged spermatozoa is thought to be a complex process and can be derived from various chemical processes, organelles or cells even from outside the cell². If the production of ROS is excessive and

International Journal of ChemTech Research, 2018,11(03): 13-18

DOI : http://dx.doi.org/10.20902/IJCTR.2018.110303

Suherni Susilowati et al /International Journal of ChemTech Research, 2018,11(03): 13-18. 14

unable to be neutralized by the antioxidant defense system present in the spermatozoa or seminal plasma can cause damage poly unsaturated fatty acid which is an important component of the phospholipid membrane membrane spermatozoa cells, the inactivation of glycolytic enzymes, motility and spermatozoa³. According to¹, L-Arginine is an amino acid that plays an important role in stimulating the motility of spermatozoa in mammals under in vitro conditions. L-Arginine also acts as an antioxidant that produces nitric oxide from enzyme synthesis reactions thus reducing peroxidation of lipid membrane spermatozoa that occurs due to free radicals when spermatozoa interact directly with oxygen. Nitric oxide is a biological molecule that plays an important role in physiological spermatozoa such as spermatozoa motility, spermatozoa-ovum interaction and spermatogenesis. Nitric oxide also plays a role in the defense mechanism of spermatozoa against the formation of reactive oxygen under the freezing point in liquid nitrogen at -196°C, which maintains sperm motility and post thawing viability.

According to⁶, L-Arginine is a substrate that produces nitric oxide with an intermediate enzyme nitrate oxide synthase. The enzym is in the acrosome section and the spermatozoa tail. L-Arginine stimulates capacitation and spermatozoa acrosome reaction¹¹. At low concentration, NO improve sperm motility. Beside promoting sperm motility, NO is also know to enhance capacitation and acrosom reaction in mouse and human spermatozoa ^{11;13}. However, higher concentrations of L-Arginin can have adverse effect on motility and fertility of human¹⁴ and rat ²³.Research on L-Arginine has been done, but research on the addition of L-Arginine in goat spermatozoa media has not been studied.

2. Materials and Methods

2. 1.Experimental animal

Semen is collected from three male goats by using an artificial vagina. The average body weight was 45 kg and average age was 3-4 years. An artificial vagina is prepared by placing both shrouded and sterilized containers, while the space between the outer and inner sheath is filled with warm water at 45 ° C in order to give the temperature to the inner sheath of 42-43 °C and the front third of the envelope in the vaginal artificial Vaseline smeared . After the artificial vagina is prepared, the male is given a stimulus with the angler females then carried out semen storage. Immediately after the shelter, the semen was taken by a laboratory to be separated from the spermatozoa and the plastics were used to be added to the goat semen diluent. The goat semen is accommodated with an artificial vagina and examined macroscopically and microscopically. Macroscopic examination includes volume, color, odor, consistency and pH and microscopic examination including mass movement, individual movement, viability, concentration and test resistance.

2.2. L-Arginine application on spermatozoa capacitation media

Good quality semen (percentages of spermatozoa motility and viability should be 70% or above), as much as 0.5 ml plus HEPES medium 1 ml and centrifugation at 1800 rpm for 10 minutes. Calculated $3x10^6$ spermatozoa for treatment.Spermatozoa of each centrifugation divided by 4 groups.Tube I filled with spermatozoa and HEPES medium, tube II filled with spermatozoa + HEPES medium + L-Arginin 0.002 M / ml HEPES. Tubes III filled with spermatozoa + HEPES medium + L-Arginin 0.004 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES. Tubes IV filled with spermatozoa + HEPES medium + L-Arginin 0.006 M / ml HEPES.

2.3. Assesment of motility

The motility of spermatozoa was analyzed by mixing the semen gently and placing a 10μ l drop of diluted semen on warm slide covered with a glass cover slip (18 x 18 mm) from five selected representative fields. One hundred spermatozoa were evaluated in at least five different fields in each smear under a light microscope. Individual sperm were recorded as being motil or non motil²¹.

2.4. Sperm viability

Eosin – negrosin staining was used to evaluate sperm viability as described by²¹. One drops of semen was placed on a tempered glass slide and this samples was mixex with one drop of eosin negrosin solution. The mixture was smeared on the glass slide and allowed to air dry. One hundred spermatozoa were evaluated in at least five different fields in each smear under a light microscope. Eosin penetrates non viable, non stained cells ²¹.

2.5. Assessment of sperm capacitation with Tetracycline Chlorine Staining

Examination of sperm capacitation with ChlorTetracyclin (CTC) staining. Preparations are examined under a fluorescent microscope with a magnification of 400x. The visible picture is: the whole head of the spermatozoa is fluorescent, the spermatozoa are non-capacitated. The upper half spermatozoa head fluorescent is a spermatozoa that undergoes capacitation.

2.6. Design and Statistical Analysis

The research design used is complete randomized design and data analisis done with Anova Test, if there is difference followed by Duncan Test ¹⁷.

3. Result

3.1.Collection of Goat semen for Treatment

This research requires fresh semen of goats that are first examined in macroscopic and microscopic. The macroscopic examination includes: volume, color, odor, consistency and degree of acidity or pH. Microscopic examination includes mass movement, individual movement, concentration and survival (viability) of spermatozoa. The results of the macroscopic and microscopic examination of goat's semen (Table 1). **Table 1.** Macroscopic and microscopic examination of fresh goat semen

Indicator	Character
Colour	White-yellowish
Smell	Typical
Consistency pH	Thick
Volume (ml)	7,00
Conceentration (juta) spz/mm ³	2,5±0,35
Mass Motility	3985x10 ⁶
Individual Motility (%)	+++
Viability (%)	Progressive (87±3,50)
Whole Plasma Membrane (%)	93±2,45
	85±1,55

The results showed that the goat semen that was accommodated was yellowish white, typical smell, thick consistency, pH \pm 7, volume 2.5 \pm 0.35 ml, concentration 3985 \pm 106, mass motility +++ (movement of large and large waves), individual motility progressive move 87 \pm 3.50%, viability 93 \pm 2.45 and intact plasma membrane 85 \pm 1.55%. In general, the volume of semen will increase according to age, body size, changes in state, reproductive health and frequency of semen shelter. The color, consistency and concentration of spermatozoa are closely related to each other. The more dilute a semen the concentration of spermatozoa will be lower and the color of the semen is getting pale. While the consistency of semen depends on the ratio of spermatozoa and plasma semen ⁴. The degree of acidity (pH) greatly affects the life-force of spermatozoa. When a high or low pH will cause the spermatozoa die. The degree of acidity of semen is influenced by the concentration of lactic acid produced in the metabolic process. According⁸, spermatozoa metabolism in anaerobic state will produce lactic acid buried and increase or decrease the pH of semen.

3.2 Examination of motility, viability and capacitation of goat sperm in HEPES + L-Arginine media

Individual motility was observed using a 400 times magnification light microscope. Observations were made on the number of progressive motile spermatozoa and spermatozoa velocity from one field of view and expressed in percent can be seen in table 2.

Observation of spermatozoa viability was done using a light microscope with 400 times magnification. The percentage of viability of spermatozoa can be seen in table 2. The live spermatozoa are characterized by clear or transparent colored spermatozoa heads while the dead spermatozoa are characterized by pink spermatozoa heads.

Suherni Susilowati et al /International Journal of ChemTech Research, 2018,11(03): 13-18. 16

The observed capacities and acrosome reactions were performed by Chlortetracyclin (CTC) staining and were observed with a magnification fluorescence microscope 400 times. The visible picture is: the whole head of the spermatozoa is fluorescent, the spermatozoa are not capacitated. The upper half-fluorescent spermatozoa head is a spermatozoa that has matured or capacitated. Observations were made from one field of view and expressed in percentages can be seen in table 2.

Table 2.Percentage of motility, viability and capacitation of goat spermatozoa in HEPES-added
LArginine media.

Parameter	HEPES	HEPES + 0,002	HEPES + 0,004	HEPES + 0,006
		M/mL L-Arginin	M/mL L-Arginin	M/mL L-Arginin
Motility P (%)	50,10±2,55 ^a	35,05±2,25 ^b	56,75±1,50 ^a	68,35±3,05°
Viability (%)	58,35±2,30 ^a	44,55±1,05 ^b	65,75±2,50 ^a	72,25±1,75°
Capacitation (%)	54,45±0,75 ^a	37,45±2,25 ^b	60,45±0,80 ^a	82,55±0,85°

Different letter notations on the same line are significantly different (p < 0.05)

In Table 2 it can be seen that the addition of L-Arginine in the capacitation medium (HEPES) yields a distinct pattern of difference in progressive motility, viability and spermatozoa capacitance. Motility, viability and spermatozoa capacitation in HEPES + medium 0.002 M / mL L-Arginine showed the lowest percentage. Between medium HEPES and HEPES + 0,004 M / ml L-Arginine there was no difference, meanwhile the highest on medium HEPES + 0,006 M / ml L-Arginin.UjiMultivariant to motility, viability and spermatozoa capacitation there was significant difference p<0,05 between the treatments.

4. Discussion

The movement of spermatozoa can occur because of the energy (ATP) that is product by mitochondria and the existence motor dynein (Cytoskeleton) for flagela or tail of sperm. The movement organized by Ca²⁺ (Cyclic Adenosin Mono Phosphat)²². Absolute spermatozoa survival require the integrity of the membrane. During the centrifugation process in addition to the antioxidant present in the seminal plasma disappeare there is also membrane molecular materials resulting in membrane destabilization thereby decreacing the physiological function of the membrane under normal condition, living cells need a source of energy formed from ATP through oxidative fosforylation requiring oxygen in process of reduction of oxygen in the transfer 4 electron. Inextain situation this transfer of 4 electron is running less than perfect, so that the reactive oxygen compound is very dangerous and damage the cells if not muted. The reaction oxygen compound is power full oxidant than can damage cell components that are essential to maintain cell integrity in live ²⁰.

Capacitation showed a correlation with change in fluidity of plasma membrane spermatozoa, intracellular ion concentration, metabolism and motility²². Molecularly during capacitation there are modifications of calcium intracellular ions and other ions, changes in lipid and phospholipid membrane spermatozoa, changes in protein phosphorylaion and activity of protein kinase. Capacitation is the rearrangement of biochemical membrane spermatozoa where as capacitance increases in intracellular Ca^{2+} level ¹⁵. The increasing of Ca^{2+} activity activates spermatozoa (hyperactivation).

The results showed that the highest percentage of motility, viability and capacitance of spermatozoa in HEPES + 0,006M / ml L-Arginine. The statement is appropriate with the research ⁶, which writing L-Arginine can acts as an antioxidant that protects spermatozoa from free radicals and lipid peroxidation so that the spermatozoa plasma membrane remains intact. In addition, because the dose of L-Arginine given in the medium of capacitance is to balance the occurrence of lipid peroxidation or free radicals.

L-Arginine is an amino acid that can stimulate the motility of mammalian spermatozoa under in vitro conditions and play an important role in the defense of cellular immunity ¹⁹. L -Arginin protects spermatozoa against lipid peroxidation by increasing the production of nitric oxide and deactivating free radicals. Nitric oxide has been shown to be an antioxidant and beneficial to reactive oxygen species (ROS) such as hydrogen peroxide (H202) and superoxide anion (O2) ¹. The mechanism of L-Arginin action is through the production of NO in the presence of NOS, and it was confirmed by the inhibition of the enzyme action as well as by scavenging the NO free radical. A similar role of NO has been demonstrated in human spermatozoa where this ROS has been shown to trigger tyrosine phosphorylation and double serine/threonine phosphorylation (Herrero et al, 2000) both events

being closely realted with capacitation¹⁰. Moreover, ⁹and ⁷mentioned that the arginin acts as an antioxidant which protects spermatozoa against lipid peroxidation during storage through increasing nitric oxide production which reduces lipid peroxidation by activating free radicals.

L-Arginine is a secondary antioxidant that works by chopping off a chain-oxidation reaction from free radicals or by capturing free radicals. L-Arginine can also increase spermatooza glycolysis. The process of glycolysis can produce energy in the form of adenosine triphosphate (ATP), which is used by spermatozoa as a source of energy in the process of movement so that it can remain motile and maintain its life force. Futhermore, ⁵reported that the nitric oxide have a major role in stimulation of lactic dehydrogenize which was a key for metabolic process in sperm motility that may be presumably given as the sperm direct effects responses individual motility.

The conclusion of this study, group (PIII) is the addition of L-Arginine in spermatozoa (HEPES) capacitation medium by 0,006 M / m increase of motility, viability and capacitation reaction.

Acknowledgement

This research was supported by funding from the Research and Technology, and Directorate General of Higher Education Ministry (RISTEK-DIKTI) (2017). The authors are also grateful to all of the laboratory staff in the Department of Reproduction, Faculty of Veterinary Medicine Airlangga University.

References

- 1. Al-Ebady,AS., SO. Hussain, KI.,Al-Badry and FF. Ibrahim.2012. Effect of Adding L-Arginin on Some Parameters of Bull Sperms After Freezing in Liquid Nitrogen (-1960C). Al-Qadisiya Journal of Vet Med Sci. 11(2):15-161.
- 2. Agaerwal, R; R.A Saleh and M.A Bedalwy, 2003. Role of Reactive Oxygen Species, Lipid Peroxidation and Human Sperm Function. Biology Reprod. 40: 183-197.
- 3. Alvarez, J.G and B.T, Storey, 1995. Differential Incorporation of Fatty Acid into and Peroxidative Loss of Fatty Acid From Phospholipid of Human Spermatozoa. Mol. Reprod. Dev. 42 : 334-345. 4. Evans,G and M.W.C.Maxwell, 1988. Salomon's Artificial Insemination of Sheep and Goat. Butherworths, Sydney.
- 5. Clancy RM; Levatovsky D; Leszczynska-Piziak J. Yegudin J, Abramson SB, 1994. Nitric Oxide Reacts with Intracellular Gtutathion and Activates the Hexose Monophosphate Shunt in Human Neutrophils, Evidence for S-nitro so Glutathione as a Bioactve Intermediary. Proc Natl. Acad. Sci. USA. 91: 36803684.
- 6. Flaherty, C.O; Pablo R and SudhaSrivastava.(2004). L-Arginin Promotes Capacitation and Acrosom Reaction in Cryopreserved Bovine Spermatozoa. Biochimia at BiophysicaActa. 1674: 215-221.
- 7. Govil G; Phadke RS, Srivastava S, 1992. Physical/chemical studies of vitamin E in membranes. In Ong ASH, Packer L (eds). Lipid Soluble Antioxidant. Biochemistry and Clinical Applications. Birkhautser Verlag. Basel, Switzerland. pp.27-46.
- 8. Hardjopranjoto, 2006. Perkembangan Bioteknologi Reproduksi Pada Ternak. Pidato Ilmiah. Pada Acara TemuIlmiahSehariDalamRangka Purnabakti Prof. Dr. Soehartojo Hardjopranjoto MSc,Drh. Dari Fakultas Kedokteran Hewan Universitas Airlangga.
- 9. Hogg N, Kalyanaraman B ,1999. Nitric Oxide and Lipid Peroxide Ion. Biochem.Biophys.Acta1411 : 378-384.
- 10. Thindathil, J; E. de Lamirande, C Gagnon 2003. Nitric Oxide Regulates the phosphorylation of The Threonine-Glutamine-Tyrosin motif in proteins of human spermatozoa during capacittion. Biol. Reprod. 68:1291-1298.
- 11. Herrero, M.B; S, Chaterjou, Letfieve, E. de Lamirande, C.Gagnon, 2000. Nitric Oxide Interacts with The cAMP pathway to Modulate capacitation of Human Spermatozoa. Free Radic. Biol.Mol. 29:522-536.
- 12. Laughlin, Mac; E. A and W.C.L Ford, 1994. Adenosin Triphosphate and Motility Characteristics of Fresh and Cryopreservation of Human Spermatozoa. Int. J.Adrol. 17:19-23. Frequencies of Fertilization. Theriogenology. 23: 216.
- Revelli, A; C. Costamagna, F. Moffa, E. Aldieri, S Ochesti, A Bosia, M Massobrio, B.Lindbiom, D. Ghigo, 2001. Signaling Pathway of Nitric –Oxide- Induces Acrosom Reaction in Human Spermatozoa. Biol. Reprod. 64: 1708-1712.
- 14. Roselli, M; R.K. Dubey, B, Imthum, E, Macas, P. Keller, 1995. Effects of Nitric Oxide on Human Spermatozoa : Evidence that Nitric Oxide Decreases Sperm Motility and Induce Sperm Toxicity . Human Reprod. 10: 1786-1790.

- 15. Parrish, J.J; J.J. Susko Parish and N.L.Fist, 1985. In Vitro Fertilization of Bovine Oocytes Using Heparin Treated and Swim Up Separated Frozen – Thawed Bovine Semen is Repeatable and Result In High
- 16. Palmer, R.M.J; D.S; Ashton S, Moncada(1988). Vascular Endothelial Cell Synthesize Nitric Oxide from L-Arginin. Nature 33:664—668.
- 17. Santoso, S dan Fandy, T, 2001. Riset Pemasaran, Konsep dan Aplikasi dengan SPSS. PT Gramedia . Jakarta.
- 18. Salisbury, G.W and N.L VanDemark, 1985. FisiologiReproduksi dan Inseminasi Buatan pada Sapi. Terjemahan R. Djanuar. Gadjah Mada UniversityPress. Yogyakarta.
- 19. Srivastava, S., Prashant D., Evans C., Girjesh G. 2006. Mechanism of Action of L-Argininon The Vitality of Spermatozoais Primarily Through Increased Byosynthesis of Nitric Oxide. Biology of Reproduction 74: 954-958.
- 20. Suryohudoyo, P, 2000., Ilmu Kedokteran Molekuler. Cetakan Pertama. Jakarta. CV. Sagung Seto. 3147.
- 21. Susilowati, S; Hardiyanto; TriWahyu Suprayogi; Trilas Sardjito; TatikHernawati , 2010. Penuntun Praktikum Inseminasi Buatan. Airlangga UniversityPress.
- 22. Visconti, P.E; G.D. Moore: J.L. Bailey; P.Leciere; S.A. Connors; D.P.P Old Clarke and G.S. Kopf, 1995. Capacitation of Mouse Spermatozoa Division of Reproductive Biology Departement of Obstretics and Gynecology. University of Pennylvania School of Medicine. Philadelphia. Reprod. 5: 234-245
- 23. Ratnassooriya, W.D; M.G. Dharmasiri, 2001. L-Argininthe Substrate of Nitric Oxide Synthase ,Inhibits Fertility of MaleRats. Asian Journal Androl. 3 : 97-103.
- 24. Yanagimachi, R, 1994. Mammalian Fertilization. In.knobil, E. Nell, JD (eds). ThePhysiology of Reproduction. New York. Raven Press. Ltd. 189-318.
