

## DAFTAR PUSTAKA

- Afifi, S. A. *et al.* (2015) ‘Nanosuspension: An emerging trend for bioavailability enhancement of etodolac’, **International Journal of Polymer Science**, 2015. doi: 10.1155/2015/938594.
- Allen, L. V and Ansel, H. C. (2011) ‘**Ansel’s Pharmaceutical Dosage Forms and Drug Delivery Systems**’, p. 537. doi: 978-0-7817-7934-0.
- American Pharmaceutical Assosiation, 2018. **The United States Pharmacopeia/ The National Formulary (USP41/NF36) Volume 2.** Rockville : The United States Pharmacopeial Convention, Inc.
- Ansel, H. C., Allen, L. V., and Popovich, N. G., 2005, **Ansel’s Pharmaceutical Dosage Form and Drug Delivery Systems, Eight Edition**, 230, 239-241, Lippincott Williams & Wilkins a wotters Kluver Company, Philadelphia.
- Article, R., Jaspreet, K. and Sandeep, K. (2018) ‘**Development Of Nanocrystal Formulation With Improved Dissolution**’, 8(5), pp. 118–129.
- Ashari, A. (2018) **Pengaruh Penambahan PVP K-30 Terhadap Karakteristik Fisik Tablet Hasil Solidifikasi Nanosuspensi Hesperetin-Poloksamer 407 yang Dibuat dengan Metode Fluid Bed Granulation.** Departemen Farmasetika. Universitas Airlangga (Skripsi)
- Aulton, M.E., Taylor, K.M.G., 2013, **Aulton’s Pharmaceutics: The Design and Manufacture of Medicines**, Fourth Edition, Churcihill Livingstone Elsevier, 465-476

- Bandelin, F.J., 1989. Compressed Tablets by Wet Granulation, In: Lachman, L., Lieberman, H.A., Schwartz, J.B., **Pharmaceutical Dosage Form : Tablets**, Vol.1, New York : Marcell Dekker, Inc, p.205-209.
- Bernard Van Eerdenbrugh, Jan Vermant, Johan A. Martens, Ludo Froyen, Jan Van Humbeeck, Patrick Augustijns, G. V. D. M. (2009) 'A Screening Study of Surface Stabilization during the Production of Drug Nanocrystals', **JOURNAL OF PHARMACEUTICAL SCIENCES**, 98(10), pp. 2091–210.
- Bhakay, A. *et al.* (2018) 'Bioavailability enhancement of poorly water-soluble drugs via nanocomposites: Formulation–Processing aspects and challenges', **Pharmaceutics**, 10(3). doi: 10.3390/pharmaceutics10030086.
- Bose, S. *et al.* (2012) 'Application of spray granulation for conversion of a nanosuspension into a dry powder form', **European Journal of Pharmaceutical Sciences**. Elsevier B.V., 47(1), pp. 35–43. doi: 10.1016/j.ejps.2012.04.020.
- Budavari, S., 1996. **The Merck Index**. Edisi 12. WhiteHouse USA: Merck & Co.Inc.
- Cantor, S., Augsburger, L. and Gerhardt, A. (2008) 'Pharmaceutical Granulation Processes, Mechanism and the Use of Binders', **Pharmaceutical Dosage Forms - Tablets, Third Edition**, (June), pp. 261–301. doi: 10.1201/b15115-9.
- Carstensen, J.T., 1977. **Pharmaceutical of Solids and Solid Dosage Form**.New York : John Wiley and Sonic, p.133.
- Chaudhari, S. P. and Dugar, R. P. (2017) 'Application of surfactants in solid dispersion technology for improving solubility of poorly water

- soluble drugs', **Journal of Drug Delivery Science and Technology**. Elsevier Ltd, 41(June), pp. 68–77. doi: 10.1016/j.jddst.2017.06.010.
- Choudhary, O. P. and ka, P. (2017) 'Scanning Electron Microscope: Advantages and Disadvantages in Imaging Components', **International Journal of Current Microbiology and Applied Sciences**, 6(5), pp. 1877–1882. doi: 10.20546/ijcmas.2017.605.207.
- DFE pharma (2012) 'Introduction to tableting by wet granulation', **DMV-Fonterra Excipients**, pp. 1–10. doi: 10.1111/j.1399-6576.2011.02426.x.
- DM Parikh (2016) **Handbook of pharmaceutical granulation technology**.
- Figueroa, C. E. and Bose, S. (2013) 'Spray granulation: Importance of process parameters on in vitro and in vivo behavior of dried nanosuspensions', **European Journal of Pharmaceutics and Biopharmaceutics**. Elsevier B.V., 85(3 PART B), pp. 1046–1055. doi: 10.1016/j.ejpb.2013.07.015.
- Gülsün, T., Gürsoy, R. N. and Öner, L. (2009) 'Nanocrystal technology for oral delivery of poorly water-soluble drugs', **Fabad Journal of Pharmaceutical Sciences**, 34(1), pp. 55–65.
- Hadisoewignyo, L. and Fudholi, A., 2013. **Sediaan Solida**. Yogyakarta : Pustaka Pelajar
- Junyaprasert, V. B. and Morakul, B. (2015) 'Nanocrystals for enhancement of oral bioavailability of poorly water-soluble drugs', **Asian Journal of Pharmaceutical Sciences**. Elsevier Ltd, 10(1), pp. 13–23. doi: 10.1016/j.ajps.2014.08.005.
- Kanaze, F. I. *et al.* (2006) 'Thermal analysis study of flavonoid solid

- dispersions having enhanced solubility', **Journal of Thermal Analysis and Calorimetry**, 83(2), pp. 283–290. doi: 10.1007/s10973-005-6989-9.
- Kannuri, R., Chamarthi, H., Kumar, S.M., Challa, T., Goud, A.B. 2011. Formulation Development and In Vitro Evaluation of Escitalopram Oxalate Orally Disintegrating Tablets. **Internasional Journal of Pharmaceutical, Chemical and Biological Sciences**. Vol 1(1): 57-65
- Kementerian Kesehatan RI. 2014. **Famakope Indonesia Edisi V**. Jakarta: Kementerian Kesehatan RI. Hal 52
- Lachman, L., Lieberman, Lee, J. and Cheng, Y. (2006) „Critical freezing rate in freeze drying nanocrystal dispersions“, 111, pp. 185–192.
- Lestari, M. L. A. D., Müller, R. H. and Möschwitzer, J. P. (2015) ‘Systematic screening of different surface modifiers for the production of physically stable nanosuspensions’, **Journal of Pharmaceutical Sciences**, 104(3), pp. 1128–1140. doi: 10.1002/jps.24266.
- Liu, P. et al. (2011) „Nanosuspensions of poorly soluble drugs : Preparation and development by wet milling“, **International Journal of Pharmaceutics**. Elsevier B.V., 411(1–2), pp. 215–222.
- Merisko-Liversidge, E., Liversidge, G. G. and Cooper, E. R. (2003) ‘Nanosizing: A formulation approach for poorly-water-soluble compounds’, **European Journal of Pharmaceutical Sciences**, 18(2), pp. 113–120. doi: 10.1016/S0928-0987(02)00251-8.
- Mishra, P. R. et al. (2009) „Production and characterization of Hesperetin nanosuspensions for dermal delivery“, **International Journal of Pharmaceutics**, 371(1–2), pp. 182–189.

- Morin, G. and Briens, L. (2014) 'A Comparison of Granules Produced by High-Shear and Fluidized-Bed Granulation Methods', **AAPS PharmSciTech**, 15(4), pp. 1039–1048. doi: 10.1208/s12249-014-0134-7.
- Oulahna, D. *et al.* (2003) 'Wet granulation: The effect of shear on granule properties', **Powder Technology**, 130(1–3), pp. 238–246. doi: 10.1016/S0032-5910(02)00272-3.
- Pande, V. V. and Abhale, V. N. (2016) 'Nanocrystal technology: A particle engineering formulation strategy for the poorly water soluble drugs', **Der Pharmacia Lettre**, 8(5), pp. 384–392. doi: 10.1016/j.ijpharm.2013.01.019.
- Parrot, E. (1970). **Pharmaceutical Technology Fundamental Pharmaceutics**. Burgess Publishing Company. United States of America.
- Peltonen, L. and Hirvonen, J. (2010) 'Pharmaceutical nanocrystals by nanomilling: Critical process parameters, particle fracturing and stabilization methods', **Journal of Pharmacy and Pharmacology**, 62(11), pp. 1569–1579. doi: 10.1111/j.2042-7158.2010.01022.x.
- REDC/ADF (2018) 'Dissolution Testing and Acceptance Criteria for Immediate-Release Solid Oral Dosage Form Drug Products Containing High Solubility Drug Substances Guidance for Industry', **FDA**.
- Rowe, R., Sheskey, P. and Quinn, M. (2009) „Handbook of Pharmaceutical Excipients“, **Handbook of pharmaceutical excipients, Sixth edition**, pp. 549–553.
- Rudnic, E.M., and Schwartz, J.D., 2000. Oral Solid Dosage Form. In: Gennaro, A.R., **Remington : The Science and Practice of**

- Pharmacy**, 20th Eddition, Vol. 1, Philadelphia: Lippincott Williams & Wilkins, p.860-861, 868, 1625.
- Salazar, J., Müller, R. H. and Möschwitzer, J. P. (2013) ‘Application of the combinative particle size reduction technology H 42 to produce fast dissolving glibenclamide tablets’, **European Journal of Pharmaceutical Sciences**. Elsevier B.V., 49(4), pp. 565–577. doi: 10.1016/j.ejps.2013.04.003.
- Singare, D. S., Kulkarni, G. T. and Gowthamrajan, K. (2013) ‘Nanosuspension solidification technique: Evaluation of high shear granulation as per industrial perspective’, **International Journal of Pharma and Bio Sciences**, 4(3).
- Srirangam, R. and Majumdar, S. (2010) „Passive asymmetric transport of hesperetin across isolated rabbit cornea”, **International Journal of Pharmaceutics**. Elsevier B.V., 394(1–2), pp. 60–67.
- Steeden, S. *et al.* (1994) ‘Flavonoids Antioxidants’, **J. Am. Chem. Soc**, (5), pp. 4846–4851.
- Thadkala, K., Sailu, C. and Aukunuru, J. (2015) ‘Formulation, optimization and evaluation of oral nanosuspension tablets of nebivolol hydrochloride for enhancement of dissolution rate’, **Der Pharmacia Lettre**, 7(3), pp. 71–84.
- Tommasini, S. *et al.* (2004) ‘Improvement in solubility and dissolution rate of flavonoids by complexation with β-cyclodextrin’, **Journal of Pharmaceutical and Biomedical Analysis**, 35(2), pp. 379–387. doi: 10.1016/S0731-7085(03)00647-2.
- Tsai, Y. *et al.* (2010) „In vitro permeation and in vivo whitening effect of topical hesperetin microemulsion delivery system”, 388, pp. 257–262.