

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12 H-pyrano[2,3-a]xanthen-12-one from
Calophyllum pseudomole

by Mulyadi Tanjung

Submission date: 08-May-2018 06:42PM (UTC+0800)

Submission ID: 960735515

File name: Molbank2017-M961.pdf (254.25K)

Word count: 2563

Character count: 13286

Short Note

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one from *Calophyllum pseudomole*

¹⁰ Mulyadi Tanjung *, Ratih Dewi Saputri and Tjitjik Srie Tjahjandarie

Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya 60115, Indonesia; duffputri@gmail.com (R.D.S.); tjitjiktjahjandarie@fst.unair.ac.id (T.S.T.)

* Correspondence: mulyadi-t@fst.unair.ac.id; Tel.: +62-31-5936501; Fax: +62-31-5936502

¹⁸ Received: 1 September 2017; Accepted: 27 September 2017; Published: 30 September 2017

Abstract: A new pyranocoumarin, namely 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one **1**, was isolated from the stem bark of *Calophyllum pseudomole*. The structure of compound **1** was elucidated based on its ultraviolet (UV); infrared (IR); high resolution electro spray ionization mass spectrometry (HRESIMS); 1D and 2D nuclear magnetic resonance (NMR) spectral data.

Keywords: 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one; *Calophyllum pseudomole*; pyranoxanthone

²⁶ 1. Introduction

The genus *Calophyllum* belongs to the Clusiaceae family which comprises about 180 species found mainly in Southeast Asia. This plant is endemic to Kalimantan Island, Indonesia. This genus has been shown to possess a number of secondary metabolites such as xanthenes [1–3], coumarins [4,5], and chromanone acids [6,7]. Many of these compounds have shown a wide range of biological and pharmacological properties such as anti-HIV [8,9], anticancer [10,11], and antimalarial properties [12].




Figure 1. Structures of 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one.

²² In continuation of our phytochemical investigation on bioactive xanthone, we wish to report the isolation and structural elucidation of a new pyranoxanthone, 5,9,11-trihydroxy-

3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one (Figure 1) from the stem bark of *Calophyllum pseudomole*. The cytotoxicity against murine leukemia P-388 and antiplasmodial activity against *Plasmodium falciparum* of **1** are also briefly described.


2. Result and Discussion

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one was isolated as a yellow solid, and its UV absorption maxima at λ_{max} 245, 268, and 320 nm which is characteristic of xanthone chromophore [2]. Based on HRESIMS (see data supplementary Figure S1), compound **1** showed deprotonated molecule ion $[M-H]^-$ at m/z 477.1917 corresponding to the molecular formula $C_{28}H_{30}O_7$. The IR spectrum (data supplementary Figure S2) showed absorption for hydroxyl (3423 cm^{-1}), conjugated carbonyl (1645 cm^{-1}), and aromatic (1577 and 1460 cm^{-1}) groups [13]. The $^1\text{H-NMR}$ spectrum (Table 1, data supplementary Figure S3) of **1** showed the presence of a singlet of chelated hydroxyl at δ_{H} 13.82 (11-OH), a singlet of hydroxyl at δ_{H} 6.14 (5-OH), and two singlets of the isolated aromatic proton at δ_{H} 6.81 (H-6) and δ_{H} 6.39 (H-8) which is typical for a xanthone with six substituents [1]. The $^1\text{H-NMR}$ spectrum also revealed the presence of a substituent of a monosubstituted 2,2-dimethylpyrano, 2-methylbut-3-en-2-yl, and 2-hydroxy-3-methylbut-3-en-1-yl group. The presence of a monosubstituted 2,2-dimethylpyrano was determined from resonances of vinyl at δ_{H} 8.13 (s, 1H), and two signals of methyl at δ_{H} 1.54 (s, 3H) and 1.52 (s, 3H). The $^1\text{H-NMR}$ spectrum of 2-methylbut-3-en-2-yl showed the presence of a methyl at δ_{H} 1.43 (s, 6H), a vinyl at δ_{H} 5.98 (dd, 1H), and a methylene at δ_{H} 5.13 (d, 1H) and δ_{H} 5.08 (d, 1H). The $^1\text{H-NMR}$ spectrum of 2-methylbut-3-en-2-yl also showed the presence of a methyl at δ_{H} 1.87 (s, 3H), two proton signals of methylene at δ_{H} 3.16 (dd, 1H) and 2.93 (dd, 1H), a methyne at δ_{H} 4.43 (d, 1H), and two proton signals of vinylic methylene at δ_{H} 4.97 (s, 1H) and 4.86 (s, 1H).

The $^{13}\text{C-NMR}$ spectrum (Table 1, Figure S4 data supplementary) of **1** showed 28 carbon signals and their assignments were determined by heteronuclear multiple quantum coherence (HMQC) and heteronuclear multiple bond correlation (HMBC) spectra. The placement of hydroxyl, a monosubstituted 2,2-dimethylpyrano, 2-methylbut-3-en-2-yl, and 2-hydroxy-3-methylbut-3-en-1-yl group was confirmed by HMQC and HMBC spectra (Table 1). Based on the HMBC spectrum, a chelated hydroxyl signal at δ_{H} 13.82 (11-OH) was correlated with two quaternary carbons [δ_{C} 107.4 (C-10); 103.4 (C-11a)], and an oxyaryl carbon δ_{C} 161.1 (C-11) indicated that C-10 has a substituent. Two proton signals of methylene (H-1'') of the 2-hydroxy-3-methylbut-3-en-1-yl group at δ_{H} 3.18 and δ_{H} 2.95 showed long-range correlations with two oxyaryl carbons [δ_{C} 161.1; 163.2 (C-9)], two quaternary carbons [δ_{C} 107.4; 146.7 (C-3'')], and a methine carbon of alcohol [δ_{C} 77.5 (C-2'')]. The correlations showed that the 2-hydroxy-3-methylbut-3-en-1-yl group attached at C-10. The position of 2-hydroxy-3-methylbut-3-en-1-yl group at C-10 is reinforced by the correlation of the isolated aromatic proton signals at δ_{H} 6.39 (H-8) with two oxyaryl carbons [δ_{C} 155.8 (C-7a); 163.2 (C-9)], and two quaternary carbons [δ_{C} 107.4 (C-10); 103.4 (C-11a)]. Furthermore, the proton signal of isolated aromatic at δ_{H} 6.81 (H-6) correlates with two oxyaryl carbons [δ_{C} 153.4 (C-5), 150.6 (C-6a)] and a quaternary carbon signal (δ_{C} 108.5, C-4a), which showed that 2,2-dimethylpyrano was fused at C-4a and C-12b and the hydroxyl group attached at C-5 from the xanthone skeleton. The position of the hydroxyl group at C-5 is reinforced by the correlation of the hydroxyl proton signal at δ_{H} 6.15 (5-OH) with an oxyaryl carbon [δ_{C} 153.4 (C-5)], and a methine carbon [δ_{C} 101.8 (C-6)]. The methine signal of the vinyl group at δ_{H} 8.13 (H-4) of a monosubstituted 2,2-dimethylpyrano group showed long-range correlations with four quaternary carbons [δ_{C} 80.6 (C-2), 135.7 (C-3), 108.5 (C-4a), 42.1 (C-2'')] which showed that a methyl but-3-en-2-yl group attached at C-3. The HMBC correlation of compound **1** can be seen on Figure 2. Therefore, compound **1** was identified as 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one and a novel compound.

Table 1. NMR spectroscopic data of 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one in CDCl₃.

No.C	δ_H (Mult, J in Hz)	δ_C	HMBC
2	-	80.6	-
2a	1.54 (s, 3H)	27.4	C-2; C-2b; C-12b
2b	1.53 (s, 3H)	27.4	C-2; C-2a; C-12b
3	-	135.7	-
4	8.13 (s, 1H)	118.0	C-2; C-3; C-4a; C-2'
4a	-	108.5	-
5	-	150.6	-
6	6.81 (s, 1H)	101.8	C-3; C-4a; C-5; C-6a
6a	-	153.4	-
7a	-	155.8	-
8	6.39 (s, 1H)	94.5	C-7a; C-9; C-10; C-11a
9	-	163.2	-
10	-	107.4	-
11	-	161.1	-
11a	-	103.4	-
12	-	182.5	-
12a	-	121.7	-
12b	-	148.8	-
1'	1.43 (s, 3H)	28.3	C-2; C-3; 2'-CH ₃
2'	-	42.7	-
2'-CH ₃	1.43 (s, 3H)	28.2	C-1'; C-2'; C-3'
3'	5.98 (dd, 10.6; 17.6, 1H)	146.8	C-1'; C-2'; 2'-CH ₃
4'	5.08 (d, 10.6, 1H) 5.13 (d, 17.6, 1H)	112.0	C-2'; C-3'
1''	3.16 (d, 15.1, 1H) 2.93 (dd, 7.6; 15.1, 1H)	28.1	C-9; C-10; C-11; C-2''; C-3''
2''	4.42 (d, 7.3, 1H)	77.5	C-10; C-1''; C-3''; C-4''; C-5''
3''	-	146.7	-
4''	4.97 (s, 1H) 4.86 (s, 1H)	110.2	C-2''; C-5''
5''	1.87 (s, 3H)	18.7	C-2''; C-3''; C-4''
5-OH	6.15 (s, 1H)	-	C-4a; C-5; C-6
9-OH	9.15 (s, 1H)	-	-
11-OH	13.48 (s, 1H)	-	C-10; C-11; C-11a

**Figure 2.** Selected HMBC correlation for 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-a]xanthen-12-one.

The cytotoxic activity of compound **1** against murine leukemia P-388 cells, as determined by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay, showed IC₅₀ values of 4.10 µg/mL and was categorized as moderate activity. The antiplasmodial activity of **1** against *P. falciparum* showed IC₅₀ values of 2.88 µg/mL and was categorized as moderate activity.

3. Experiment Section

3.1. General

NMR spectra were recorded on a JEOL 400 ECA spectrophotometer (Tokyo, Japan) in CDCl₃ at 400 (¹H) and 100 (¹³C) MHz using tetramethyl silane (TMS) as the internal standard. The mass spectra were recorded using a Waters LCT Premier XE (Santa Clara, CA, USA). The UV spectra were measured with a Shimadzu series 1800 spectrophotometer (Kyoto, Japan). The IR spectra were recorded with Perkin-Elmer spectrum-100 FT-IR (Waltham, MA, USA). Column chromatography and planar radial chromatography were carried out using silica gel 60 G 1.07734.1000 and Si gel 60 PF₂₅₄ 1.07749.1000 (Merck, Darmstadt, Germany).

3.2. Plant Material

The stem bark of *C. Pseudomole* was collected in Sungai Mendawak, anak Sungai Kapuas, District Kubu Raya, Kalimantan, Indonesia on April 2015. The specimen was identified at the Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia.

3.3. Extraction and Isolation

The dried stem bark of *C. pseudomole* (3.0 kg) was macerated in MeOH twice for 2 days each time. After evaporation of the solvent in a rotary evaporator, 260 g of pale brown semi-solid was obtained. Further, the MeOH extract was partitioned first with *n*-hexane (1:1 *v/v*); water was added (10% *v/v*) in the second step to increase the polarity; then the MeOH extract was partitioned with EtOAc (1:1 *v/v*). The EtOAc extract (35 g) was subjected to column chromatography over silica gel and eluted with *n*-hexane-ethyl acetate (from 9:1 to 1:1) to give fractions A–C. Fraction C was then subjected further to column chromatography and eluted with *n*-hexane-EtOAc (from 9:1 to 1:1) to produce three subfractions C₁–C₃. Subfraction C₂ was purified by planar radial chromatography using *n*-hexane-CHCl₃ (from 3:7 to 7:3), CHCl₃ and CHCl₃-EtOAc 9:1 to yield compound 5,9,11-trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en-1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12H-pyrano[2,3-*a*]xanthen-12-one (18 mg).

3.4. Cytotoxic Assay

The cytotoxic properties of **1** against murine leukemia P-388 cells were evaluated according to the MTT method as previously described [14–16]. Artonin E was used as the positive control.

3.5. Antiplasmodial Assay

The antiplasmodial properties of **1** against *P. falciparum* were evaluated according to the Trager–Jensen method as previously described [17,18]. Chloroquine was used as the positive control.

Supplementary Materials: The following are available online at 1422-8599/2017/4/M961, IR, HRESIMS, 1D and 2D NMR spectra are reported in the supplementary materials as Figures S1–S6.

Acknowledgments: This research was supported by Universitas Airlangga, Ministry of Research, Technology and Higher Education, Republic of Indonesia (Penelitian Hibah Kompetensi, Universitas Airlangga, 2017).

Author Contributions: Mulyadi Tanjung designed the whole experiment of bioactivity and wrote the manuscript. Tjitjik Srie Tjahjandarie researched data, analyzed the NMR and HRESIMS spectra and contributed to the manuscript, Ratih Dewi Saputri designed the whole experiment. All authors read and approved the final manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Tanjung, M.; Saputri, R.D.; Tjahjandarie, T.S. 5,9,11-Trihydroxy-2,2-dimethyl-10-(3'-methyl-2'-butenyl)-3-(2''-methyl-3''-butenyl)pyrano[2,3-a]xanthen-12(2H)-one from the stem bark of *Calophyllum pseudomole*. *Molbank* **2016**, M906.
2. Tjahjandarie, T.S.; Saputri, R.D.; Tanjung, M. 5,9,11-Trihydroxy-2,2-dimethyl-3-(2-methylbut-3-en-2-yl)pyrano[2,3-a]xanthen-12(2H)-one from the stem bark of *Calophyllum tetrapterum* Miq. *Molbank* **2017**, M936.
3. Wei, D.J.; Mei, W.L.; Zhong, H.M.; Zeng, Y.B.; Wu, X.D.; Dai, H.F. A new prenylated xanthone from the branches of *Calophyllum inophyllum*. *J. Asian Nat. Prod. Res.* **2011**, *13*, 265–269.
4. Daud, S.B.; Ee, G.C.L.; Malek, E.A.; Teh, S.S.; See, I. A new coumarin from *Calophyllum hosei*. *Nat. Prod. Res.* **2014**, *28*, 1534–1538.
5. Joshi, S.P.; Kulkarni, S.R.; Phalgune, U.D.; Puranik, V.G. New dipyrano-coumarin from the leaves of *Calophyllum apetalum* Willd. *Nat. Prod. Res.* **2013**, *27*, 1896–1901.
6. Ha, L.D.; Hansen, P.E.; Duus, F.; Pham, H.D.; Nguyen, L.D. A new chromanone acid from the bark of *Calophyllum dryobalanoides*. *Phytochem. Lett.* **2012**, *5*, 287–291.
7. Lim, C.K.; Subramaniam, H.; Say, Y.H.; Jong, V.Y.M.; Khaledi, H.; Chee, C.F. A new chromanone acid from the stem bark of *Calophyllum teysmannii*. *Nat. Prod. Res.* **2015**, *29*, 1970–1977.
8. McKee, T.C.; Covington, C.D.; Fuller, R.W.; Bokesch, H.R.; Young, S.; Cardellina, J.H.; Kadushin, M.R.; Soejarto, D.D.; Stevens, P.F.; Cragg, G.M. Pyranocoumarins from tropical species of the genus *Calophyllum*: Chemotaxonomic study of extracts in the national cancer institute collection. *J. Nat. Prod.* **1998**, *61*, 1252–1256.
9. Reyes, M.H.; Basualdo, M.C.; Abe, F.; Estrada, M.J.; Soler, C.; Chilpa, R.R. HIV-1 inhibitory compounds from *Calophyllum brasiliense* leaves. *Biol. Pharm. Bull.* **2004**, *27*, 1471–1475.
10. Xiao, Q.; Zeng, Y.-B.; Mei, W.L.; Zhao, Y.-X.; Deng, Y.-Y.; Dai, H.-F. Cytotoxic prenylated xanthenes from *Calophyllum inophyllum*. *J. Asian Nat. Prod. Res.* **2008**, *10*, 993–997.
11. Mah, S.H.; Ee, G.C.L.; Teh, S.S.; Sukari, M.A. *Calophyllum inophyllum* and *Calophyllum soulattri* source of anti-proliferative xanthenes and their structure-activity relationships. *Nat. Prod. Res.* **2013**, *27*, 98–101.
12. Hay, A.; Helesbeux, J.; Duval, O.; Labaied, M.; Grellier, P.; Richomme, P. Antimalarial xanthenes from *Calophyllum caledonicum* and *Garcinia vieillardii*. *Life Sci.* **2004**, *75*, 3077–3085.
13. Tanjung, M.; Saputri, R.D.; Tjahjandarie, T.S. Antioxidant activity of two isomeric benzoxepinderivatives from the stem bark of *Bauhinia acuelata* L. *J. Chem. Pharm. Res.* **2014**, *6*, 705–708.
14. Tanjung, M.; Hakim, E.H.; Syah, Y.M. Prenylated dihydrostilbenes from *Macaranga rubiginosa*. *Chem. Nat. Compd.* **2017**, *53*, 215–218.
15. Tanjung, M.; Tjahjandarie, T.S.; Sentosa, M.H. Antioxidant and cytotoxic agent from the rhizomes of *Kaempferia pandurata*. *Asian Pac. J. Trop. Dis.* **2013**, *3*, 401–404.
16. Tanjung, M.; Hakim, E.H.; Elfahmi, Latip, J.; Syah, Y.M. Dihydroflavonol and flavonol derivatives from *Macaranga recurvata*. *Nat. Prod. Commun.* **2012**, *10*, 1309–1310.
17. Tanjung, M.; Saputri, R.D.; Wahjoedi, R.A.; Tjahjandarie, T.S. 4-Methoxy-3-(3-methylbut-2-en-1-yl)-7-(3-methylbut-2-en-1-yl)oxyquinolin-2(1H)-one from *Melicope moluccana* T.G. Hartley. *Molbank* **2017**, M939.
18. Tjahjandarie, T.S.; Pudjiastuti, P.; Saputri, R.D.; Tanjung, M. Antimalaria and antioxidant activity of phenolic compounds isolated from *Erythrina crista-galli* L. *J. Chem. Pharm. Res.* **2014**, *6*, 786–790.



© 2017 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en -1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12 H-pyrano[2,3-a]xanthen-12-one from Calophyllum pseudomole

ORIGINALITY REPORT

22%

SIMILARITY INDEX

15%

INTERNET SOURCES

20%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | mdpi.com
Internet Source | 2% |
| 2 | repository.um.edu.my
Internet Source | 2% |
| 3 | www.sphinxsai.com
Internet Source | 1% |
| 4 | researchbank.rmit.edu.au
Internet Source | 1% |
| 5 | Hui Xiong, Yanan Zheng, Guangzhong Yang, Huixia Wang, Zhinan Mei. "Triterpene saponins with anti-inflammatory activity from the stems of <i>Entada phaseoloides</i> ", <i>Fitoterapia</i> , 2015
Publication | 1% |
| 6 | Block, S.. "Diterpenes from the leaves of <i>Croton zambesicus</i> ", <i>Phytochemistry</i> , 200404
Publication | 1% |
| 7 | www.thieme-connect.de
Internet Source | |

1%

1%

8

Chan, Gomathi, Khalijah Awang, Nor Hadiani Ismail, Seik Weng Ng, and Edward R. T. Tiekink. "6-[(2E)-3,7-Dimethylocta-2,6-dien-1-yl]-5,7-dihydroxy-8-(2-methylbutanoyl)-4-phenyl-2H-chromen-2-one–6-[(2E)-3,7-dimethylocta-2,6-dien-1-yl]-5,7-dihydroxy-8-(3-methylbutanoyl)-4-phenyl-2H-chromen-2-one (1/1) from *Mesua elegans*", *Acta Crystallographica Section E Structure Reports Online*, 2012.

Publication

1%

9

Alain Meli Lannang, Denis Kehdinga Sema, Simplicie J. N. Tatsimo, V. F. Tsague Tankeu et al. " A new depsidone derivative from the leaves of ", *Natural Product Research*, 2017

Publication

1%

10

Tanjung, Mulyadi, Ratih Dewi Saputri, Faiz Fakhriah Fitriati, and Tjitjik Srie Tjahjandarie. "Antimalarial and Antioxidant Activities of Isoprenylated Coumarins from the Stem Bark of *Mesua borneensis* L.", *Journal of Biologically Active Products from Nature*, 2016.

Publication

1%

11

Jaeyoung Kwon, Nguyen Tuan Hiep, Dong-Woo Kim, Sungeun Hong et al. " Chemical

Constituents Isolated from the Root Bark of and
Their Potential Neuroprotective Effects ",
Journal of Natural Products, 2016

Publication

12

Mah, Siau Hui, Gwendoline Cheng Lian Ee,
Soek Sin Teh, Mawardi Rahmani, Yang Mooi
Lim, and Rusea Go. "Phylattrin, a New Cytotoxic
Xanthone from *Calophyllum soulattri*",
Molecules, 2012.

Publication

1%

13

dict.cnki.net

Internet Source

1%

14

www.dovepress.com

Internet Source

1%

15

www.freepatentsonline.com

Internet Source

1%

16

ijddr.in

Internet Source

1%

17

Yan, C.. "A new sesterterpenoid from the fruits
of *Phellodendron chinense* var. *glabriusculum*
Schneid", Chinese Chemical Letters, 200808

Publication

1%

18

Guo-Liang Li, Wei-Jie Guo, Guang-Bao Wang,
Rong-Rong Wang, Yu-Xue Hou, Kun Liu, Yang
Liu, Wei Wang. "Sterols from the Green Alga
Ulva australis", Marine Drugs, 2017

<1%

19

Nasir, Nadiah Mad, Mawardi Rahmani, Khozirah Shaari, Gwendoline Cheng Lian Ee, Rusea Go, Nur Kartinee Kassim, Siti Noor Kamilah Muhamad, and Mohd Johadi Iskandar. "Two New Xanthones from *Calophyllum nodusum* (Guttiferae)", *Molecules*, 2011.

Publication

<1%

20

weather.ninemsn.com.au

Internet Source

<1%

21

www.jstage.jst.go.jp

Internet Source

<1%

22

Wu, Bin, Wen Hui Lin, Hui Yuan Gao, Lu Zheng, Li Jun Wu, and Chul Sa Kim. "Four New Antibacterial Constituents from *Senecio cannabifolius*.", *Pharmaceutical Biology*, 2006.

Publication

<1%

23

hzi.openrepository.com

Internet Source

<1%

24

sistemas.ufms.br

Internet Source

<1%

25

Dai, Hao-Fu, Yan-Bo Zeng, Qi Xiao, Zhuang Han, You-Xing Zhao, and Wen-Li Mei. "Caloxanthones O and P: Two New Prenylated Xanthones from *Calophyllum inophyllum*", *Molecules*, 2010.

<1%

26 García-Zebadúa Julio César, Magos-Guerrero Gil Alfonso, Mumbrú-Massip Marius, Estrada-Muñoz Elizabeth et al. "Inhibition of HIV-1 reverse transcriptase, toxicological and chemical profile of *Calophyllum brasiliense* extracts from Chiapas, Mexico", *Fitoterapia*, 2011
Publication

27 ijcc.iregway.com
Internet Source

28 www.phytojournal.com
Internet Source

29 repositorio.unesp.br
Internet Source

30 lib.ugent.be
Internet Source

31 Alexander Pontius, Anja Krick, Stefan Kehraus, Reto Brun, Gabriele M. König. " Antiprotozoal Activities of Heterocyclic-Substituted Xanthones from the Marine-Derived Fungus sp. ", *Journal of Natural Products*, 2008
Publication

32 M. M. M. Pinto. "Synthesis of Prenylated Xanthones: An Overview", *Current Organic Chemistry*, 08/01/2009

33 pertanika.upm.edu.my <1%
Internet Source

34 www.jove.com <1%
Internet Source

35 Tag Wang. "New 2-(2-Phenylethyl) Chromones from Bothriochloa Ischaemum", Journal of Asian Natural Products Research, 3/1/2001 <1%
Publication

36 Adonis Bello Alarcón, Osmany Cuesta-Rubio, Jorge Cárdenas Pérez, Anna Lisa Piccinelli, Luca Rastrelli. " Constituents of the Cuban Endemic Species ", Journal of Natural Products, 2008 <1%
Publication

37 Chihiro Ito, Masataka Itoigawa, Yoshiaki Miyamoto, K. Sundar Rao et al. " A New Biflavonoid from with Antitumor-Promoting Activity ", Journal of Natural Products, 1999 <1%
Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

5,9,11-Trihydroxy-10-(2''-hydroxy-3''-methylbut-3''-en -1-yl)-2,2-dimethyl-3-(2'-methylbut-3'-en-2'-yl)-2H,12 H-pyrano[2,3-a]xanthen-12-one from Calophyllum pseudomole

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
