

also developed by scimago:

 SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal & Country Ran

Enter Journal Title, ISSN or Publisher Name



[Home](#)

[Journal Rankings](#)

[Country Rankings](#)

[Viz Tools](#)

[Help](#)

[About Us](#)

Natural Product Sciences

Country [South Korea](#) -  [SJR Ranking of South Korea](#)

22

Subject Area and Category [Chemistry](#)
[Organic Chemistry](#)

[Pharmacology, Toxicology and Pharmaceutics](#)
[Drug Discovery](#)

H Index

Publisher [Korean Society of Pharmacognosy](#)

Publication type Journals

ISSN 12263907

Coverage 1997-ongoing

Scope Natural Product Sciences is the official publication of the Korean Society of Pharmacognosy which was launched in 1995. Natural Product Sciences will primarily publish research papers on original work, either experimental or theoretical, that advance our understanding of natural product sciences, including important questions of phytochemistry, chemistry, and bio- chemistry of natural resources. Timely reviews and commentaries on recent progress in active areas of natural products will be also published.


 [Homepage](#)

[How to publish in this journal](#)

[Contact](#)

 [Join the conversation about this journal](#)


Quartiles



 SJR





I'm not a robot



reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the p
 specific journal. The purpose is to have a forum in which
 journal, experiences and other issues derived from the pu
 articles, maintain the dialogue through the usual channel:



Powered by:




Scopus

@ScimagoJR

2019. Data Source: Scopus®

IN REBUS

e 1,1,106)



Natural Product Sciences

Q3

Drug Discovery

best quartile

SJR 2018

0.27

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scim
```

Online Submission System

Click →

Aims and Scope

Editorial Board

Archives

Instructions to Authors

Contact Us

KC-ST
한국과학기술단체총연합회

NPS Journal
Homepage QR Code



Editorial Board

Editor in Chief

Jee H. Jung	<i>Pusan National Univ.</i>
--------------------	-----------------------------

Associate Editor

Dae-Keun Kim	<i>Woosuk Univ.</i>
Won Keun Oh	<i>Seoul National Univ.</i>

Editorial Board

Sun Yeou Kim <i>Gachon Univ.</i>	Youn Chul Kim <i>Wonkwang Univ.</i>	Jin Sook Kim <i>Korea Institute of Oriental Me</i>
Byung-Sun Min <i>Catholic Univ. of Daegu</i>	Nam-In Baek <i>Kyung-Hee Univ.</i>	Eun-Kyung Seo <i>Ewha Womans Univ.</i>
Hyun Ok Yang <i>KIST, Gangneung</i>	Eun-Rhan Woo <i>Chosun Univ.</i>	Shi Yong Ryu <i>KRICT</i>
Dongho Lee <i>Korea Univ.</i>	Min Won Lee <i>Chung-Ang Univ.</i>	Sang Kook Lee <i>Seoul National Univ.</i>
Seung Ho Lee <i>Yeungnam Univ.</i>	Ik-Soo Lee <i>Chonnam National Univ.</i>	Hyun sun Lee <i>KRIBB</i>
Bang Yeon Hwang <i>Chungbuk National Univ.</i>	Hee Jae Shin <i>KIOST</i>	

Advisory Board

Yeong Shik Kim <i>Seoul National Univ.</i>	Young Ho Kim <i>Chungnam National Univ.</i>	Jinwoong Kim <i>Seoul National Univ.</i>
Jeoung Hill Park <i>Seoul National Univ.</i>	Hee-Juhn Park <i>Sangji Univ.</i>	Kang Ro Lee <i>Sungkyunkwan Univ.</i>
Hyeong-Kyu Lee <i>KRIBB</i>	Jongheon Shin <i>Seoul National Univ.</i>	Jae Sue Choi <i>Pukyong National Univ.</i>

International Advisory Board

Zheng Cui <i>Shenyang Pharmaceutical Univ.</i>	Masayuki Mikage <i>Kanazawa Univ.</i>	Shigetoshi Kadota <i>Toyama Med. Pharmaceut. U</i>
Shao-Qing Cai <i>Peking Univ.</i>	William Gerwick <i>University of California at San Diego</i>	Kirk Gustafson <i>National Cancer Institute</i>
Shigeki Matsunaga <i>Tokyo Univ.</i>	Masami Ishibashi <i>Chiba Univ.</i>	Yuewei Guo <i>Shanghai Institute of Materia</i>
Guowei Qin <i>Shanghai Institute of Materia Medica</i>	Yukihiro Shoyama <i>Nagasaki International Univ.</i>	Chengbin Cui <i>Ocean Univ. of China</i>

Natural Product Sciences

Published by The Korean Society of Pharm

IS
IS

Online Submission System

Click →

Aims and Scope

Editorial Board

Archives

Instructions to Authors

Contact Us

KC-ST
한국과학기술단체총연합회

NPS Journal
Homepage QR Code



Archives

Quick search

Full search

Search in each volume

• Publication Natural Product Sciences

• Years 2018

• Vol. No. Vol. 24 No. 4

※ Please click a search button, after select Volume and number.

• Search in this vol.

Search

11. Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of *Acronychia pedunculata* (PDF file / 4 p View | Do

Mulyadi Tanjung , Intan Nurmalasari , Aisyah Kanti Wilujeng , Ratih Dewi Saputri , Fida Rachmadiarti , Tjitjik Srie Tjahjandarie

12. Efficient Isolation of Dihydrophaseic acid 3'-O-β-D-Glucopyranoside from *Nelumbo nucifera* Seeds Using High-performance Countercurrent Chromatography and Reverse-phased High-performance Liquid Chromatography (PDF file / 5 p View | Do

Taewoong Rho , Kee Dong Yoon

13. Effect of *Sasa quelpaertensis* Nakai Extracts and its Constituent p-coumaric Acid on the Apoptosis of Human Cancer Cell Lines (PDF file / 5 p View | Do

Mi Gyeong Jang , Hee Chul Ko , Se Jae Kim

< 1 2 >

2013 ©Copyright The Korean Society of Pharmacognosy. All Rights Reserved.

Seoul National University, 1, Gwanak-ro, Gwanak-gu, Seoul 08826, Korea

Tel: +82-2-880-7872 | E-mail: journal@e-nps.or.kr | Powered by [INFORang Co., Ltd](http://www.inforang.com)



Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of *Acronychia pedunculata*

Mulyadi Tanjung^{1,*}, Intan Nurmalasari¹, Aisyah Kanti Wilujeng¹, Ratih Dewi Saputri¹, Fida Rachmadiarti², and Tjitjik Srie Tjahjandarie¹

¹Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya, Surabaya, Indonesia

Abstract – A new isoprenylated acetophenone, acronyculatin P (**1**) as well as two known compounds, 3',5'-diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (**2**) and 3'-isoprenyl-2',4',6'-trihydroxyphenylethanone (**3**) were isolated from the stem bark of *Acronychia pedunculata* (L.) Miq. The structures were determined by HRESIMS, 1D and 2D NMR. The inhibitory activity of the isoprenylated acetophenone derivatives against murine leukemia P-388 cells showed compound **1** moderate activity with IC₅₀ 15.42 μM.

Keywords – Acronyculatin P, isoprenylated acetophenone, *Acronychia pedunculata*, P-388 cells

Introduction

Acronychia pedunculata is one species belongs to the Rutaceae family found in all of Indonesia. The stem bark have been used in traditional medicine for the treatment of fever, asthma, diarrhea, and rheumatism.¹ According to previous studies, the most common secondary metabolites isolated from *A. pedunculata* are alkaloids,² coumarins,³ and isoprenylated acetophenone derivatives.^{4,5} Isoprenylated acetophenone derivatives in the genus *Acronychia* indicate not their value as chemotaxonomic markers of the genus. Isoprenylated acetophenone derivatives were reported to possess cytotoxic,⁴ anti-inflammatory,⁶ and antioxidant⁷ activities. In the present study, a phytochemical investigation is reported of the stem bark of *A. pedunculata* focused on the isolation and structural elucidation of a new isoprenylated acetophenone derivatives, acronyculatin P (**1**) along with two known compounds, 3',5'-diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (**2**) and 3'-isoprenyl-2',4',6'-trihydroxyphenylethanone (**3**). The cytotoxic activity of compounds **1** - **3** against murine leukemia P-388 cells from this plant are also reported.

Experimental

General experimental procedures – Column chromatography and radial chromatography were carried out using silica gel 60 and silica gel 60 PF₂₅₄ (Merck, Darmstadt, Germany). UV spectra were recorded in MeOH on a Shimadzu series 1800 UV-VIS spectrophotometer (Kyoto, Japan). IR spectra were recorded in KBr on a One Perkin Elmer instrument (Waltham, MA, USA). NMR spectra were measured on a JEOL JNM-ECA 400 MHz FTNMR spectrophotometer (Tokyo, Japan) in CDCl₃ with TMS as the internal standard. Mass spectra were measured on an ESI-TOF Waters LCT Premier XE producing pseudo-molecular ions, [M-H]⁻ negative ion mode (Santa Clara, CA, USA).

Plant materials – The dried and powdered of stem bark of *A. pedunculata* was collected in July 2017 from Gunung Salak, Bogor, West Java, Indonesia by Mr. Ismail Rachman. The plant material was identified at the Herbarium Bogoriense, Bogor. A voucher specimen (AP 60329) was deposited in Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia.

Extraction and isolation – The stem bark of *A. pedunculata* (1.5 kg) was extracted with methanol at room temperature two times and then extracts were concentrated in vacuo. The methanol extract (350 g) was

*Author for correspondence

Mulyadi Tanjung, Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia
Tel: +62-31-5936501; E-mail: mulyadi-t@fst.unair.ac.id




Fig. 1. Isoprenylated acetophenones **1** - **3** isolated from *A. pedunculata*.

suspended in H₂O and partitioned with *n*-hexane (11.6 g) and ethyl acetate (7.7 g). The *n*-hexane extract (11 g) was further fractionated by column chromatography on silica gel (200 g) eluted with *n*-hexane-ethyl acetate by increasing polarity (from 9:1, 4:1; 7:3, and 1:1) to give three major fractions A-C. Fraction A (0.5 g) was separated by planar radial chromatography eluted with *n*-hexane-CHCl₃ (from 9:1 to 4:1) to produce subfractions A₁-A₂. Subfraction A₂ was purified by planar radial chromatography using *n*-hexane-diisopropylether (from 9:1 to 4:1) to yield compound **1** (46 mg). Fraction B (5.3 g) was refractionated using column chromatography and eluted with *n*-hexane-ethyl acetate (from 9:1 to 1:1) to produce three subfractions B₁-B₃. Subfraction B₂ was purified by planar radial chromatography using *n*-hexane-diisopropylether (from 9:1 to 4:1) to afford **2** (28 mg). The ethyl acetate extract (7.5 g) was fractionated over silica gel column chromatography eluted with *n*-hexane-ethyl acetate (from 9:1, 4:1; and 1:1) to give five major fractions D-H. Fraction E (0.75 g) was purified with *n*-hexane-ethyl acetate (from 4:1, 7:3; and 1:1) to afford **3** (24 mg).

Acronyculatin P (1) – Yellowish syrup. UV (MeOH) λ_{\max} nm (log ϵ) : 252 (4.25), and 283 (4.23). IR (KBr) ν_{\max} cm⁻¹: 3300, 1619, 1587 and 1184. ¹H and ¹³C NMR see Table 1. HRESIMS: *m/z* [M-H]⁻ calcd. for C₂₀H₂₈O₄ 331.1924, found 331.1909.

3',5'-Diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (2) – Yellowish syrup. UV (MeOH) λ_{\max} nm (log ϵ) : 224 (4.38), and 290 (4.21). HRESIMS: *m/z* [M-H]⁻ calcd. for C₁₉H₂₅O₄ 317.1536, found 317.1532.

3'-Isoprenyl-2',4',6'-trihydroxyphenylethanone (3) – Pale yellowish solid, mp. 173 - 175 °C. UV (MeOH) λ_{\max} nm (log ϵ) : 236 (4.32), and 280 (4.20). HRESIMS: *m/z* [M-H]⁻ calcd. for C₁₃H₁₅O₄ 235.2569, found 235.2560.

Cytotoxic activity – The human tumor cell used in this work was P-388 cells (murine leukemia) and cultured in RPMI 1640 medium. The P-388 cells were seeded into each 96-well cell culture plate at a density of 3 × 10⁴ cells/well and incubated at 37 °C for 48 h against murine

leukemia P-388 cells using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) reduction assay. Compounds **1-3** dissolved in DMSO by variations in concentration of 100; 30; 10; 3; 1; 0.3 and 0.1 µg/mL with triplicate treatment. The number of cells inhibited by each of compounds **1-3** were measured using microplate reader spectrometer at λ 540 nm. Artonin E was used as positive control and DMSO 1% was used as negative control.⁸⁻¹¹ The IC₅₀ values of the compounds were calculated through extrapolation 50% absorption lines to various concentrations using regression analysis.


Result and Discussion

The structures of **1-3** were elucidated by UV, IR, HRESIMS, 1D and 2D NMR spectroscopy. To our knowledge, compound (**1**) is a new compound. The ¹H and ¹³C NMR spectral data of compounds **2-3** are consistent with published data.¹²⁻¹³

Compound (**1**) was isolated as yellowish syrup. The HRESIMS of **1** exhibited a negative molecular ion peak [M-H]⁻ at *m/z* 331.1924 indicating a molecular formula of C₂₀H₂₇O₄ implying seven degrees of unsaturation. The UV maximum absorption at λ_{\max} 252 (4.25), 283 (4.23) nm and IR bands (ν_{\max} , cm⁻¹) at 3300, 1619, 1587 and 1184⁴. The ¹H NMR spectrum (Table 1) showed an acetyl signal at δ_{H} 2.71 (3H, s, H-2), a chelated hydroxyl group at δ_{H} 13.43 (1H, s, 2'-OH), and two methoxyl groups at δ_{H} 3.71 (6H, s, 4'/6'-OCH₃). In addition, compound **1** showed 3-methyl-1-butenyl proton signals at δ_{H} 6.40 (1H, d, *J* = 15.6 Hz, H-1''), 6.55 (1H, dd, *J* = 15.6; 7.1 Hz, H-2''), 2.48 (1H, dq, *J* = 15.6; 8.2 Hz, H-3''), 1.68 (3H, s, H-4''), 1.10 (6H, d, *J* = 6.8 Hz, H-4''/5''), and a 3-methyl-2-butenyl (isoprenyl) proton signals at δ_{H} 5.16 (1H, t, *J* = 6.6 Hz, H-2'''), 3.29 (2H, d, *J* = 6.6 Hz, H-1'''), 1.78 (3H, s, H-4'''), and 1.69 (3H, s, H-5'''). The placement of hydroxyl, methoxyl, 3-methyl-1-butenyl, and 3-methyl-2-butenyl groups of **1** were established by HMQC and HMBC spectra (Fig. 2). The ¹³C NMR spectrum of **1**

Table 1. NMR Spectroscopic data (400 MHz in CDCl₃) for acronyculatin P (1)

No.C	δ_H (mult, <i>J</i> in Hz)	δ_C	HMBC
1	-	204.7	-
2	2.71 (s, 3H)	31.4	C-1
1'	-	111.9	-
2'	-	161.5	-
3'	-	116.4	-
4'	-	163.5	-
5'	-	120.4	-
6'	-	159.6	-
1''	6.40 (<i>d</i> , 15.6, 1H)	116.7	C-4', C-2'', C-3''
2''	6.55 (<i>dd</i> , 7.1; 15.6, 1H)	143.1	C-1'', C-3'', C-4''/5''
3''	2.48 (<i>dq</i> , 8.2; 15.6, 1H)	32.9	C-1'', C-2'', C-4''/ C-5''
4''	1.10 (<i>d</i> , 6.8, 3H)	22.6	C-2'', C-3'', C-5''
5''	1.10 (<i>d</i> , 6.8, 3H)	22.6	C-2'', C-3'', C-4''
1'''	3.29 (<i>d</i> , 6.6, 2H)	23.2	C-4', C-5', C-6', C-2''', C-3'''
2'''	5.16 (<i>t</i> , 6.6, 1H)	123.8	C-4''', C-5'''
3'''	-	131.6	-
4'''	1.78 (<i>s</i> , 3H)	17.9	C-2''', C-3''', C-5'''
5'''	1.69 (<i>s</i> , 3H)	25.8	C-2''', C-3''', C-4'''
2'-OH	13.43 (<i>s</i> , 1H)	-	C-1', C-2', C-3'
4'-OCH ₃	3.71 (<i>s</i> , 3H)	60.4	C-4'
6'-OCH ₃	3.71 (<i>s</i> , 3H)	63.0	C-6'

**Fig. 2.** Selected HMBC correlations for compound 1.

revealed the signals for an acetyl group at δ_C 31.4 and δ_C 204.7 as well as three oxyaryl carbons at δ_C 163.5, δ_C 161.5, and δ_C 159.6 characteristic for a 2',4',6'-trioxygenated acetophenone derivatives.⁵ The HMBC spectrum revealed a cross-peak correlation between δ_H 2.71 (H-2) and an carbonyl carbon at δ_C 204.7. The proton signal of a chelated hydroxyl group (δ_H 13.43, 2'-OH) correlated with three quaternary carbons [δ_C 111.9 (C-1'); 161.5 (C-2'); 116.9 (C-3')]. The proton signal of methoxyl group at δ_H 3.71 (4'/6'-OCH₃) correlated with two oxyaryl carbons [δ_C 163.5, and δ_C 159.6] showing the presence of two

methoxyl groups at C-4' and C-6'. Furthermore, the proton signal of methylene of 3-methyl-2-butenyl chain at δ_H 3.29 has correlation with two oxyaryl carbons [δ_C 163.5 (C-4'), and δ_C 159.6 (C-6')], two quaternary carbons [δ_C 120.4 (C-5'), and δ_C 131.6 (C-3''')], and a methine carbon at δ_C 123.8 (C-2''') confirmed that the 3-methyl-2-butenyl chain is located at C-5'. The presence of long-range correlations between the proton signal of a vinylic of 3-methyl-1-butenyl chain at δ_H 6.40 (H-1'') was correlated to a oxyaryl carbon at δ_C 163.5 (C-4'), and two methine carbons at δ_C 143.1 (C-2''), and 32.9 (C-3'') reinforces the location of 3-methyl-1-butenyl chain at C-3'. Therefore, compound 1 was identified as (*E*)-1-(2'-hydroxy-4',6'-dimethoxy-3'-(3''-methylbut-1''-enyl)-5'-(3'''-methylbut-2'''-enyl)phenylethanone and given the trivial name acronyculatin P. Other HMBC correlations consistent with the structure 1 are shown in Table 1 and Fig. 2.

Compounds 1 - 3 were assessed for their cytotoxicity and results are shown in Table 2. The IC₅₀ values were 15.42, 27.26, and 80.59 μ M, respectively (artoinin E as a positive control, IC₅₀ 3.05 μ M). Compound 1 was more active than others, But, all the compounds were less active when compared with the positive control.

Table 2. Cytotoxic activity of compounds **1 - 3** against murine leukemia P-388 cells

Compounds	IC ₅₀ (μM)
Acronyculatin P (1)	15.42 ± 0.51
3',5'-Diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (2)	27.26 ± 1.23
3'-Isoprenyl-2',4',6'-trihydroxyphenylethanone (3)	80.59 ± 1.67

Acknowledgments

This research was supported by Universitas Airlangga through Hibah Riset Mandat 2018 research.

References

- (1) Hartley, T. G. *J. Arnold Arbor.* **1974**, *55*, 469-523.
- (2) de Silva, L. B.; de Silva, U. L. L.; Mahendran, M.; Jenings, R. M. *Phytochem.* **1979**, *18*, 1255-1256.
- (3) De Silva, L. B.; Herath, W. M.; Liyanage, C.; Kumar, V.; Ahmad, V. U.; Sultana, A. *Phytochem.* **1991**, *30*, 1709-1710.
- (4) Kozaki, S.; Takenaka, Y.; Mizushima, Y.; Yamaura, T.; Tanahashi, T. *J. Nat. Med.* **2014**, *68*, 421-426.
- (5) Kouloura, E.; Halabalaki, M.; Lallemand, M. C.; Nam, S.; Jove, R.; Litaudon, M.; Awang, K.; Hadi, H. A.; Skaltsounis, A. L. *J. Nat. Prod.* **2012**, *75*, 1270-1276.
- (6) Pathmasiri, W.; el-Seedi, H. R.; Han, X.; Janson, J. C.; Huss, U.; Bohlin, L. *Chem. Biodivers.* **2005**, *2*, 463-469.
- (7) Su, C. R.; Kuo, P. C.; Wang, M. L.; Liou, M. J.; Damu, A. G.; Wu, T. S. *J. Nat. Prod.* **2003**, *66*, 990-993.
- (8) Tanjung, M.; Rachmadiarti, F.; Saputri, R. D.; Tjahjandarie, T. S. *Nat. Prod. Res.* **2018**, *32*, 1062-1067.
- (9) Tanjung, M.; Hakim, E. H.; Syah, Y. M. *Chem Nat Compd.* **2017**, *53*, 215-218.
- (10) Marliana, E.; Astuti, W.; Kosala, K.; Hairani, R.; Tjahjandarie, T. S.; Tanjung, M. *Asian J. Chem.* **2018**, *30*, 795-798.
- (11) Tanjung, M.; Saputri, R. D.; Tjahjandarie, T. S. *Molbank.* **2016**, M906, 3, 1-5.
- (12) Han, X.; Pathmasiri, W.; Bohlin, L.; Janson, J. C. *J. Chromatogr. A.* **2004**, *1022*, 213-216.
- (13) Basabe, P.; de Román, M.; Marcos, I. S.; Diez, D.; Blanco, A.; Boderó, O.; Mollinedo, F.; Sierra, B. G.; Urones, J. G. *Eur. J. Med. Chem.* **2010**, *45*, 4258-4269.

Received May 31, 2018

Revised August 17, 2018

Accepted August 18, 2018