

Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of *Acronychia pedunculata*

by Mulyadi Tanjung

Submission date: 05-Feb-2019 12:04PM (UTC+0800)

Submission ID: 1073198178

File name: ed_Acetophenone_from_the_Stem_Bark_of_Acronychia_pedunculata.pdf (323.85K)

Word count: 2176

Character count: 10739



6

Natural Product Sciences

24(4) : 1-4 (2018)

<https://doi.org/10.20307/nps.2018.24.4.1>

Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of *Acronychia pedunculata*

Mulyadi Tanjung^{1,*}, Intan Nurmalasari¹, Aisyah Kanti Wilujeng¹, Ratih Dewi Saputri¹,
Fida Rachmadiarti², and Tjitjik Srie Tjahjandarie¹

¹Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Surabaya, Surabaya, Indonesia

Abstract – A new isoprenylated acetophenone, acronyculatin P (**1**) as well as two known compounds, 3',5'-diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (**2**) and 3'-isoprenyl-2',4',6'-trihydroxyphenylethanone (**3**) were isolated from the stem bark of *Acronychia pedunculata* (L.) Miq. The structures were determined by HRESIMS, 1D and 2D NMR. The inhibitory activity of the isoprenylated acetophenone derivatives against murine leukemia P-388 cells showed compound **1** moderate activity with IC₅₀ 15.42 μM.

Keywords – Acronyculatin P, isoprenylated acetophenone, *Acronychia pedunculata*, P-388 cells

Introduction

Acronychia pedunculata is one species belongs to the Rutaceae family found in all of Indonesia. The stem bark have been used in traditional medicine for the treatment of fever, asthma, diarrhea, and rheumatism.¹ According to previous studies, the most common secondary metabolites isolated from *A. pedunculata* are alkaloids,² coumarins,³ and isoprenylated acetophenone derivatives.^{4,5} Isoprenylated acetophenone derivatives in the genus *Acronychia* indicate not their value as chemotaxonomic markers of the genus. Isoprenylated acetophenone derivatives were reported to possess cytotoxic,⁴ anti-inflammatory,⁶ and antioxidant⁷ activities. In the present study, a phytochemical investigation is reported of the stem bark of *A. pedunculata* focused on the isolation and structural elucidation of a new isoprenylated acetophenone derivatives, acronyculatin P (**1**) along with two known compounds, 3',5'-diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (**2**) and 3'-isoprenyl-2',4',6'-trihydroxyphenylethanone (**3**). The cytotoxic activity of compounds **1** - **3** against murine leukemia P-388 cells from this plant are also reported.

Experimental

General experimental procedures – Column chromatography and radial chromatography were carried out using silica gel 60 and silica gel 60 PF₂₅₄ (Merck, Darmstadt, Germany). UV spectra were recorded in MeOH on a Shimadzu series 1800 UV-VIS spectrophotometer (Kyoto, Japan). IR spectra were recorded in KBr on a One Perkin Elmer instrument (Waltham, MA, USA). NMR spectra were measured on a JEOL JNM-ECA 400 MHz FTNMR spectrophotometer (Tokyo, Japan) in CDCl₃ with TMS as the internal standard. Mass spectra were measured on an ESI-TOF Waters LCT Premier XE producing pseudo-molecular ions, [M-H]⁻ negative ion mode (Santa Clara, CA, USA).

Plant materials – The dried and powdered of stem bark of *A. pedunculata* was collected in July 2017 from Gunung Salak, Bogor, West Java, Indonesia by Mr. Ismail Rachman. The plant material was identified at the Herbarium Bogoriense, Bogor. A voucher specimen (AP 60329) was deposited in Herbarium Bogoriense, Center of Biological Research and Development, National Institute of Science, Bogor, Indonesia.

Extraction and isolation – The stem bark of *A. pedunculata* (1.5 kg) was extracted with methanol at room temperature two times and then extracts were concentrated in vacuo. The methanol extract (350 g) was

*Author for correspondence

Mulyadi Tanjung, Natural Products Chemistry Research Group, Organic Chemistry Division, Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, Indonesia
Tel: +62-31-5936501; E-mail: mulyadi-t@fst.unair.ac.id




Fig. 1. Isoprenylated acetophenones 1 - 3 isolated from *A. pedunculata*.

suspended in H₂O and partitioned with *n*-hexane (11.6 g) and ethyl acetate (7.7 g). The *n*-hexane extract (11 g) was further fractionated by column chromatography on silica gel (200 g) eluted with *n*-hexane-ethyl acetate by increasing polarity (from 9:1, 4:1; 7:3, and 1:1) to give three major fractions A-C. Fraction A (0.5 g) was separated by planar radial chromatography eluted with *n*-hexane-CHCl₃ (from 9:1 to 4:1) to produce subfractions A₁-A₂. Subfraction A₂ was purified by planar radial chromatography using *n*-hexane-diisopropylether (from 9:1 to 4:1) to yield compound **1** (46 mg). Fraction B (5.3 g) was refractionated using column chromatography and eluted with *n*-hexane-ethyl acetate (from 9:1 to 1:1) to produce three subfractions B₁-B₃. Subfraction B₂ was purified by planar radial chromatography using *n*-hexane-diisopropylether (from 9:1 to 4:1) to afford **2** (28 mg). The ethyl acetate extract (7.5 g) was fractionated over silica gel column chromatography eluted with *n*-hexane-ethyl acetate (from 9:1, 4:1; and 1:1) to give five major fractions D-H. Fraction E (0.75 g) was purified with *n*-hexane-ethyl acetate (from 4:1, 7:3; and 1:1) to afford **3** (24 mg).

Acronyculatin P (1) – Yellowish syrup. UV (MeOH) λ_{\max} nm (log ϵ): 252 (4.25), and 283 (4.23). IR (KBr) ν_{\max} cm⁻¹: 3300, 1619, 1587 and 1184. ¹H and ¹³C NMR see Table I. HRESIMS: m/z [M-H]⁻ calcd. for C₂₀H₂₈O₄ 331.1924, found 331.1909.

3',5'-Diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (2) – Yellowish syrup. UV (MeOH) λ_{\max} nm (log ϵ): 224 (4.38), and 290 (4.21). HRESIMS: m/z [M-H]⁻ calcd. for C₁₉H₂₅O₄ 317.1536, found 317.1532.

3'-Isoprenyl-2',4',6'-trihydroxyphenylethanone (3) – Pale yellowish solid, mp. 173 - 175 °C. UV (MeOH) λ_{\max} nm (log ϵ): 236 (4.32), and 280 (4.20). HRESIMS: m/z [M-H]⁻ calcd. for C₁₃H₁₅O₄ 235.2569, found 235.2560.

Cytotoxic activity – The human tumor cell used in this work was P-388 cells (murine leukemia) and cultured in RPMI 1640 medium. The P-388 cells were seeded into each 96-well cell culture plate at a density of 3 × 10⁴ cells/well and incubated at 37 °C for 48 h against murine

leukemia P-388 cells using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-2H-tetrazolium bromide (MTT) reduction assay. Compounds **1-3** dissolved in DMSO by variations in concentration of 100; 30; 10; 3; 1; 0.3 and 0.1 µg/mL with triplicate treatment. The number of cells inhibited by each of compounds **1-3** were measured using microplate reader spectrometer at λ 540 nm. Artonin E was used as positive control and DMSO 1% was used as negative control.⁸⁻¹¹ The IC₅₀ values of the compounds were calculated through extrapolation 50% absorption lines to various concentrations using regression analysis.


Result and Discussion

The structures of **1-3** were elucidated by UV, IR, HRESIMS, 1D and 2D NMR spectroscopy. To our knowledge, compound (**1**) is a new compound. The ¹H and ¹³C NMR spectral data of compounds **2-3** are consistent with published data.¹²⁻¹³

Compound (**1**) was isolated as yellowish syrup. The HRESIMS of **1** exhibited a negative molecular ion peak [M-H]⁻ at m/z 331.1924 indicating a molecular formula of C₂₀H₂₇O₄ implying seven degrees of unsaturation. The UV maximum absorption at λ_{\max} 252 (4.25), 283 (4.23) nm and IR bands (ν_{\max} , cm⁻¹) at 3300, 1619, 1587 and 1184⁴. The ¹H NMR spectrum (Table 1) showed an acetyl signal at δ_H 2.71 (3H, s, H-2), a chelated hydroxyl group at δ_H 13.43 (1H, s, 2'-OH), and two methoxyl groups at δ_H 3.71 (6H, s, 4'/6'-OCH₃). In addition, compound **1** showed 3-methyl-1-butenyl proton signals at δ_H 6.40 (1H, d, J = 15.6 Hz, H-1''), 6.55 (1H, dd, J = 15.6; 7.1 Hz, H-2''), 2.48 (1H, dq, J = 15.6; 8.2 Hz, H-3''), 1.68 (3H, s, H-4''), 1.10 (6H, d, J = 6.8 Hz, H-4''/5''), and a 3-methyl-2-butenyl (isoprenyl) proton signals at δ_H 5.16 (1H, t, J = 6.6 Hz, H-2'''), 3.29 (2H, d, J = 6.6 Hz, H-1'''), 1.78 (3H, s, H-4'''), and 1.69 (3H, s, H-5'''). The placement of hydroxyl, methoxyl, 3-methyl-1-butenyl, and 3-methyl-2-butenyl groups of **1** were established by HMQC and HMBC spectra (Fig. 2). The ¹³C NMR spectrum of **1**

Table 1. NMR Spectroscopic data (400 MHz in CDCl₃) for acronyculatin P (**1**)

No.C	δ_H (mult, <i>J</i> in Hz)	δ_C	HMBC
1	-	204.7	-
2	2.71 (s, 3H)	31.4	C-1
1'	-	111.9	-
2'	-	161.5	-
3'	-	116.4	-
4'	-	163.5	-
5'	-	120.4	-
6'	-	159.6	-
1''	6.40 (<i>d</i> , 15.6, 1H)	116.7	C-4', C-2'', C-3''
2''	6.55 (<i>dd</i> , 7.1; 15.6, 1H)	143.1	C-1'', C-3'', C-4''/5''
3''	2.48 (<i>dq</i> , 8.2; 15.6, 1H)	32.9	C-1'', C-2'', C-4''/ C-5''
4''	1.10 (<i>d</i> , 6.8, 3H)	22.6	C-2'', C-3'', C-5''
5''	1.10 (<i>d</i> , 6.8, 3H)	22.6	C-2'', C-3'', C-4''
1'''	3.29 (<i>d</i> , 6.6, 2H)	23.2	C-4', C-5', C-6', C-2''', C-3'''
2'''	5.16 (<i>t</i> , 6.6, 1H)	123.8	C-4''', C-5'''
3'''	-	131.6	-
4'''	1.78 (s, 3H)	17.9	C-2''', C-3''', C-5'''
5'''	1.69 (s, 3H)	25.8	C-2''', C-3''', C-4'''
2'-OH	13.43 (s, 1H)	-	C-1', C-2', C-3'
4'-OCH ₃	3.71 (s, 3H)	60.4	C-4'
6'-OCH ₃	3.71 (s, 3H)	63.0	C-6'

**Fig. 2.** Selected HMBC correlations for compound **1**.

revealed the signals for an acetyl group at δ_C 31.4 and δ_C 204.7 as well as three oxyaryl carbons at δ_C 163.5, δ_C 161.5, and δ_C 159.6 characteristic for a 2',4',6'- trioxy-generated acetophenone derivatives.⁵ The HMBC spectrum revealed a cross-peak correlation between δ_H 2.71 (H-2) and an carbonyl carbon at δ_C 204.7. The proton signal of a chelated hydroxyl group (δ_H 13.43, 2'-OH) correlated with three quaternary carbons [δ_C 111.9 (C-1'); 161.5 (C-2'); 116.9 (C-3')]. The proton signal of methoxy group at δ_H 3.71 (4'/6'-OCH₃) correlated with two oxyaryl carbons [δ_C 163.5, and δ_C 159.6] showing the presence of two

methoxy groups at C-4' and C-6'. Furthermore, the proton signal of methylene of 3-methyl-2-butenyl chain at δ_H 3.29 has correlation with two oxyaryl carbons [δ_C 163.5 (C-4'), and δ_C 159.6 (C-6')], two quaternary carbons [δ_C 120.4 (C-5'), and δ_C 131.6 (C-3'')], and a methine carbon at δ_C 123.8 (C-2''') confirmed that the 3-methyl-2-butenyl chain is located at C-5'. The presence of long-range correlations between the proton signal of a vinylic of 3-methyl-1-butenyl chain at δ_H 6.40 (H-1'') was correlated to a oxyaryl carbon at δ_C 163.5 (C-4'), and two methine carbons at δ_C 143.1 (C-2''), and 32.9 (C-3'') reinforces the location of 3-methyl-1-butenyl chain at C-3'. Therefore, compound **1** was identified as (*E*)-1-(2'-hydroxy-4',6'-dimethoxy-3'-(3''-methylbut-1''-enyl)-5'-(3'''-methylbut-2'''-enyl)phenylethanone and given the trivial name acronyculatin P. Other HMBC correlations consistent with the structure **1** are shown in Table 1 and Fig. 2.

Compounds **1** - **3** were assessed for their cytotoxicity and results are shown in Table 2. The IC₅₀ values were 15.42, 27.26, and 80.59 μ M, respectively (artoinin E as a positive control, IC₅₀ 3.05 μ M). Compound **1** was more active than others, But, all the compounds were less active when compared with the positive control.

Table 2. Cytotoxic activity of compounds **1-3** against murine leukemia P-388 cells

Compounds	IC ₅₀ (μM)
Acronyculatin P (1)	15.42 ± 0.51
3',5'-Diisoprenyl-2',4'-dihydroxy-6'-methoxyphenylethanone (2)	27.26 ± 1.23
3'-Isoprenyl-2',4',6'-trihydroxyphenylethanone (3)	80.59 ± 1.67

Acknowledgments

This research was supported by Universitas Airlangga through Hibah Riset Mandat 2018 research.

References

- (1) Hartley, T. G. *J. Arnold Arbor*. **1974**, *55*, 469-523.
- (2) de Silva, L. B.; de Silva, U. L. L.; Mahendran, M.; Jenings, R. M. *Phytochem*. **1979**, *18*, 1255-1256.
- (3) De Silva, L. B.; Herath, W. M.; Liyanage, C.; Kumar, V.; Ahmad, V. U.; Sultana, A. *Phytochem*. **1991**, *30*, 1709-1710.
- (4) Kozaki, S.; Takenaka, Y.; Mizushima, Y.; Yamaura, T.; Tanahashi, T. *J. Nat. Med*. **2014**, *68*, 421-426.
- (5) Kouloura, E.; Halabalaki, M.; Lallemand, M. C.; Nam, S.; Jove, R.; Litaudon, M.; Awang, K.; Hadi, H. A.; Skaltsounis, A. L. *J. Nat. Prod*. **2012**, *75*, 1270-1276.
- (6) Pathmasiri, W.; el-Seedi, H. R.; Han, X.; Janson, J. C.; Huss, U.; Bohlin, L. *Chem. Biodivers*. **2005**, *2*, 463-469.
- (7) Su, C. R.; Kuo, P. C.; Wang, M. L.; Liou, M. J.; Damu, A. G.; Wu, T. S. *J. Nat. Prod*. **2003**, *66*, 990-993.
- (8) Tanjung, M.; Rachmadiarti, F.; Saputri, R. D.; Tjahjandarie, T. S. *Nat. Prod. Res*. **2018**, *32*, 1062-1067.
- (9) Tanjung, M.; Hakim, E. H.; Syah, Y. M. *Chem Nat Compd*. **2017**, *53*, 215-218.
- (10) Marlina, E.; Astuti, W.; Kosala, K.; Hairani, R.; Tjahjandarie, T. S.; Tanjung, M. *Asian J. Chem*. **2018**, *30*, 795-798.
- (11) Tanjung, M.; Saputri, R. D.; Tjahjandarie, T. S. *Molbank*. **2016**, M906, 3, 1-5.
- (12) Han, X.; Pathmasiri, W.; Bohlin, L.; Janson, J. C. *J. Chromatogr. A*. **2004**, *1022*, 213-216.
- (13) Basabe, P.; de Román, M.; Marcos, I. S.; Diez, D.; Blanco, A.; Boderó, O.; Mollinedo, F.; Sierra, B. G.; Urones, J. G. *Eur. J. Med. Chem*. **2010**, *45*, 4258-4269.

Received November 0, 2018

Revised December 0, 2018

Accepted December 0, 2018

Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of Acronychia pedunculata

ORIGINALITY REPORT

23%

SIMILARITY INDEX

7%

INTERNET SOURCES

20%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Mulyadi Tanjung, Fida Rachmadiarti, Anindita Prameswari, Villa Ultha Wustha Agyani et al. "Airlanggins A-B, two new isoprenylated benzofuran-3-ones from the stem bark of ", *Natural Product Research*, 2017
Publication 15%
- 2 www.amse-aixmarseille.fr
Internet Source 2%
- 3 pubs.acs.org
Internet Source 1%
- 4 www.db-thueringen.de
Internet Source 1%
- 5 Xue, Jinghua, Ping Wu, Liangxiong Xu, and Xiaoyi Wei. "Penicillitone, a Potent in Vitro Anti-inflammatory and Cytotoxic Rearranged Sterol with an Unusual Tetracycle Core Produced by *Penicillium purpurogenum*", *Organic Letters*, 2014.
Publication 1%

6	orbi.uliege.be Internet Source	1%
7	tacoutdoors.net Internet Source	<1%
8	Linnert, M.. "Synthesis, structure and characterization of [Mg ⁷ (@m ³ -OCH ² CH ² OMe) ⁶ (@m-OCH ² CH ² OMe) ⁶] [Al(n-Bu) ⁴] ² - A cationic heptanuclear magnesium complex consisting of discrete cations and tetrabutylaluminate anions", Polyhedron, 20080120 Publication	<1%
9	www.kochi-tech.ac.jp Internet Source	<1%
10	www.tdx.cat Internet Source	<1%
11	Mulyadi Tanjung, Tjitjik Srie Tjahjandarie, Mulya Hadi Sentosa. "Antioxidant and cytotoxic agent from the rhizomes of Kaempferia pandurata", Asian Pacific Journal of Tropical Disease, 2013 Publication	<1%
12	Xiaoyu LIU, Wen LIU, Ling YANG, Beili XIA, Jinyan LI, Ji ZUO, Xiaotian LI. "Increased Connexin 43 Expression Improves the Migratory and Proliferative Ability of H9c2 Cells by Wnt-3a Overexpression", Acta Biochimica et Biophysica	<1%

Sinica, 2007

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Acronyculatin P, A New Isoprenylated Acetophenone from the Stem Bark of *Acronychia pedunculata*

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
