

# PHENOLIC COMPOUNDS FROM *Aquilaria microcarpa* STEM BARK

*by* Mulyadi Tanjung

---

**Submission date:** 08-May-2018 06:06PM (UTC+0800)

**Submission ID:** 960728578

**File name:** 16\_17\_33\_Alfinda\_1111\_1115.pdf (361.01K)

**Word count:** 3019

**Character count:** 14158

## PHENOLIC COMPOUNDS FROM *Aquilaria microcarpa* STEM BARK

Alfinda Novi Kristanti, Mulyadi Tanjung, Okky P. Rahayu, Erika Herdiana

13

Department of Chemistry, Faculty of Science and Technology  
Universitas Airlangga, Indonesia  
E-mail: alfinda-n-k@fst.unair.ac.id

Received 05 January 2017  
Accepted 20 July 2017

### ABSTRACT

This paper reports a study of secondary metabolites contained in *Aquilaria microcarpa*, a species belonging to Thymelaceae. This species has not been investigated so far. *Aquilaria microcarpa* is one of *Aquilaria* species that grows in Indonesia. The sample plant used is taken from east Kalimantan. Extraction of stem bark is done using methanol. Two phenolic compounds, namely 5,3',4'-trihydroxy-7-methoxyflavon or known as 3'-hydroxy genkwanin and 6-hydroxy-2-(2-phenylethyl)chromone are isolated and identified. The chemical structure of these compounds is determined based on spectroscopic data, as well as HR-ESI-MS and NMR spectra. Both compounds were reported previously but they were extracted from another species of *Aquilaria*.

**Keywords:** *Aquilaria microcarpa*, Thymelaeaceae, chromone, flavonoid.

### INTRODUCTION

*Aquilaria* is a genus belonging to Thymelaeaceae family. The species of *Aquilaria* are widely distributed in Asia. Certain trees of *Aquilaria* species produce the fragrant resinous wood, known in the different regions as agarwood, eaglewood, gaharu, kanankoh, jinkoh, chen xiang or tram. Other people also call it aloeswood or agalloch [1, 2]. The *Aquilaria* genus is rich in a variety of different classes of natural products, especially sesquiterpenes and chromones. Flavonoid, benzophenone, diterpenoid, triterpenoid and lignin compounds are also present.

Agarwood preparations are used in Kampo medicine in Japan because of their sedative, analgesic or digestive properties [3]. *Aquilaria* leaves are applied in China topically to treat injuries such as fractures and bruises [4], while in Korea agarwood is used for the treatment of cough, asthma, and as a sedative among others [5]. In Saudi Arabia and other Arabic countries, the wood of *Aquilaria* trees is used as incense at important religious occasions [6, 7].

Some species of *Aquilaria* which are widely studied

are *A. sinensis*, *A. malaccensis*, *A. crassna* and *A. agallocha*, whereas *A. beccariana*, *A. hirta*, *A. cumingiana*, *A. filaria* dan *A. microcarpa* are *Aquilaria* species that grow in Indonesia and are not studied so far.

This paper reports research results referring to *A. microcarpa* as one of these species. Two phenolic compounds are isolated from stem bark of *A. microcarpa*. They are identified as 5,3',4'-trihydroxy-7-methoxyflavon or known as 3'-hydroxy genkwanin (**1**) and 6-hydroxy-2-(2-phenylethyl)chromone (**2**). Their structure of is determined spectroscopically. It is presented in Fig. 1.

### EXPERIMENTAL

Non infected stem bark of *A. microcarpa* obtained from Bukit Bangkirai forest conservation, Samboja, Samarinda, Kalimantan Timur was used.

Methanol, *n*-hexane, ethyl acetate, diisopropyl ether, chloroform, cerium sulfate, acetone, silica gel 60 GF<sub>254</sub> (Merck), silica gel 60 PF<sub>254</sub> (Merck), silica gel 60 GF<sub>254</sub> 0.25 mm (Merck) were the chemical reagents applied.

A rotary vacuum evaporator, column and radial




Fig. 1. Structure of isolated compounds.

chromatography were used in the course of the experimental procedure applied. The latter was as follows. 1.69 kg of stem bark was cut up and then subjected to extraction with n-hexane. The residue was separated from the filtrate and then macerated again using methanol. The extract methanol was fractionated with ethyl acetate. The ethyl acetate fraction was then separated using a column chromatography on silica gel. The fractions of interest were subsequently separated again using several methods of chromatography until a pure product was obtained. Compound 1 was obtained after a double separation using gravity column chromatography followed by a double separation using radial chromatography. Mixtures of hexane - ethyl acetate, hexane - acetone, hexane - chloroform and hexane - diisopropyl ether whose polarity was gradually increased were used as an eluent in those separations. Meanwhile, compound 2 was obtained after a double separation using flash chromatography and a single separation using radial chromatography. A mixture of hexane - ethyl acetate whose polarity was gradually increased was the eluent used in those separations. Compounds 1 and 2 were obtained as much as 3 mg each.

The structure of each compound was elucidated using UV-Vis spectroscopy, HRESI-MS and NMR. UV-Vis Shimadzu 1800, HR-ESI-MS Waters LCT XE ESITOF (electro spray ionization-time of flight), and NMR JEOL ECA 400 ( $^1\text{H-NMR}$ , 400 MHz) and ( $^{13}\text{C-NMR}$ , 100 MHz) were used.

Table 1.  $^1\text{H}$  and  $^{13}\text{C}$ -NMR data of compound 1 and 2.

No	Compound 1 ( $\text{CDCl}_3$ )		Compound 2 (acetone- $d_6$ )	
	$\delta_{\text{H}}$ (m, J Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (m, J Hz)	$\delta_{\text{C}}$
2	-	158,0	-	169,1
3	6,18 (s)	103,9	6,04 (s)	109,7
4	-	183,2	-	182,7
4a	-	102,0	-	124,6
5	-	163,6	7,48 (d, J = 2,1 Hz)	108,9
6	6,32 (d, J = 2,2 Hz)	101,0	-	156,6
7	-	166,7	7,24 (dd, J = 8,3 & 2,1 Hz)	120,2
8	6,46 (d, J = 2,2 Hz)	100,1	6,88 (d, J = 8,3 Hz)	115,8
8a	-	160,5	-	151,2
1'	-	122,9	-	141,3
2'	7,36 (d, J = 2,2 Hz)	119,1	7,27 (m)	129,2
3'	-	147,0	7,27 (m)	129,2
4'	-	139,4	7,27 (m)	129,2
5'	7,54 (d, J = 8,6 Hz)	124,4	7,27 (m)	129,2
6'	7,13 (dd, J = 8,6 & 2,2 Hz)	124,0	7,27 (m)	129,2
7'	-	-	3,08 (m)	33,4
8'	-	-	2,96 (m)	36,4
5-OH	11,14 (s)	-	-	-
7-OCH <sub>3</sub>	3,67 (s)	55,7	-	-

## RESULTS AND DISCUSSION

**5,3',4'-Trihydroxy-7-methoxyflavone (1)** is a pale yellow solid;  $\lambda_{\text{max}}$  (MeOH) nm (log  $\epsilon$ ): 278 (4.09), 287 sh (3.97), and 326 (3.97); HRESI-MS:  $[\text{M-H}]^-$  at  $m/z$  301.0718 in accordance with the molecular formula  $\text{C}_{16}\text{H}_{13}\text{O}_6$ ; the NMR data is showed in Table 1.

**6-Hydroxy-2-(2-phenylethyl)chromone (2)** is a pale yellow solid;  $\lambda_{\text{max}}$  (MeOH) nm (log  $\epsilon$ ): 242 (4.33), 314 (3.58), and 369 (3.39); HRESI-MS:  $[\text{M-H}]^-$  at  $m/z$  265.0805 in accordance with the molecular formula  $\text{C}_{17}\text{H}_{14}\text{O}_3$ ; NMR data is shown in Table 1.

The  $^1\text{H-NMR}$  spectrum of the isolated flavonoid compounds 1 in  $\text{CDCl}_3$  shows two units of aromatic proton signals that refer to three aromatic proton signals of ABX system and two aromatic proton signals of AX system. The three aromatic proton signals of ABX system are attributed to the protons in the chemical shifts



Table 2. Correlation one bond from HMQC analysis.

Compound 1		Compound 2	
$\delta_H$ (mult, J Hz)	$\delta_C$	$\delta_H$ (mult, J Hz)	$\delta_C$
7,54 (1H, d, J = 8,6 Hz)	119,1	7,48 (d, J = 2,1, 1H)	108,9
7,36 (1H, d, J = 2,2 Hz)	124,4	7,27 (m, 5H)	129,3 129,2 127,1
7,13 (1H, dd, J = 8,6 & 2,2 Hz)	124,0	7,24 (dd, J = 8,3 dan 2,1, 1H)	120,2
6,46 (1H, d, J = 2,2 Hz)	100,1	6,88 (d, J = 8,3, 1H)	115,8
6,32 (1H, d, J = 2,2 Hz)	101,0	6,04 (s, 1H)	109,7
6,18 (1H, s)	103,9	3,08 (m, 2H)	33,4
3,67 (3H, s)	55,7	2,96 (m, 2H)	36,4

$\delta_H$  (ppm) of 7.54 ( $d, J = 8.6$  Hz), 7.36 ( $d, J = 2.2$  Hz) and 7.13 ( $dd, J = 8.6$  and 2.2 Hz). The pair of doublet signals in AX system with *meta* coupling constants ( $J = 2.2$  Hz) is attributed to the proton in the chemical  $\delta_H$  (ppm) shifts of 6.46 and 6.32. Those units' signals indicate that the isolated flavonoid compounds are substituted at C-5, C-7, C-3 and C-4'. The singlet signal at  $\delta_H$  6.18 ppm shows that the isolated compound is derived of flavones (Zhao *et al.*, 2005). The  $^1\text{H-NMR}$  spectrum gives also one singlet signal attributed to a methoxy group that appears at  $\delta_H$  (ppm) 3.67. The deshielding proton signal at  $\delta_H$  11.14 ppm characterizes the hydroxyl group at C-5 which is capable of forming a hydrogen bond with the carbonyl group at C-4.

The  $^{13}\text{C-NMR}$  spectrum of (1) shows 16 signals attributed to six methine carbons CH ( $\delta_C$  124.4; 124.0; 119.1; 103.9; 101.0 and 100.1 ppm), one methoxy carbon ( $\delta_C$  55,7 ppm), and nine quaternary carbons ( $\delta_C$  183.2; 166.7; 163.6; 160.5; 158.0; 147.0; 139.4; 122.9 and 102.0 ppm). Six of the latter are aryloxy carbons ( $\delta_C$  166.7; 163.6; 160.5; 158.0; 147.0; and 139.4 ppm). The presence of a carbonyl carbon atom is evidenced by a signal of  $\delta_C$  183.2 ppm. It follows that there is only one methoxy group. The three other substituents refer to hydroxy groups.  $^1\text{H}$  and  $^{13}\text{C-NMR}$  data of compound 1 is listed in Table 1.

The  $^1\text{H-NMR}$  spectrum of isolated compound (2) in acetone- $d_6$  shows two units of aromatic proton signals referring to three ABX system aromatic proton signals and five monosubstituted aromatic proton signals. The

three ABX system aromatic proton signals appear in the chemical shifts  $\delta_H$  of 6.88 ( $d, J = 8.3$  Hz), 7.24 ( $dd, J = 8.3$  and 2.1 Hz), and 7.48 ( $d, J = 2.1$  Hz). The proton signals of the other aromatic unit (five protons) appear as a multiplet at  $\delta_H$  7.27 ppm. The proton singlet signal at  $\delta_H$  6.04 ppm and the two proton multiplet signals at  $\delta_H$  3.08 and 2.96 show a characteristic chromone class of compounds that have a hydroxy substituent at C-6 or C-7.

The  $^{13}\text{C-NMR}$  spectrum analysis shows fifteen carbons that are perfectly separated. Based on HR - ESI - MS analysis there should be seventeen carbons. This suggests the existence of two symmetric carbons. Two symmetric carbons are located in ring B at C-2' / 6' and C-3' / 5'. Fifteen carbon signals refer to six quaternary carbons ( $\delta_C$  (ppm) 182.7; 168.9; 156.6; 151.2; 141.2; 124.6), seven methine carbons CH ( $\delta_C$  (ppm) 109.7; 108.9; 120.2; 115.8; 129.2; 127.1; 129.3) and two methylene carbons  $\text{CH}_2$  ( $\delta_C$  (ppm) 33.4 and 36.4). Four of these six quaternary carbon signals consist of three aryloxy carbon signals ( $\delta_C$  (ppm) 169.2; 156.6; 151.2) and one of carbonyl carbon signal ( $\delta_C$  182.7 ppm). This indicates that compound 2 has the structure of 2-(2-phenylethyl)chromone with one hydroxyl substituent. The  $^1\text{H}$  and  $^{13}\text{C-NMR}$  data of compound 2 is presented in Table 1. A correlation of one bond in the HMQC spectrum analysis of compound 1 shows seven correlations as presented in Table 2. The same correlation occurs also in compound 2.

The presence of a methoxy and hydroxyl groups in the structure of compound (1) is evidenced by the HMBC spectrum. It is confirmed that the two hydroxyl groups in ring B are positioned at 3' and 4'. The position of the methoxy group and the remaining hydroxyl group in ring A has to be determined. The positions to be filled refer to numbers 5 and 7. The proton signal of the hydroxyl group ( $\delta_H$  of 11.14 ppm) shows correlations with two quaternary carbon signals ( $\delta_C$  of 163.6 ppm and 102.0 ppm) and one methine carbon signal at  $\delta_C$  of 101.0 ppm. It has been stated above that a deshielding proton signal at  $\delta_H$  of 11.14 ppm is characteristic for the hydroxyl group at C-5 which is capable of forming a hydrogen bond with the carbonyl group at C-4. Hence it follows that the signal at 163.6 ppm refers to C-5, that at 102.0




Fig. 2. HMBC correlation of compound **1** and compound **2**.

ppm refers to C-4a, while that at 101.0 ppm refers to C-6. The singlet proton signal of the methoxy group ( $\delta_{\text{H}}$  of 3.67 ppm) gives one correlation with the quaternary carbon signal at  $\delta_{\text{C}}$  of 166.7 ppm. The latter is a signal referring to C-7. The results of HMBC spectrum analysis of compound **1** are illustrated in Fig. 2.

HMBC analysis is required to verify the hydroxy group position (at C-6 or C-7) in compound **2** structure. A singlet proton signal at  $\delta_{\text{H}}$  of 6.04 ppm (H-3) shows correlations with two quaternary carbon signals at  $\delta_{\text{C}}$  of 169.1 ppm (C-2) and 124.6 ppm (C-4a) and one methine carbon at  $\delta_{\text{C}}$  of 36.4 ppm (C-8'). Another correlation is revealed between the aromatic proton in ABX system and the two aryloxy carbons and one quaternary carbon. The proton considered appears as a doublet with a coupling constant of 2.1 Hz at  $\delta_{\text{H}}$  of 7.48 ppm (H-5). The two aryloxy carbons mentioned above show signals at  $\delta_{\text{C}}$  of 156.6 (C-6), 151.2 (C-8a) ppm respectively, while the quaternary carbon appears at  $\delta_{\text{C}}$  of 124.6 ppm (C-4a). A correlation is also observed between the doublet ( $J = 8.3$  Hz) proton signal at  $\delta_{\text{H}}$  of 6.88 ppm (H-8) with one aryloxy carbon at  $\delta_{\text{C}}$  of 151.2 ppm (C-8a) and one methine carbon at  $\delta_{\text{C}}$  of 120.2 (C-7). The overall correlations data indicates that the hydroxyl group is attached at C-6. The results of HMBC spectrum analysis of compound **2** are shown in Fig. 2.

All data obtained is compared to that found in the

literature. Compound **1** is reported by Qi [8] who succeeds isolating this compound from the leaves of *A. sinensis*. In that report, Qi mentions also that this 3'-hydroxy genkwanin has a significant inhibitory activity in respect to neutrophils respiratory burst stimulated by PMA with  $\text{IC}_{50}$  value  $0.80 \pm 0.13$   $\mu\text{mol/l}$ . Likewise, compound **2** is compared to the same compound isolated [9] from an ether extract of powdered agarwood from Kalimantan. In 1989 Yang [10] is the first to obtain this compound from the ether soluble fraction of an alcoholic extract of *A. sinensis* (Lour.) Gilg. The same compound is also successfully isolated from a 70 % MeOH extract of *A. malaccensis* agarwood chips [11] and from an EtOAc extract of Chinese agarwood induced by artificial holing from *A. sinensis* [12]. The latter reference mentions also that this compound exhibits acetylcholinesterase inhibitory activity with percentage of inhibition of  $19.3 \pm 0.8$  %. Bioactivity tests referring to compound **1** are not performed in the course of the present investigation because of the small quantity isolated. However an anticancer activity test against T47d cell line is carried out with the participation of compound **2**. The result indicated that the  $\text{IC}_{50}$  is 2884.03  $\mu\text{g/mL}$ . The same test is also carried out with the ethyl acetate fraction used for the isolation of compound **1** and compound **2**. The result shows that this fraction has a weak activity with  $\text{IC}_{50}$  value of  $26.48 \pm 0.02$   $\mu\text{g/mL}$ .

## CONCLUSIONS

A plant sample of *Aquilaria microcarpa* was used to isolate two compounds. Their structure was determined based on analyses, including UV, HR-ESI-MS and NMR. The compounds obtained were identified as 5,3',4'-trihydroxy-7-methoxyflavon or known as 3'-hydroxy genkwanin **1** and 6-hydroxy-2-(2-phenylethyl)chromon **2**. They were isolated by other research groups using *A. sinensis* and *A. malaccensis*, which are different species of *Aquilaria*.

## Acknowledgements

This work was financially supported by 2016 grant from the Ministry of Research and Higher Education. The authors are grateful to Bukit Bangkirai Forest Conservation, Samboja, Samarinda, Kalimantan Timur for the sample material provision.

## REFERENCES

1. W. Li, C.H. Cai, W.H. Dong, Z.K. Guo, H. Wang, W.L. Mei, H.F. Dai, 2-(2-phenylethyl) chromone derivatives from Chinese agarwood induced by artificial holing, *Fitoterapia*, 98, 2014, 117-123.
2. J. Ueda, L. Imamura, Y. Tezuka, Q.L. Tran, M. Tsuda, S. Kadota, New sesuiterpene from Vietnamese agarwood and its induction effect on brain-derived neurotrophic factor mRNA expression in vitro, *Bioorg. Med. Chem.*, 14, 2006, 3571-3574.
3. T. Yagura, N. Shibayama, M. Ito, F. Kiuchi, G. Honda, *Tetrahedron Lett.*, 46, 2005, 4395-4398.
4. M. Zhou, H. Wang, Suolangjiba, J. Kou, B. Yu, Antinociceptive and anti-inflammatory activities of *Aquilaria sinensis* (Lour.) Gilg. leaves extract, *J. Ethnopharmacol.*, 117, 2008, 345-350.
5. Y.C. Kim, E.H. Lee, Y.M. Lee, H.K. Kim, B.K. Song, E.J. Lee, H.M. Kim, Effect of the aqueous extract of *Aquilaria agallocha* stems on the immediate hypersensitivity reactions, *J. Ethnopharmacol.*, 58, 1997, 31-38.
6. H.Z. Alkhathlan, H.M. Al-Hazimi, F.S. Al-Dhalaan, A.A. Mousa, Three 2-(2-phenylethyl) chromones and two terpenes from agarwood, *Nat. Prod. Res.*, 19, 4, 2005, 367-372.
7. D.L. Yang, H. Wang, Z.K. Guo, W.H. Dong, W.L. Mei, H.F. Dai, A new 2-(2-phenylethyl) chromone derivative in Chinese agarwood "Qi-Nan" from *Aquilaria sinensis*, *J. Asian Nat. Prod. Res.*, 16, 7, 2014, 770-776.
8. J. Qi, J.J. Lu, J.H. Liu, B.Y. Yu, Flavonoids and a rare benzophenone glycoside from the leaves of *Aquilaria sinensis*, *Chem. Pharm. Bull.*, 57, 2, 2009, 134-137.
9. Y. Shimada, T. Tominaga, T. Konishi, S. Kiyosawa, Studies on the agarwood (Jinko). I. Structures of 2-(2-phenylethyl) chromone derivatives, *Chem. Pharm. Bull.*, 30, 10, 1982, 3791-3795.
10. J.S. Yang, Y.L. Su, Studies on the chemical constituents of *Aquilaria sinensis* (Lour.) Gilg. IV. Isolation and characterization of 2-(2-phenylethyl) chromone derivatives, *Yao Xue Xue Bao*, 24, 9, 1989, 678-683.
11. B. Wu, S.W. Kwon, G.S. Hwang, J.H. Park, Eight new 2-(2-phenylethyl) chromone (=2-(2-phenylethyl)-4H-1-benzopyran-4-one) derivatives from *Aquilaria malaccensis* agarwood, *Helv. Chim. Acta*, 95, 2012, 1657-1665.
12. W. Li, C.H. Cai, W.H. Dong, Z.K. Guo, H. Wang, W.L. Mei, H.F. Dai, 2-(2-Phenylethyl) chromone derivatives from Chinese agarwood induced by artificial holing, *Fitoterapia*, 98, 2014, 117-123.



# PHENOLIC COMPOUNDS FROM *Aquilaria microcarpa* STEM BARK

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	3%
2	<a href="http://www.ekouhou.net">www.ekouhou.net</a> Internet Source	1%
3	Lin Yang, Lirui Qiao, Dan Xie, Yuhe Yuan, Naihong Chen, Jungui Dai, Shunxing Guo. "2-(2-Phenylethyl)chromones from Chinese eaglewood", <i>Phytochemistry</i> , 2012 Publication	1%
4	<a href="http://www.freshpatents.com">www.freshpatents.com</a> Internet Source	1%
5	<a href="http://tpauxpages)%2fec506a59176be044c1256e9e00unrise1.org">tpauxpages)%2fec506a59176be044c1256e9e00unrise1.org</a> Internet Source	1%
6	Maya Mori-Hongo, Hiroyuki Takimoto, Takayuki Katagiri, Makoto Kimura, Yu Ikeda, Toshio Miyase. " Melanin Synthesis Inhibitors from ", <i>Journal of Natural Products</i> , 2009 Publication	1%

7	Ibrahim, Sabrin R.M., and Gamal A. Mohamed. "Natural occurring 2-(2-phenylethyl) chromones, structure elucidation and biological activities", <i>Natural Product Research</i> , 2014.	1%
Publication		
8	Taha Sarg. "Phytochemical and Pharmacological Studies of <i>Dalbergia sissoo</i> Growing in Egypt", <i>Pharmaceutical Biology (Formerly International Journal of Pharmacognosy)</i> , 1/1/1999	1%
Publication		
9	Wei Li, Cai-Hong Cai, Wen-Hua Dong, Zhi-Kai Guo, Hao Wang, Wen-Li Mei, Hao-Fu Dai. "2-(2-Phenylethyl)chromone derivatives from Chinese agarwood induced by artificial holing", <i>Fitoterapia</i> , 2014	<1%
Publication		
10	<a href="http://www.cropwatch.org">www.cropwatch.org</a>	<1%
Internet Source		
11	<a href="http://spandidos-publications.com">spandidos-publications.com</a>	<1%
Internet Source		
12	<a href="http://www.forestfocus.org">www.forestfocus.org</a>	<1%
Internet Source		
13	Tjahjandarie, Tjitjik, Ratih Saputri, and Mulyadi Tanjung. "Methyl 2,5-Dihydroxy-4-(3'-methyl-2'-butenyl)benzoate", <i>Molbank</i> , 2016.	<1%



14

Huang, Y.L.. "Antioxidant flavonoids from the rhizomes of *Helminthostachys zeylanica*", *Phytochemistry*, 200312

Publication

---

<1%

15

Wu, Bo, Sung Won Kwon, Gwi Seo Hwang, and Jeong Hill Park. "Eight New 2-(2-Phenylethyl)chromone (=2-(2-Phenylethyl)-4H-1-benzopyran-4-one) Derivatives from *Aquilaria malaccensis* Agarwood", *Helvetica Chimica Acta*, 2012.

Publication

---

<1%

16

[beilstein-journals.org](http://beilstein-journals.org)

Internet Source

---

<1%

17

[repository.unair.ac.id](http://repository.unair.ac.id)

Internet Source

---

<1%

18

"Fogacin, a Novel Cyclic Octaketide Produced by *Streptomyces* Strain Tü 6319 †", *The Journal of Antibiotics*, 05/2006

Publication

---

<1%

19

[www.patents.com](http://www.patents.com)

Internet Source

---

<1%

20

[patentscope.wipo.int](http://patentscope.wipo.int)

Internet Source

---

<1%

21

[data.epo.org](http://data.epo.org)

Internet Source

<1%

22

[www.thieme-connect.com](http://www.thieme-connect.com)

Internet Source

&lt;1%

23

Ge Liao, Wen-Li Mei, Wen-hua Dong, Wei Li, Pei Wang, Fan-Dong Kong, Cui-Juan Gai, Xi-Qiang Song, Hao-Fu Dai. "2-(2-Phenylethyl)chromone derivatives in artificial agarwood from *Aquilaria sinensis*", *Fitoterapia*, 2016

Publication

&lt;1%

24

Jeong Seon Yoon, Mi Kyeong Lee, Sang Hyun Sung, Young Choong Kim. "Neuroprotective 2-(2-Phenylethyl)chromones of ", *Journal of Natural Products*, 2006

Publication

&lt;1%

25

Hashim, Yumi Zuhanis Has-Yun, Philip G. Kerr, Phirdaous Abbas, and Hamzah Mohd Salleh. "Aquilaria spp. (agarwood) as source of health beneficial compounds: A review of traditional use, phytochemistry and pharmacology", *Journal of Ethnopharmacology*, 2016.

Publication

&lt;1%

26

Bai, Laxinamujila, Hai Ying Bao, and Tolgor Bau. "Isolation and identification of a new benzofuranone derivative from *Usnea longissima*", *Natural Product Research*, 2014.

Publication

&lt;1%

27

Lin Yang, Lirui Qiao, Chengxue Ji, Dan Xie, Ning-Bo Gong, Yang Lu, Jianjun Zhang, Jungui Dai, Shunxing Guo. "Antidepressant Abietane Diterpenoids from Chinese Eaglewood", *Journal of Natural Products*, 2013

Publication

---

&lt;1%

28

Fozia A. Adem, Victor Kuete, Armelle T. Mbaveng, Matthias Heydenreich et al. "Cytotoxic benzylbenzofuran derivatives from *Dorstenia kameruniana*", *Fitoterapia*, 2018

Publication

---

&lt;1%

29

Reza Alavi, Seyed Hamid, Narguess Yassa, Abbas Shafiee, and Fatemeh Fouladi. "A New Furanocoumarin from *Peucedanum ruthenicum*.", *Pharmaceutical Biology*, 2008.

Publication

---

&lt;1%

30

Swarna A. Gamage, Nisana Tepsiri, Prapon Wilairat, Stanley J. Wojcik, David P. Figgitt, Raymond K. Ralph, William A. Denny. "Synthesis and in vitro Evaluation of 9-Anilino-3,6-diaminoacridines Active Against a Multidrug-Resistant Strain of the Malaria Parasite *Plasmodium falciparum*", *Journal of Medicinal Chemistry*, 1994

Publication

---

&lt;1%

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      On



# PHENOLIC COMPOUNDS FROM *Aquilaria microcarpa* STEM BARK

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---