

DAFTAR PUSTAKA

- Katz, B. H., Quencer, R. v. & Hinks, R. S., 1989. Comparison of Gradient-Recalled-Echo and T2-Weighted Spin-Echo Pulse Sequences in Intramedullary Spinal Lesions. *AJNR*, pp. 815-821.
- Algra, P. R., Valk, J. & Heimans, J. J., 1998. Diagnosis and Therapy of Spinal Tumors. In: *Diagnosis and Therapy of Spinal Tumors*. Berlin: Springer, p. 72.
- Dura, T. S., 2008. *Atlas Of Anatomy By Sectional Imaging*. Berlin: Bayer Health Care.
- Fatimah, Dahjono, J. & Sativa, M. R., 2016. Optimasi Field of View (FOV) Terhadap kualitas Citra Pada T2Wi FSE MRI Lumbal Sagital. pp. 1-5.
- Fleischman, D. et al., 2012. Cerebrospinal Fluid Pressure Decrease with Older Age. *Plos One*, VII(12), pp. 1-9.
- Hutapea & Marintan, D., 2017. Analisis Kualitas Citra Lumbal pada Pengaruh Perubahan Nilai Time Repetition dan Time Echo Untuk Kasus Hernia Nukeus Pulposus dengan Menggunakan Magnetic Resonance Imaging, Medan: Universitas Sumatera Utara.
- Japardi, I., 2002. *Radikulopati Thorakalis*, Medan: USU Digital Library.
- Leon, P. A. et al., 2016. Comparison of Sagittal FSE T2, STIR, and T1-Weighted Phase-Sensitive Inversion Recovery in the Detection of Spinal Cord Lesions in MS at 3T. *AJNR*, pp. 1-6.

- Leon, P. A. et al., 2016. Comparison of Sagittal FSE T2, STIR, and T1-Weighted Phase-Sensitive Inversion Recovery in The Detection of Spinal Cord Lesion. *American Journal Neuroradiology*, pp. 970-974.
- Moulopoulos, L. A. & Koutoulidis, V., 2015. Bone Marrow MRI. In: *Bone Marrow MRI*. Italia: Springer, p. 1.
- Muzamil, A. & Firmansyah, A. H., 2017. Optimization Image of Magnetic Resonance Imaging (MRI) T2 Fast Spin Echo (FSE) with Variation Echo Train Length (ETL) on the Rupture Tendon Achilles Case. *International Conference on Physical Instrumentation and Advanced Materials*, pp. 1-9.
- Nakatsu, M. et al., 2000. Comparison of Short Time Inversion Recovery (STIR) and Fat-Saturated (chemsat) Techniques for Bckground Fat Intensity Suppression in Cervical and Thoracic MR Imaaging. *Journal of Magnetic Resonance Imaging*, pp. 56-60.
- Papinutto, N. et al., 2015. Age, Gender and Normalization Covariates for Spinal Cord Gray Matter and Total Cross-Sectional Areas at Cervical and Thoracic Levels: A 2D Phase Sensitive Inversion Recovery Imaging Study. *Plos One*.
- Pfirmann, C. W. et al., 2006. Effect of Aging and Degeneration on Disc Volume and Shape: A Quantitive Studiyy in Assynmpomatic Volunteers. *Journal of Orthopaedic Research*, Volume XXIV, pp. 1086-1094.
- Serdamayanti & Hidayat , S., 2002. *Metodologi Penelitian*. s.l.:Mandar Maju.

- Shah, L. M. & Hanrahan, C. J., 2011. MRI of Spinal Bone Marrow: Part I, Techniques and Normal Age-Related Appereances. *American Journal of Radiology*, pp. 1298-1308.
- Simanjuntak, J., Nur, M. & Hidayanto, E., 2014. Studi Analisis Echo Train Length Dalam K-Space serta Pengaruhnya Terhadap Kualitas Citra Pembobotan T2 FSE pada MRI 1.5T. *Berkala Fisika*, Volume XVII, pp. 7-12.
- Supranto, J., 2000. *Teknik Sampling untuk Survei dan Eksperimen*. Jakarta: Rineka Cipta.
- Westbrook, C., 2008. *Handbook of MRI Technique Third Edition*. United Kingdom: Willey Blackwell.
- Westbrook, C., 2014. *Handbook Of MRI Technique Fourth Edition*. United Kingdom: Willey Blackwell.
- Westbrook, C., Roth, C. K. & John, T., 2011. *MRI in Practice Fourth Edition*. United Kingdom: Willey Blackwell.
- Westbrook, C. & Talbot, J., 2019. *MRI In Practice Fifth Edition*. United Kingdom: Willey Blackwell.
- Wijojongko, S. et al., 2016. *Protokol Radiologi CT Scan dan MRI*. Magelang: Inti Medika Pustaka.
- Zhao, J. et al., 2009. MRI of the Spine : Image Quality and Normal Bone Marrow Contrast at 3T Versus 1,5T. *American Journal of Radiology*, pp. 873-880.