

**DAFTAR PUSTAKA**

- Abad ANA, Nouri MHK, Gharjanie A, *et al*, 2011, Effect of Matricaria chamomilla Hydroalcoholic Extract on Cisplatin-induced neuropathy in mice, *Chin J Nat Med*; 9(2): 126-131.
- Abad ANA, Nouri MHK, Tavakoli F, 2011, Effect of Salvia officinalis Hydroalcoholic Extract on Vincristine-induced Neuropathy in Mice, *Chin J Nat Med*; 9(5): 354-8.
- Adam-Vizi, V. and Chinopoulos, C, 2006, Bioenergetics and the formation of mitochondrial reactive oxygen species, *Trends Pharmacol. Sci*, 27, 639–645.
- Addington, J. Dan Freimer, M., 2016. Chemotherapy-induced peripheral neuropathy: an update on the current understanding. *F1000 Faculty Review*, 5 (1466), PP. 1-7.
- Apkarian, A.V., Bushnell, M.C., Treede, R.D., Zubieta, J.K., 2005, Human brain mechanisms of pain perception and regulation in health and disease, *Eur. J. Pain* 9, 463–484.
- Andreyev, A. Y., Kushnareva, Y. E. & Starkov, A. A, 2005, Mitochondrial metabolism of reactive oxygen species, *Biochemistry (Moscow)* 70, 200–214.
- Areti Aparna, Prashanth Komirishetty, Anil Kumar Kalvala, *et al*, 2018, Rosmarinic Acid Mitigates Mitochondrial Dysfunction and Spinal Glial Activation in Oxaliplatin-Induced Peripheral Neuropathy.
- Argyriou Andreas A, Athanasios P Kyritsis, Thomas Makatsoris, *et al*, 2014, Chemotherapy-Induced Peripheral Neuropathy in Adults, Comprehensive Update of the Literature, *Cancer Management and Research*.
- Azevedo, M. I., Pereira, A. F., Nogueira, R. B., Rolim, F. E., Brito, G. A., Wong, D. V. T., Vale, M. L. (2013). The Antioxidant Effects of the Flavonoids Rutin and Quercetin Inhibit Oxaliplatin-Induced Chronic Painful Peripheral Neuropathy. **Molecular Pain**, 9, 1744–8069–9–53.

- Balaban, R. S., Nemoto, S. & Finkel, T, 2005, Mitochondria, oxidants, and aging, *Cell* 120, 483–495.
- Balayssac, D., Ferrier, J., Descoeur, J., Ling, B., Pezet, D., Eschalier, A., et al, 2011, Chemotherapy-induced peripheral neuropathies: from clinical relevance to preclinical evidence, *Expert Opin. Drug Saf.* 10, 407–417. doi: 10.1517/14740338.2011.543417.
- Baliki, M.N., Geha, P.Y., Apkarian, A.V., Chialvo, D.R., 2008. Beyond feeling: chronic pain hurts the brain, disrupting the defaultmode network dynamics. *J. Neurosci.* 28, 1398–1403.
- Basbaum AI, Bautista DM, Scherrer G, et al, 2009, Cellular and molecular mechanisms of pain. *Cell*;139:267–284.
- Bhadri N, Sanji T, Guggilla HM, et al, 2013, Amelioration of Behavioural, Biochemical, and Neurophysiological Deficits by Combination of Monosodium Glutamate with Resveratrol/Alpha-Lipoic Acid/Coenzyme Q10 in Rat Model of Cisplatin-Induced Peripheral Neuropathy, *The Scientific World Journal*.
- Boatright Kelly M & Guy S Salvesen, 2003, Mechanism of Caspase activation. *Current Opinion in Cell Biology*, 15:725-731.
- Boyette-Davis, J., Xin, W., Zhang, H., et al, 2011, *Intraepidermal nerve fiber loss corresponds to the development of taxol-induced hyperalgesia and can be prevented by treatment with minocycline*. *Pain* 152, 308–313. doi: 10.1016/j.pain.2010.10.030.
- Boyette-Davis JA, Cata JP, Driver LC, et al, 2013, *Persistent chemoneuropathy in patients receiving the plant alkaloids paclitaxel and vincristine*, *Cancer Chemother Pharmacol* 71:619-626.
- Budihardjo Imawati, Holt Oliver, Michael Lutter, Xu Luo, et al, 1999, *Biochemical Pathways of Caspase Activation during Apoptosis*. *Howard Hughes Medical Institute and Department of Biochemistry*, University of Texas Southwestern Medical Center, Dallas, Texas 75235; Annu, Rev, *Cell Dev, Biol*, 1999, 15:269-90.
- Burgos, E., Gomez-Nicola, D., Pascual, D., Martin, M. I, et al, 2012, *Cannabinoid agonist WIN 55,212-2 prevents the development of paclitaxel-induced peripheral neuropathy in rats*, Possible

- involvement of spinal glial cells. *Eur. J. Pharmacol.* 682, 62–72, doi: 10.1016/j.ejphar,2012,02,008.
- Cadenas, E. and Davies, K. J., 2000, Mitochondrial free radical generation, oxidative stress, and aging, *Free Radical Biol. Med.*, 29, 222–230.
- Casey, K.L., 1999, Forebrain mechanisms of nociception and pain: analysis through imaging, *Proc. Natl. Acad. Sci. U. S. A.* 96, 7668–7674.
- Chance, B., Sies, H. and Boveris, A, 1979, Hydroperoxide metabolism in mammalian organs. *Physiol. Rev.* 59, 527–605.
- Chollet P, Bensmaïne MA, Brienza S, et al. Single agent activity of oxaliplatin in heavily pretreated advanced epithelial ovarian cancer. *Ann Oncol* 1996, 7, 1065–1070.
- Conklin, K. A. (2004) ‘Chemotherapy-associated oxidative stress: Impact on Chemotherapy effectiveness’, *Integrative Cancer Therapies*, 3(4), pp. 294–300. doi: 10.1177/1534735404270335.
- Costigan M, Scholz J, & Woolf CJ, 2009, *Neuropathic pain: a maladaptive response of the nervous system to damage.* Annu Rev Neurosci;32:1–32. 9.
- Culy CR, Clemett D, Wiseman LR. Oxaliplatin. A review of its pharmacological properties and clinical efficacy in metastatic colorectal cancer and its potential in other malignancies. *Drugs* 2000, 60, 895–924.
- deCharms, R.C., Maeda, F., Glover, G.H., Ludlow, D., Pauly, J.M., Soneji, D., Gabrieli, J.D., Mackey, S.C., 2005, Control over brain activation and pain learned by using real-time functional MRI, *Proc. Natl. Acad. Sci. U. S. A.* 102, 18626–18631.
- de Gramont A, Vignoud J, Tournigand C, et al. Oxaliplatin with high-dose leucovorin and 5-fluorouracil 48-hour continuous infusion in pretreated metastatic colorectal cancer. *Eur J Cancer* 1997, 33, 214–219.

- de Gramont A, Figer A, Seymour M, et al. Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol* 2000, 18, 2938–2947.
- Decosterd Isabelle and Clifford J. Woolf, 2000, Spared nerve injury: an animal model persistent peripheral neuropathic pain, *Pain*, 149-158.
- Dong, Y., Wang, J., Feng, D., Qin, H., Wen, H., Yin, Z., Gao, G., & Li, C., 2014. Protective effect of quercetin against oxidative stress and brain edema in an experimental rat model of subarachnoid hemorrhage. *Journal of Medical Science*, 11(3), pp. 282-290.
- Dubner R & Ruda MA, 1992, *Activity-dependent neuronal plasticity following tissue injury and inflammation*, Trends Neurosci;15: 96–103. 6.
- Erichsen Helle K and Gordon Blackburn-Munro, 2002, Pharmacological characterisation of the spared nerve injury model of neuropathic pain, *Pain*, 151-161.
- Fallon M.T., 2013, Neuropathic pain in cancer, *British Journal Anaesthesia* 111(1): 105-11, doi:10.1093/bja/aet208.
- Ferrandiz ML & Alcaraz MJ, 1991, *Anti-inflammatory activity and inhibition of arachidonic acid metabolism by flavonoids*, *Agents Actions*;32:283–288.
- Flor, H., Turk, D.C., 2006. Cognitive and learning aspects. In: McMahon, S.B., Koltzenburg,M. (Eds.), *Textbook of Pain*. Elsevier, Amsterdam, pp. 241–258.
- Fuji, J., Y. Iuchi, S. Matsuki, and T. Ishii. 2003. Cooperative function of antioxidant and redox system against oxidative stress in male reproductive tissue. *Asian J. Androl.* 5: 231-242.
- Gallagher, R.M., Verma, S., 2004. Mood and anxiety disorders in chronic pain. *Prog. Pain Res. Manag.* 27, 139–178.
- Gao W et al, 2016, Quercetin ameliorates paclitaxel-induced neuropathic pain by stabilizing mast cells, and subsequently blocking PKC $\epsilon$ -dependent activation of TRPV1, *Acta Pharmacologica Sinica*; 37: 1166–1177.

- Gata J.P, H.-R. Weng, B N. Lee, J. M. Et al, 2006, *Clinical and Experimental Finding in Humans and Animals with Chemotherapy-Induced Peripheral Neuropathy*, Department of Anesthesiology and Pain Medicine, 72 : 51-69.
- George Paxinos and Keith B.J. Franklin, 2001, Te Mouse Brain in Stereotaxic Coordinates, second edition, Academic Press, Sydney Australia, Halaman 49 dan 77.
- Glordano Catia, Livio Luongo, Lulgia Cristina, Stefania Petrosino *et al*, 2011, TRPV1-Dependent and Independent Alterations in the Limbic Cortex of Neuropathic Mice: Impact on Glial Caspases and Pain Perception, Cerebral Cortex, doi: 10.1093/cercor/bhr328.
- Goel, S., Goldberg, G. L., Kuo, D. Y., Muggia, F., Arezzo, J., and Mani, S, 2008, Novel neurosensory testing in cancer patients treated with the epothilone B analog, ixabepilone, Ann. Oncol. 19, 2048–2052. doi: 10.1093/annonc/ mdn420.
- Guindon J, Deng L, Fan B, Wager-Miller J et al, 2014, Optimization of a cisplatin model of chemotherapy-induced peripheral neuropathy in mice: use of vitamin C and sodium bicarbonate pretreatments to reduce nephrotoxicity and improve animal health status, *Molecular Pain*, 10: 56-69.
- Gupta, R., & Bhaskar, A 2016, ‘Chemotherapy-induced peripheral neuropathic pain’, *BJA Education*, 16(4), pp. 115–119. doi:10.1093/bjaed/mkv044
- Han yaqin and Maree T. Smith., 2013, Pathobiology of cancer chemotherapy-induced peripheral neuropathy (CIPN), Frontiers in Pharmacology, volume 4, hal.5
- Hershman DL, Lacchetti C, Dworkin RH, Smith EML et al. 2014. Prevention and Management of Chemotherapy-Induced Peripheral Neuropathy in Survivors of Adult Cancers, *American Society of Clinical Oncology Clinical Practice Guideline*. J Clin Oncol; 32: 1941-67.
- Herzog, Christian, Gur P.Kaushal, Randy S. Haun, et al, 2014, *Caspase Protocols in Mice*. Department of International Medicine, Method Mol Biol ; 1133 : 141-154.

- Hoy, S. M, 2013, Subcutaneous bortezomib: in multiple myeloma. *Drugs* 73, 45–54. doi: 10.1007/s40265-013-0006-6.
- Hsieh, J.C., Belfrage, M., Stone-Elander, S., Hansson, P., Ingvar, M. (1995). Central representation of chronic ongoing neuropathic pain studied by positron emission tomography. *Pain*, 63, pp. 225–236.
- Ji RR, Kohno T, Moore KA, et al, 2003, *Central sensitization and LTP: do pain and memory share similar mechanisms?* Trends Neurosci;26:696–705. 10.
- Kannarkat G, Lasher EE, Schiff D, Neurologic complication of chemotherapy agents, *Current Opinion in Neurology*, 2007;20(6):719-25.
- Kauffman Scott H & William C. Earnshaw, 2000, Induction of apoptosis by cancer chemotherapy, *Experimental Cell Research* 256, 42-49.
- Kaushal, V., Herzog, C., Haun, R. S., & Kaushal, G. P. (2014). Caspase protocols in mice, *Methods in molecular biology* (Clifton, N.J.), 1133, 141-54.
- Kocer, B., Sucak, G., Kuruoglu, R., Aki, Z., Haznedar, R., and Erdoganmus, N. I, 2009, Clinical and electrophysiological evaluation of patients with thalidomide-induced neuropathy, *Acta Neurol. Belg.* 109, 120–126.
- Lee, J., N. Koo, and D. B. Min. 2004. Reactive oxygen species, aging, and antioxidative nutraceuticals. *Compre Rev. In food sci. And food safety.* 3: 21-33.
- Levi F, Misset JL, Brienza S, et al. A chronopharmacologic phase II clinical trial with 5-fluorouracil, folinic acid, and oxaliplatin using an ambulatory multichannel programmable pump. High antitumor effectiveness against metastatic colorectal cancer. *Cancer* 1992, 69, 893–900.
- Li P, Nijhawan D, Budihardjo I, Srinivasula SM, Ahmad M, et al, 1997, Cytochrome c and dATP-dependent formation of Apaf-1/caspases-9 complex initiates an apoptotic protease cascade, *Cell* 91:479-89.
- Mackey, S.C., Maeda, F., 2004, Functional imaging and the neural systems of chronic pain. *Neurosurg. Clin. N. Am.* 15, 269–288.

- Makker P, G, S., Samuel S, Duffy., justin G. Lees *et al.*, 2017, characterisation of immune and neuroinflammatory changes associated with chemotherapy-induced peripheral neuropathy, research article, PLOS ONE | DOI:10.1371/journal.pone.0170814.
- Mantyh PW, Clohisy DR, Koltzenburg M, et al, 2002, *Molecular mechanisms of cancer pain*, Nat Rev Cancer;2:201–209.
- Martini FH (2005) Anatomy and physiology, 1st ed. Pearson Education, Inc
- Masocha, W, 2015, Astrocyte activation in the anterior cingulate cortex and altered glutamatergic gene expression during paclitaxel-induced neuropathic pain in mice, PeerJ 3:e1350. doi: 10.7717/peerj.1350
- Mayer, E.A., Naliboff, B.D., Craig, A.D., 2006, Neuroimaging of the brain-gut axis: from basic understanding to treatment of functional GI disorders, Gastroenterology 131, 1925–1942.
- Meacham Kathleen., Andrew Shepherd., Durga P. Mohapatra and Simon Haroutounian., 2017., Neuropathic Pain: Central vs. Peripheral Mechanisms, Curr Pain Headache Rep 21:28, DOI 10.1007/s11916-017-0629-5.
- Mira, L., Fernandez, M. T., Santos. M., Rocha, R., Florencio, M. H., Jennings, K.R., 2002. Interaction of flavonoids with iron and copper ions; a mechanism for their antioxidant activity. Free Radical Research 36, 1199-1028.
- Moudi, M., Go, R., Yien, C.Y.S, & Nazre, M., 2013, Vinca Alkaloids, *International Journal of Preventive Medicine*, 4(11), PP. 1231-1235.
- Muller, F, 2000, The nature and mechanism of superoxide production by the electron transport chain. J. Am, Aging Assoc, 23, 227–253.
- National Institute for Health and Care Excellent, 2017, Neuropathic pain
- Neugebauer Volker, Vasco Galhardo, Sabatino Maione and Sean C. Mackey, 2009, Forebrain Pain Mechanism, Brain Research Reviewer, 226-242.

- Ohara, P.T., Vit, J.P., Jasmin, L., 2005. Cortical modulation of pain. *Cell. Mol. Life Sci.* 62, 44–52.
- Park, H.J., 2014, Chemotherapy induced peripathic *pain*, *Korean Journal anesthesiol*, 67 (1), PP. 4-7
- Pérez-Garijo, A. (2017). When dying is not the end: Apoptotic caspases as drivers of proliferation. *Seminars in Cell & Developmental Biology*. doi:10.1016/j.semcd.2017.11.036.
- Peyron R, Laurent B, Garcia-Larrea L. (2000). Functional imaging of brain responses to pain. A review and meta-analysis, *Neurophysiologie Clinique/Clinical Neurophysiology*, 30(5), pp. 263-288. Doi:10.1016/s0987-7053(00)00227-6.
- Pubchem struktur quercetin, Diakses dari <https://pubchem.ncbi.nlm.nih.gov/compound/quercetin#section=Top> pada tanggal 10 Februari 2019.
- Raha, S. & Robinson, B. H, 2000, Mitochondria, oxygen free radicals, disease and ageing. *Trends Biochem*, 25, 502–508.
- Rhudy, J.L., Meagher, M.W., 2001. The role of emotion in pain modulation. *Curr. Opin. Psychiatry* 14, 241–245.
- Rhudy, J.L., Williams, A.E., McCabe, K.M., Russell, J.L., Maynard, L.J, 2008, Emotional control of nociceptive reactions (ECON): do affective valence and arousal play a role?, *Pain* 136, 250–261.
- Rice-Evans CA, Miller NJ, Bolwell PG, et al, 1995, *The relative antioxidant activities of plant-derived polyphenolic flavonoids*, *Free Radic Res*, 22(4):375–383.
- Rothwell, J.A., Day, A.J., & Morgan, M.R.A., 2005. Experimental determination of octanol – water partition coefficients of quercetin and related flavonoids. *Journal Agricultural and Food Chemistry*, 53, pp. 4355-4360.
- Ruiz-Medina, J., Baulies, A., Bura, S. A., et al , 2013, Paclitaxel-induced neuropathic pain is age dependent and devolves on glial response, *Eur. J. Pain* 17, 75–85. doi: 10.1002/j.1532-2149.2012. 00172.

- Salat K, Cios A, Wyska E, Salat R et al, 2014, Antiallodynic and antihyperalgesic activity of 3-[4-(3-trifluoromethyl-phenyl)-piperazin-1-yl]-dihydrofluran-2-one compared to pregabalin in chemotherapy-induced neuropathic pain in mice, *Pharmacology, Biochemistry and Behavior*; 122: 173-181.
- Seminowicz, D.A., Mikulis, D.J., Davis, K.D., 2004, Cognitive modulation of pain-related brain responses depends on behavioral strategy, *Pain* 112, 48–58.
- Seminowicz, D.A., Davis, K.D., 2007, A re-examination of pain-cognition interactions: Implications for neuroimaging, *Pain* 130, 8–13.
- Seretny, M. et al., 2014. Incidence, prevalence, and predictors of chemotherapy induced peripheral neuropathy: A systematic review and meta analysis', *Pain*, 155(12), pp. 2461–2470. doi:10.1016/j.pain.2014.09.020.
- Starobova, H. Dan Vetter, I., 2017, Pathophysiology of Chemotherapy-Induced Peripheral Neuropathy, *Frontiers in Molecular Neuroscience*, 10 (174), PP. 1-21.
- Steeds Charlotte E, 2016, The anatomy and physiology of pain, *Anaesthesia and Pain Medicine*, Basic Science, 55-59.
- Tall JM & Raja SN, 2004, Dietary constituents as novel therapies for pain, *Clin J Pain*, 20:19–26.
- Toma, W., Kyte, S.L., Bagdas, D., Alkhlaif, Y., et al, 2017, Effect of paclitaxel on the development of neuropathy and effective behaviors in the mouse, *Neuropharmacology*, 117, PP. 305-315.
- Tracey, I., Mantyh, P.W., 2007, The Cerebral Signature for Pain Perception and Its Modulation, *Neuron* 55, 377–391.
- Turrens, J. F, 2003, Mitochondrial formation of reactive oxygen species, *J. Physiol*, 552, 335–344.
- Tseng MT, Chiang MC, Chao CC, Tseng WY, Hsieh ST. (2013). fMRI evidence of degeneration-induced neuropathic pain in diabetes: enhanced limbic and striatal activations. *Human Brain Mapping*, 34(10), pp. 2733-2746, doi:10.1002/hbm.22105.

- Ushio S, Egashira N, Sada H, Kawashiri T et al, 2012, Goshajinkigan reduces oxaliplatin-induced peripheral neuropathy without affecting anti-tumor efficacy in rodents, *European Journal of Cancer*; 48: 1407-13.
- Vaello Boadas P, S Castany, J Homas, B Alvarez-Perez, M Deulofeu and E Verdu, 2016, Neuroplasticity of ascending and descending pathway after somatosensory system injury: reviewing knowledge to identify neuropathic pain therapeutic targets, *Neuroplasticity in neuropathic pain*, 54, 330-340.
- Wang G, Zhang J, Liu L, Sharma S, Dong Q, 2012, *Quercetin potentiates doxorubicin mediated antitumor effects against liver cancer through*, p53/Bcl-xl. PLoS One 7:e51764.
- Wang G, Song L, Wang H, Xing N, 2013, *Quercetin synergizes with 2-methoxyestradiol inhibiting cell growth and inducing apoptosis in human prostate cancer cells*. Oncol Rep 30:357– 363.
- Windebank AJ, Grisold W, Chemotherapy-induced neuropathy, *J peripher Nerv Syst*, 2008;13(1):27-46.
- World Health Organization, 2012, Guidelines on the pharmacological treatment of persisting pain in children with medical illnesses, *World Health Organization*.
- Wolf Sherry, Debra Barton, Lisa Kottschade, et al, 2008, Chemotherapy-Induced Peripheral Neuropathy:Prevention and Treatment Strategies, *European Jurnal of Cancer*., 44 : 1507-1515.
- Woolf CJ & Mannion RJ, 1999, *Neuropathic pain etiology, symptoms, mechanisms, and management*, Lancet;353:1959–1964. 5.
- Wrigley et. al. (2009). Neuropathic pain and primary somatosensory cortex reorganization following spinal cord injury. *Pain*, 141(1), pp. 52-59. doi:10.1016/j.pain.2008.10.007.
- Xu H, Wu LJ, Wang H, Zhang X, Vadakkan KI, Kim SS, Steenland HW, Zhuo M. (2008). Presynaptic and postsynaptic amplifications of neuropathic pain in the anterior cingulate cortex. *Journal of Neuroscience*, 28(29), pp. 7445-7453. doi:10.1523/jneurosci.1812-08.2008.

Yamashita A, *et al.* (2014). Astrocytic activation in the anterior cingulate cortex is critical for sleep disorder under neuropathic pain. *Synapse*, 68(6), pp. 235-247. doi:10.1002/syn.21733.

Youdim, K.A., Qaiser, M.Z., Begley, D.J., Rice-Evans, C.A., & Abbott N.J., 2004. Flavonoid permeability across an in situ model of the blood-brain barrier. *Free Radical Biology and Medicine*, 36(5), pp. 592–604.

Zhang, H., Yoon, S. Y., and Dougherty, P. M., 2012, *Evidence that spinal astrocytes but not microglia contribute to the pathogenesis of Paclitaxel-induced painful neuropathy Pain*, 293–303. doi: 10.1016/j.jpain.2011.12.002.

Zheng Huaien, Wen Hua Xiao and Gray J. Ben net, 2011, Functional deficits in peripheral nerve mitochondria in rats with paclitaxel- and oxaliplatin-evoked painful peripheral neuropathy, *experimental Neurology* 232, 154-161.

Zhuo, M., 2008, Cortical excitation and chronic pain, *Trends Neurosci*, 31, 199–207.

Zhuo H-H., Zhang, L., Zhuo, Q-G., Fang, Y., dan Ge, W-H., 2015, Borneol attenuates oxaliplatin induced neuropathy hyperalgesia in mice, *NeuroReport*, 27, pp. 160-165.