

## Daftar Pustaka

- Abdanipour A, Tiraihi T, Delshad A, 2011. Trans-differentiation of the adipose tissue-derived stem cells into neuron-like cells expressing neurotrophins by selegiline. *Iranian Biomedical Journal*, vol. 15, no. 4, pp. 113-21.
- Ahmed E, Donovan T, Zhang LYQ, 2015. Mitochondrial targeted antioxidant in cerebral ischemia. *Journal of Neurol & Neuroscience*, vol. 6, no. 2, pp. 1-6.
- Allen KA, Brandon DH, 2011. Hypoxic ischemic encephalopathy: pathophysiology and experimental treatments. *Newborn Infant Nursing Reviews*, vol. 11, pp. 125-33.
- Alok Sharma HS, Nandini G, Prema B, Pooja K, Amruta P, 2014. Stem Cell Therapy for Cerebral Palsy – A Novel Option. In: Emira, ed. *Cerebral Palsy - Challenges for the Future*, Svraka, pp. 217-42.
- Amantea D, Micieli G, Tassorelli C, Cuartero MI, Ballesteros I, Certo M, Moro MA, Lizasoain I, Bagetta G, 2015. Rational modulation of the innate immune system for neuroprotection in ischemic stroke. *Frontiers In Neuroscience*, vol. 9, pp. 1-20.
- Amirkhani MA, Zare S, Hassani AS, Torbati E, Nilforoushzadeh MA, 2017. Literature review of adipose-derived mesenchymal cells from history to approaches. *Iran Red Crescent Medical Journal*, vol. 19, no.1, pp. 1-5.
- Anghileri E, Marconi S, Pignatelli A, Cifelli P, Galié M, Sbarbati A, Krampera M, Belluzzi O, Bonetti B, 2008. Neuronal differentiation potential of human adipose-derived mesenchymal stem cells. *Stem Cells And Development*, vol. 17, no. 5, pp. 909-16.
- Ashjian PH, Elbarbary AS, Edmonds B, DeUgarte D, Zhu M, Zuk PA, Lorenz HP, Benhaim P, Hedrick MH, 2003. In vitro differentiation of human processed lipoaspirate cells into early neural progenitors. *Plastic and Reconstructive Surgery*, vol. 111, no. 6, pp. 1922–31.
- Baglioni S, Francalanci M, Squecco R, Lombardi A, Cantini G, Angeli R, Gelmini S, Guasti D, Benvenuti S, Annunziato F, Bani D, Liotta F, Francini F, Perigli G, Serio M, Luconi M, 2009. Characterization of human adult stem-cell populations isolated from visceral and subcutaneous adipose tissue. *The FASEB Journal*, vol. 23, no. 10, pp. 3494-505.
- Banerjee S, Williamson DA, Habib N, Chataway, 2012. The potential benefit of stem cell therapy after stroke: an update. *Vascular Health & Risk Management*, vol. 8, pp. 569-80.

- Benn SC, Woolf CJ, 2004. Adult neuron survival strategies -- slamming on the brakes. *Nature Reviews Neuroscience*, vol. 5, no. 9, pp. 686-700.
- Berti R, Williams AJ, Moffett JR, Hale SL, Velarde LC, Elliott PJ, Yao C, Dave JR, Tortella FC, 2002. Quantitative real-time RT-PCR analysis of inflammatory gene expression associated with ischemiareperfusion brain injury. *Journal Cerebral Blood Flow Metabolism*, vol. 22, pp. 1068-79.
- Bonfield TL, Caplan AI, 2010. Adult mesenchymal stem cells: an innovative therapeutic for lung diseases. *Discovery Medicine*, vol. 9, no. 47, pp. 337-45.
- Bradshaw RA, Rust JM, Ibanez CF, McDonald NQ, Lapatto R, Blundell TL, 1994. Nerve growth factor: structure/function relationships. *Protein Sciencedirec*, vol. 3, pp. 1901-13.
- Broughton BRS, Reutens DC, Sobey CG, 2009. Apoptotic mechanism after cerebral ischemia, *Stroke*, vol. 40, pp. e331-9.
- Cardozo AJ, Gómez DE, Argibay PF, 2012. Neurogenic differentiation of human adipose-derived stem cells: relevance of different signaling molecules, transcription factors, and key marker genes, *Gene*, vol. 511, no. 2, pp. 427-36.
- Carroll JE, Mays RW, 2011. Update on stem cell therapy for cerebral palsy. *Expert Opinion Biological Therapy*, vol. 11, pp. 463-71.
- Chen G, Wang Y, Xu Z, Fang F, Xu R, Wang Y, Hu X, Fan L, Liu H, 2013. Neural stem cell-like cells derived from autologous bone mesenchymal stem cells for the treatment of patients with cerebral palsy. *Journal of Translational Medicine*, vol. 11, no. 21, pp 1-11.
- Chen Q, Long Y, Yuan X, Zou L, Sun J, Chen S, Perez-Polo JR, Yang K, 2005. Protective effects of bone marrow stromal cell transplantation in injured rodent brain: synthesis of neurotrophic factors. *Journal of Neuroscience Research*, vol. 80, pp. 611-9.
- Chen X, Katakowski M, Li Y, Lu D, Wang L, Zhang L, Chen J, Xu Y, Gautam S, Mahmood A, Chopp M, 2002. Human bone marrow stromal cell cultures conditioned by traumatic brain tissue extracts: growth factor production. *Journal of Neuroscience Research*, vol. 69, pp. 687-91.
- Cheng Z, Zhu W, Cao K, Wu F, Li J, Wang G, Li H, Lu M, Ren Y, He X, 2016. Anti-inflammatory mechanism of neural stem cell transplantation in spinal cord injury. *International Journal of Molecular Sciences*, vol. 17, no. 1380, pp. 1-15.
- Chiaretti A, Antonelli A, Genovese O, Fernandez E, Giuda D, Mariotti P, Riccardi R, 2008. Intraventricular nerve growth factor infusion improves cerebral blood flow and

- stimulates doublecortin expression in two infants with hypoxic-ischemic brain injury. *Neurological Research*, vol. 30, pp. 223–228.
- Chiaretti A, Barone G, Riccardi R, Antonelli A, Pezzotti P, Genovese O, Tortorolo L, Conti G, 2009. NGF, DCX, and NSE upregulation correlates with severity and outcome of head trauma in children, *Neurology*, vol. 72, no. 7, pp. 609-16.
- Chiaretti A, Genovese O, Riccardi R, Di Rocco C, Di Giuda D, Mariotti P, Pulitano S, Piastra M, Polidori G, Colafati GS, Aloe L, 2005. Intraventricular nerve growth factor infusion: a possible treatment for neurological deficits following hypoxic-ischemic brain injury in infants. *Neurological Research*, vol. 27, pp. 741–746.
- Daadi MM, Davis AS, Arac A, Li Z, Maag AL, Bhatnagar R, Jiang K, Sun G, Wu JC, Steinberg GK, 2010. Human neural stem cell grafts modify microglial response and enhance axonal sprouting in neonatal hypoxic-ischemic brain injury. *Stroke*, vol. 41, pp. 516-523.
- Dammanahalli JK, Wang X, Sun Z, 2011. Genetic interleukin-10 deficiency causes vascular remodeling via the upregulation of Nox1. *Journal of Hypertension*, vol. 29, no. 11, pp. 2116–25.
- Dawson VL, Dawson TM, Bartley DA, Uhl GR, Snyder SH, 1993. Mechanisms of Nitric Oxide-mediated Neurotoxicity in Primary Brain Cultures. *The Journal of Neuroscience*, vol. 13, no. 6, pp. 2651-61.
- de Bilbao F, Arsenijevic D, Moll T, Garcia-Gabay I, Vallet P, Langhans W, Giannakopoulos, 2009. In vivo over-expression of interleukin-10 increases resistance to focal brain ischemia in mice. *Journal of Neurochemistry*, vol. 110, no. 1, pp. 12–22.
- de Cerio F, Lara-Celador I, Alvarez A, Hilario E, 2013. Neuroprotective Therapies after Perinatal Hypoxic-Ischemic Brain Injury. *Brain Science*, vol. 3, pp. 191-214.
- Deshpandea LS, Sund DA, Sombati S, Baranova A, Wilson MS, Attkisson E, Hamm RJ, DeLorenzo RJ, 2008. Alterations in neuronal calcium levels are associated with cognitive deficits after traumatic brain injury. *Neuroscience Letters*, vol. 441, no.1, pp. 115-9.
- Dhar S, Yoon ES, Kachgal S, Evans GR, 2007. Long-term maintenance of neuronally differentiated human adipose tissue-derived stem cells. *Tissue Engineering*, vol. 13, no. 11, pp. 2625-32.
- Didion SP, Kinzenbaw DA, Schrader LI, Chu Y, Faraci FM, 2009. Endogenous interleukin-10 inhibits angiotensin II-induced vascular dysfunction. *Hypertension*, vol. 54, no. 3, pp. 619–24.

- Dixon CE, Flinn P, Bao J, Venya R, Hayes RI, 1997. Nerve growth factor attenuates cholinergic deficits following traumatic brain injury in rats. *Experimental Neurology*, vol. 146, pp. 479-90.
- Donega V, Velthoven CTJ, Nijboer CH, Bel VF, Kas MJH, Kavelaars A, Heijnen CJ, 2013. Intranasal mesenchymal stem cell treatment for neonatal brain damage: long-term cognitive and sensorimotor improvement. *Plos One*, vol. 8, no. 1, pp.1-7.
- Douglas RM, Ryu J, Kanaan A, Rivero MdC, Dugan LL, Haddad GG, Ali SS, 2010. Neuronal death during combined intermittent hypoxia/hypercapnia is due to mitochondrial dysfunction. *American Journal of Physiol Cell Physiology*, vol. 298, no. 6, pp. 1594-602.
- Eghbalian F, 2010. Frequency of Hypoxic-Ischemic Encephalopathy among Hospitalized Neonates in West Iran. *Iranian Journal Pediatrics*, vol. 20, no. 2, pp. 244-5.
- Espósito, E, Cuzzocrea, S, 2011. Anti-TNF therapy in the injured spinal cord, *Trends in Pharmacological Sciences*, vol.. 32, pp 107–15.
- Ferrarese C, Mascalucci P, Zoia C, Cavarretta R, Frigo M, Begni B, Sarinella F, Frattola L, De Simoni M, 1999. Increased cytokine release from peripheral blood cells after acute stroke. *Journal of Cerebral Blood Flow & Metabolism*, vol. 19, pp. 1004–9.
- Fluri F, Schuhmann MK, Kleinschmitz C, 2015. Animal models of ischemic stroke and their application in clinical research. *Drug Design, Development and Therapy*, vol. 9, pp. 3445-54.
- Folgiero V, Migliano E, Tedesco M, Lacovelli S, Bon G, Torre ML, Sacchi A, Marazzi M, Bucher S, Falcioni R, 2010. Purification and characterization of adipose-derived stem cells from patients with lipoaspirate transplant. *Cell Transplantation*, vol. 19, no. 10, pp. 1225–35.
- Fujimura J, Ogawa R, Mizuno H, Fukunaga Y, Suzuki H, 2005. Neural differentiation of adipose-derived stem cells isolated from GFP transgenic mice. *Biochemical and Biophysical Research Communications* vol. 333, no. 1, pp. 116-21.
- Garcia JH, Liu KF, Ho KL, 1995. Neuronal necrosis after middle cerebral artery occlusion in Wistar rats progresses at different time intervals in the caudoputamen and the cortex. *Stroke*, vol. 26, no. 4, pp. 636–43.
- Gincberg G, Shohami E, Trembovler V, Alexandrovich AG, Lazarovici P, Elchalal U, 2018. Nerve growth factor plays a role in the neurotherapeutic effect of a CD45 pan-hematopoietic subpopulation derived from human umbilical cord in a traumatic brain injury model, *Cytotherapy*, vol. 20, no. 2, pp. 245-61.

- Gingras M, Champigny MF, Berthod F, 2007. Differentiation of human adult skin-derived neuronal precursors into mature neurons. *Journal of Cellular Physiology*, vol. 210, no. 2, pp. 498-506.
- Guo MF, Yu JZ, Ma CG, 2011. Mechanisms related to neuron injury and death in cerebral hypoxic ischaemia. *Folia Neuropathologica*, vol. 49, no. 2, pp.78-87.
- Goss JR, O'Malley ME, Zou L, Styren SD, Kochanek PM, DeKosky ST, 1998. Astrocytes are the major source of nerve growth factor upregulation following traumatic brain injury in the rat. *Experimental Neurology*, vol. 149, pp. 301-9.
- Grilli M, Barbieri I, Basudev H, Brusa R, Casati C, Lozza G, Ongini E, 2000. Interleukin-10 modulates neuronal threshold of vulnerability to ischaemic damage. *European Journal of Neuroscience*, vol. 12, no. 7, pp. 2265-72.
- Haddad M, Rhinn H, Bloquel C, Coqueran B, Szabo C, Plotkine M, Scherman D, Margaill I, 2006. Antiinflammatory effects of PJ34, a poly(ADP-ribose) polymerase inhibitor, in transient focal cerebral ischemia in mice. *British Journal Pharmacology*, vol.149, pp. 23-30.
- Handel Mv, Swaab H, Vries LSD, Jongmans MJ, 2007. Long-term cognitive and behavioral consequences of neonatal encephalopathy following perinatal asphyxia: a review. *Europen Journal of Pediatrics*, vol. 166, pp. 645-54.
- Hansen AR, Soul JS, 2012. Perinatal asphyxia and hypoxic ischemic encephalopathy. In: Cloherty JP, Eichenwald EC, Hansen AR, Stark AR, editor. Manual of Neonatal Care 7. Philadelphia, Lippincott Williams and Witkins, pp. 721-6.
- Haque A, Polcyn R, Matzelle D, Bnik NL, 2018. New insights into the role of Neuron-Specific Enolase in neuro-Inflammation, neurodegeneration and neuroprotection. *Brain Science*, vol 8, no 2, pp 33.
- Harukuni I, Bhardwaj A, 2006. Mechanisms of Brain Injury after Global Cerebral Ischemia. *Neurology Clinics*, vol. 24, pp. 1-21.
- Hatfield R, McKernan R, 1992. CSF neuro-spesific enolase as a quantitative marker of neuronal damage in rat stroke, *Brain Research*, vol. 577, no. 1, pp. 249-52.
- Hossmann K-A, 2012. The two pathophysiologies of focal brain ischemia: implications for translational stroke research. *Journal of Cerebral Blood Flow & Metabolism*, vol. 32, no.7, pp. 1310–6.
- Howells DW, Porritt MJ, Rewell SS,O'Collins V, Sena ES, Worp HB, Traystman RJ, Macleod MR, 2010. Different strokes for different folks: the rich diversity of animal models of focal cerebral ischemia. *Journal of Cereb Blood Flow & Metabolism*, vol. 30, no. 8, pp. 1412–31.

- Huet O, Laemmel E, Fu Y, Dupic L, Aprico A, Andrews KL, Moore SL, Harrois A, Meikle PL, Vicaut E, Chin-Dusting JP, Duranteau J, 2013. Interleukin 10 antioxidant effect decreases leukocytes/endothelial interaction induced by tumor necrosis factor alpha. *Shock*, vol. 39, no. 1, pp. 83–8.
- Ikeda T, Xia XY, Xia YX, Ikenoune T, Han B, Choi BH, 2000. Glial cell-line-derived neurotrophic factor protects against ischemia/hypoxia-induced brain injury in neonatal rat. *Acta Neuropathologica*, vol. 100, pp. 161-7.
- Inacio A, Ruscher K, Leng L, Bucala R, Deierborg T, 2011. Macrophage migration inhibitory factor promotes cell death and aggravates neurologic deficits after experimental stroke. *Journal of Cerebral Blood Flow & Metabolism*, vol. 31, pp. 1093-106.
- Jellema RK, Wolfs TG, Lima Passos V, Zwanenburg A, Ophelders DR, Kuypers E, Hopman AHN, Dudink J, Steinbusch HW, Andriessen P, Germeraad WTV, Vanderlocht J, Kramer BW, 2013. Mesenchymal stem cells induce T-cell tolerance and protect the preterm brain after global hypoxia-ischemia, *PLoS One*, vol. 8, e73031.
- Kamel N, Tanaka N, Oishi Y, Hamasaki T, Nakanishi K, 2007. BDNF, NT-3, and NGF released from transplanted neural progenitor cells promote corticospinal axon growth in organotypic cocultures. *Spine cord*, vol. 32, no.12, pp.1272-8.
- Kang H, Yang PY, Rui YC, 2008. Adenovirus viral interleukin-10 inhibits adhesion molecule expressions induced by hypoxia/reoxygenation in cerebrovascular endothelial cells. *Acta Pharmacologica Sinica*, vol. 29, no. 1, pp. 50–6.
- Kang SK, Shin MJ, Jung JS, Kim YG, Kim CH, 2006. Autologous adipose tissue-derived stromal cells for treatment of spinal cord injury. *Stem Cells And Development*, vol. 15, no. 4, pp.583-94.
- Kinzenbaw DA, Chu Y, Pena Silva RA, Didion SP, Faraci FM, 2013. Interleukin-10 protects against aging-induced endothelial dysfunction. *Physiological Reports*, vol. 1, no. 6, pp. e00149.
- Kittaka M, Giannotta SL, Zelman V, Correale JD, DeGiorgio CM, Weiss MH, Zlokovic BV, 1997. Attenuation of brain injury and reduction of neuron-specific enolase by nicardipine in systemic circulation following focal ischemia and reperfusion in a rat model. *Journal of Neurosurgery*, vol. 87, pp 731-7.
- Kokaia Z, Martino G, Schwartz M, Lindvall O, 2012. Crosstalk between neural stem cell and immune cells: the key to better brain repair?, *Nature Neuroscience*, vol. 15, no. 8, pp. 1078-87.

- Lafon-Cazal M, Bougault I, Steinberg R, Pin JP, Bockaert J, 1992. Measurement of gamma-enolase release, a new method for selective quantification of neurotoxicity independently from glial lysis. *Brain Res*, vol. 593, pp 63-8.
- Lee, JH, Sung YB, Jang SH, 2016. Nerve growth factor expression in stroke induced rats after shock wave. *Journal of Physical Therapy Science*, vol. 28, pp. 3451-3.
- Li HL, Kostulas N, Huang YM, Xiao BG, van der Meide P, Kostulas V, Giedraitas V, Link H, 2001. Il-17 and ifn-gamma mrna expression is increased in the brain and systemically after permanent middle cerebral artery occlusion in the rat. *Journal Neuroimmunology*, vol. 116, pp. 5-14.
- Li J, Zhu H, Liu Y, Li Q, Lu S, Feng M, Huang L, Ma C, An Y, Zhao RC, Wang R, Qin, C, 2010. Human mesenchymal stem cell transplantation protects against cerebral ischemic injury and upregulates interleukin-10 expression in Macaca fascicularis, *Brain Research*, vol. 1334, pp. 65–72.
- Li Y, Chen J, Chen XG, Wang L, Gautam SC, Xu YX, Katakowski M, Zhang LJ, Lu M, Janakiraman N, Chopp M, 2002. Human marrow stromal cell therapy for stroke in rat, neurotrophins and functional recovery, *Neurology*, vol. 59, pp. 514-23.
- Li Yan, Dejin Zheng, Ren-He Xu, 2018 Critical Role of Tumor Necrosis Factor Signaling in Mesenchymal Stem Cell-Based Therapy for Autoimmune and inflammatory Diseases, *Frontiers in Immunology*. Vol. 9, article 1658.
- Li W, Ren G, Huang Y, Su J, Han Y, Li J, Chen X, Cao K, Chen Q, Shou P, Zhang L, Yuan ZR, Roberts AI, Shi S, Le AD, Shi Y, 2012. Mesenchymal stem cells: A double-edged sword in regulating immune responses. *Cell Death & Differentiation*, vol. 19, pp. 1505-13.
- Li X., Huang M, Zhao R, Zhao C, Liu Y, Zou H, Chen L, Guan Y, Zhang YA, 2018. Intravenously Delivered Allogeneic Mesenchymal Stem Cells Bidirectionally Regulate Inflammation and Induce Neurotrophic Effects in Distal Middle Cerebral Artery Occlusion Rats Within the First 7 Days After Stroke, *Cellular Physiology Biochemistry*, vol. 46, pp. 1951–70.
- Liang QJ, Jiang M, Wang XH, Le LL, Xiang M, Sun N, Meng D, Chen SF, 2015. Pre-existing interleukin 10 in cerebral arteries attenuates subsequent brain injury caused by ischemia/reperfusion. *IUBMB Life*, vol. 67, no. 9, pp. 710–9.
- Llado J, Haenggeli C, Maragakis NJ, Snyder EY, Rothstein JD, 2004. Neural stem cells protect against glutamate-induced excitotoxicity and promote survival of injured motor neurons through the secretion ofneurotrophic factors, *Molecular And Cellular Neuroscience*, vol. 27, no. 3, pp. 322-31.

- Mahmood A, Lu D, Chopp M, 2004. Intravenous administration of marrow stromal cells (MSCs) increases the expression of growth factors in rat brain after traumatic brain injury, *Journal of neurotrauma*, vol. 21, no. 1, pp. 33-9.
- Malendez MC, Yawno T, Jenkin G, Miller SL, 2013. Stem cell therapy to protect and repair the developing brain: a review of mechanisms of action of cord blood and amnion epithelial derived cells, *Front in Neuroscience*, vol. 7, pp. 1-14.
- Meirelles LDS, Simon D, Regner A, 2017. Neurotrauma: the crosstalk between neurotrophins and inflammation in the acutely injured brain, *International Journal of Molecular Sciences*, vol. 18, pp.1-22.
- Meloni BP, 2017. Pathophysiology and neuroprotective strategies in hypoxic-ischemic brain injury and stroke, *Brain science*, vol 7, no 110, pp 1-4.
- Meistrell MEI, Botchkina GI, Wang H, Di Santo E, Cockcroft KM, Vishnubhakat JM, Ghezzi P, Traces KJ, 1997. Tumor necrosis factor is a brain damaging cytokine in cerebral ischemia, *Shock*, vol. 8, pp. 341– 8.
- Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A, 2001. Interleukin-10 and the interleukin-10 receptor, *Annual Review of Immunology*, vol. 19, pp. 683–765.
- Moore TJ, Abrahamse H, 2014. Neuronal differentiation of adipose derived stem cells: progress so far, *International Journal of Image Photo*, vol. 2014, pp. 1-8.
- Morioka C, Komaki M, Taki A, Honda I, Yokoyama N, Iwasaki K, Iseki S, Morio T, Morita I, 2017. Neuroprotective effects of human umbilical cord-derived mesenchymal stem cells on periventricular leukomalacia-like brain injury in neonatal rats, *Inflammation & Regulation*, vol. 37, no. 1, pp.1-10.
- Morita Y, Takizawa S, Kamiguchi H, Uesugi T, Kawada H, Takagi S, 2007. Administration of hematopoietic cytokines increases the expression of anti-inflammatory cytokine (IL-10) mRNA in the subacute phase after stroke, *Neuroscience Research*, vol. 58, no.4, pp. 356-60.
- Nakajima M, Nito C, Sowa K, Suda S, Nishiyama Y, Takahasi A, Kasahara YN, Imagawa K, Hirato T, Ueda M, Kimura K, Okada T, 2017. Mesenchymal stem cells overexpressing Inteleuki-10 promote neuroprotection in experimental acute ischemic stroke, *Molecular Therapy Methods & Clinical Development*, vol. 15. No. 6, pp. 102-11.
- Nomoto T, Okada T, Shimazaki K, Yoshioka T, Nonaka-Sarukawa M, Ito T, Takeuchi K, Katsura KI, Mizukami H, Kume A, Ookawara S, Ikeda U, Katayama Y, Ozawa K, 2009. Systemic delivery of IL-10 by an AAV vector prevents vascular remodeling and end-

- organ damage in stroke-prone spontaneously hypertensive rat, *Gene Therapy*, vol. 16, no. 3, pp. 383–91.
- Northington FJ, 2006. Brief update on animal models of hypoxic-ischemic encephalopathy and neonatal stroke, *International League of Associations Rheumatology*, vol. 47, no. 1, pp. 32-8.
- Offner H, Subramanian S, Parker SM, Afentoulis ME, Vanderbark AA, Hurn PD, 2006. Experimental stroke induces massive, rapid activation of the peripheral immune system. *Journal of Cerebral Blood Flow & Metabolism*, vol. 26, pp. 654–65.
- Orellana JA, Froger N, Ezan P, Jiang JX, Bennett MVL, Naus CC, Giaume C, Saez JC, 2011. ATP and glutamate released via astroglial connexin43 hemichannels mediate neuronal death through activation of pannexin 1 hemichannels, *Journal Neurochemistry*, vol. 118, no. 5, pp. 826-40.
- Palmio J, Peltola J, Vuorinen P, Laine S, Suhonen J, Keränen T, 2001. Normal CSF neuron-specific enolase and S-100 protein levels in patients with recent non-complicated tonic-clonic seizures, *Journal of the Neurological Sciences*, vol .183, pp. 27-31.
- Palsdottir K, Dagbjartsson A, Thorkelsson T, Hardardottir H, 2007. Birth asphyxia and hypoxic ischemic encephalopathy, incidence and obstetric risk factors, *Laeknabladid Journal*, vol. 93, no. 9, pp. 595–601.
- Pan W, Kastin AJ, 2007. Tumor necrosis factor and stroke: role of the blood–brain barrier, *Progress in Neurobiology*, vol. 83, no. 6, pp. 363-74.
- Pang Y, Cai Z, Rhodes PG, 2003. Disturbance of oligodendrocyte development, hypomielination and white matter injury in the neonatal rat brain after intracerebral injeksiion of lypopolysaccharide, *Developmental Brain Research*, vol. 140, pp. 205-14.
- Puig IP, Miro F, Salas-Perdomo A, Bonfill-Teixidor E, Ferrer-Ferrer M, Marquez-Kisinousky L, Planas AM, 2013. IL-10 deficiency exacerbates the brain inflammatory response to permanent ischemia without preventing resolution of the lesion, *Journal of Cerebral Blood Flow & Metabolism*, vol.33, no.12, pp. 1955–66.
- Pineau, I., Lacroix, S, 2007. Proinflammatory cytokine synthesis in the injured mouse spinal cord: Multiphasic expression pattern and identification of the cell types involved, *Journal of Comparative Neurology*, vol. 500, pp. 267–85.
- Rivkin MJ, Bernard TJ, Dowling MM, Lefond CA, 2017. Guidelines for urgent management of stroke in children, *Pediatric Neurology*, vol. 56, pp. 8-17.
- Romanov YA, Darevskaya AN, Merzlikina NV, Buravkova LB, 2005. Mesenchymal stem cells from human bone marrow and adipose tissue: isolation, characterization, and

- differentiation potentialities. *Bulletin of Experimental Biology and Medicine*, vol. 140, no. 1, pp. 138-43.
- Ruff CA, Faulkner SD, Fehlings MG, 2013. The potential for stem cell therapies to have an impact on cerebral palsy: opportunities and limitations, *Developmental Medicine & Child Neurology*, vol. 55, pp. 689-97.
- Sankar R, Shin DH, Wasterlain CG, 1997. Serum specific enolase is marker for neuronal damage following status epilepticus in the rat. *Epilepsy Foundation*, vol. 28, pp. 129-36.
- Shankaran S, 2012. Hypoxic-ischemic encephalopathy and novel strategies for neuroprotection, *Clinical Perinatology*, vol. 39, pp. 919-29.
- Scheibe F, Ladhoff J, Huck J, Grohmann M, Blazej K, Oersal A, Baeva N, Seifert M, Priller J, 2012. Immune effects of mesenchymal stromal cells in experimental stroke, *Journal of cerebral blood flow & metabolism*, vol 32, pp. 1578-88.
- Shen J, Zhang H, Lin H, Su H, Xing D, Du L, 2007. Brazilein protects the brain against focal cerebral ischemia reperfusion injury correlating to inflammatory response suppression, *European Journal Pharmacology*, vol.558, pp. 88–95.
- Spera PA, Ellison JA, Feuerstein GZ, Barone FC, 1998. IL-10 reduces rat brain injury following focal stroke, *Neuroscience Letters*, vol. 251, no. 3, pp. 189–92.
- Stanimirovic D, Satoh K, 2000. Role of Inflammation Following Stroke and Neurotrauma Inflammatory Mediators of Cerebral Endothelium: A Role in Ischemic Brain Inflammation, *Brain Pathology*, vol. 10, pp.113-26
- Stirling D, Stys P, 2010. Mechanisms of axonal injury: internodal nanocomplexes and calcium deregulation, *Trends Molecular Medicine*, vol.16, no. 4, pp. 160-70.
- Stoll BJ, Kliegman RM. Hypoxia-ischemia, 2007. Dalam : Behrman RE, Kliegman RM, Jenson HB, editor. Nelson Textbook of Pediatrics. Edisi 18, Philadelphia, Saunders, pp. 566–8.
- Strem BM, Hicok KC, Zhu M, Wulur I, Alfenso Z, Schreiber RE, Fraser JK, Hedrick MH, 2005. Multipotential differentiation of adipose tissue-derived stem cells, *The Keio Journal of Medicine*, vol. 54, no. 3, pp. 132–41.
- Strle K, Zhou JH, Shen WH, Broussard SR, Johnson RW, Freund GG, Dantzer R, Kelley KW, 2001. Interleukin-10 in the brain, *Critical Reviews Immunology*, vol.21, no.5, pp. 427–49.
- Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N, 2010. A meta-analysis of cytokines in Alzheimer's disease. *Biol Psychiatry*, vol. 68, no. 10, pp. 930–41.

- Takata M, Nakagomi T, Kashiwamura S, Nakano DA, Saino O, Nakagomi N Okamura H, Minura O, Taguchi A, Matsuyama T, 2012. Glucocorticoid-induced TNF receptor-triggered T cells are key modulators for survival/death of neural stem/progenitor cells induced by ischemic stroke, *Cell Death & Differentiation*, vol. 19, no. 5, pp. 756-67.
- Tholpady SS, Katz AJ, Ogle RC, 2003. Mesenchymal stem cells from rat visceral fat exhibit multipotential differentiation in vitro, *Anatomical Record A: Discoveries in Molecular, Cellular, and Evolutionary Biology*, vol. 272, no. 1, pp. 398–402.
- Toma JG, McKenzie IA, Bagli D, Miller FD, 2005. Isolation and characterization of multipotent skin-derived precursors from human skin, *Stem Cells*, vol. 23, no. 6, pp. 727-37.
- Torio-Padron N, Huotari AM, Eisenhardt SU, Borges J, Stark GB, 2010. Comparison of pre-adipocyte yield, growth and differentiation characteristics from excised versus aspirated adipose tissue, *Cells Tissues Organs*, vol. 191, no. 5, pp. 365-71.
- Tsuchiya D, Hong S, Kayama T, Panter SS, Weinstein PR, 2003. Effect of suture size and carotid clip application upon blood flow and infarct volume after permanent and temporary middle cerebral artery occlusion in mice. *Brain Research*, vol. 970, no. 1–2, pp. 131–139.
- Van Velthoven CTJ, Kavelaars A, Bel FV, Heijnen CJ, 2010. Repeated mesenchymal stem cell treatment after neonatal hypoxia-ischemia has distinct effects on formation and maturation of new neurons and oligodendrocytes leading to restoration of damage, corticospinal motor tract activity, and sensorimotor function, *The Journal of Neuroscience*, vol. 30, no. 28, pp. 9603-11.
- Van Velthoven CTJ, Kavelaars A, Heijnen CJ, 2012. Mesenchymal stem cells as a treatment for neonatal ischemic brain damage, *Pediatric Research*, vol. 71, no. 4, pp. 474-81.
- Vannucci RC, Vanucci SJ, 2005. Perinatal hypoxic-ischemic brain damage: evolution of an animal model, *Developmental Neuroscience*, vol. 27, pp. 81-6.
- Venkataramana NK, Kumar SK, Balaraju S, Radhakrishnan RC, Bansal A, Dixit A, Rao DK, Das M, Jan M, Gupta PK, Totev SM, 2009. Open-labeled study of unilateral autologous bone-marrow-derived mesenchymal stem cell transplantation in Parkinson's disease, *Translational Research*, vol. 155, no. 2, pp. 62-70.
- Vlachantoni D, Bramall AN, Murphy MP, Taylor RW, Shu X, Tulloch B, Van Veen T, Turnbull DM, McInnes RR, Wright AF, 2011. Evidence of severe mitochondrial oxidative stress and a protective effect of low oxygen in mouse models of inherited photoreceptor degeneration, *Human Molecular Genetics*, vol. 20, pp. 322-35.

- von Heimburg D, Hemmrich K, Haydarlioglu S, Staiger H, Pallua N, 2004. Comparison of viable cell yield from excised versus aspirated adipose tissue, *Cells Tissues Organs*, vol. 178, no. 2, pp. 87-92.
- Wachtel EV, Hendricks-Munoz KD, 2011. Current management of the infant who presents with neonatal encephalopathy, *Current Problems in Pediatric and Adolescent Health Care*, vol. 41, pp. 132–53.
- Wen, TC, Rogido M, Peng H, Genetta T, Moore J, Sola A, 2006. Gender differences in long-term beneficial effects of erythropoietin given after neonatal stroke in postnatal day-7 rats. *Neuroscience*, vol. 139, no. 3, pp. 803-11.
- Williams AJ, Berti R, Dave JR, Elliot PJ, Adams J, Tortella FC, 2004. Delayed treatment of ischemia/ reperfusion brain injury: extended therapeutic window with the proteosome inhibitor MLN519, *Stroke*, vol. 35, pp. 1186–91.
- Wilson, 2015. Animal models of cerebral palsy: hypoxic brain injury in the newborn, *Iranian Journal of Child Neurology*, vol. 9, no. 2, pp. 9-16.
- Won SJ, Kim DY, Gwag BJ, 2002. Cellular and Molecular Pathways of Ischemic Neuronal Death. *Journal of Biochemistry and Molecular Biology*, vol. 35, pp. 67-86.
- Xu W, Zheng J, Gao L, Li T, Zhang J, Shao A, 2017. Neuroprotective effects of stem cells in ischemic stroke, *Stem Cells International*, article ID 4653936, pp. 1-7.
- Yamori Y, Horie R, Handa H, Sato M, Fukase M, 1976. Pathogenetic similarity of strokes in stroke-prone spontaneously hypertensive rats and humans. *Stroke*, vol. 7, no.1, pp. 46–53.
- Yardimoglu M, Ilbay G, Dalcik C, Dalcik H, Sahin D, Ates N, 2008. Immunocytochemistry of neuron specific enolase (NSE) in the rat brain after single and repeated epileptic seizures, *International Journal Neuroscience*, vol. 118, pp. 981-93.
- Yi BRY, Kim SU, Choi KC, 2013. Development and application of neural stem cells for treating various human neurological diseases in animal models, *Laboratory Animal Research*, vol. 29, no. 3, pp. 131-7.
- Ying C, Hu W, Cheng B, Zheng X, Li S, 2012 Neural Differentiation of Rat Adipose-Derived Stem Cells in Vitro, *Cellular and Molecular Neurobiology*, vol. 32, no. 8, pp. 1255–63.
- Zaleska MM, Mercado MLT, Chavez J, Fuerstein GZ, Pangalos MN, Wood A, 2009. The development of stroke therapeutics: promising mechanism and translational challenges, *Neuropharmacology*, vol. 56, pp. 329-41.

- Zandbergen EG, De Haan RJ, Hijdra A, 2001. Systematic review of prediction of poor outcome in anoxic-ischaemic coma with biochemical markers of brain damage. *Intensive Care Medicine*, vol. 27, pp. 1661-7.
- Zandt B-J, Haken Bt, Dijk JGv, Putten MJAMv, 2011. Neural dynamics during anoxia and the "wave of death". *PLoS One*, vol 6, no.22, pp.127.
- Zaremba J, Losy J, 2001. Early TNF-alpha levels correlate with ischaemic stroke severity, *Acta Neurologica Scandinavica*, vol. 104, pp. 288–95.
- Zavan B, Michelotto L, Lancerotto L, Della Puppa A, D'Avella D, Abatangelo G, Vindigni V, Cortivo R, 2010. Neural potential of a stem cell population in the adipose and cutaneous tissues, *Neurological Research*, vol. 32, no. 1, pp. 47-54.
- Zetterberg H, Smith DH, Blennow K, 2013. Biomarkers of mild traumatic brain injury in cerebrospinal fluid and blood, *Nature Review Neurology*, vol. 9, pp. 201-10.
- Zhang R, Liu Y, Yan K, Chen L, Chen XR, li P, Chen F, Jiang X, 2013. Anti-inflammatory and immunomodulatory mechanism of mesenchymal stem cell transplantation in experimental traumatic brain injury, *Journal Neuroinflammation*, vol 10, pp. 106.
- Zhang Y, Liu N, Tang Y, Yang E, Dong S, Huang M, Pan C, Zhang Y, Zhang P, Chen H, Tang Z, 2014. Efficient generation of neural stem cell-like cells from rat adipose derived stem cell after lentiviral transduction with green fluorescent protein, *Molecular Neurobiology*, vol. 50, no. 2, pp. 647-54.
- Zhu Y, Guan YM, Huang HL, Song Q, 2014. Human umbilical cord blood mesenchymal stem cell transplantation suppresses inflammatory responses and neuronal apoptosis during early stage of focal cerebral ischemia in rabbits, *Acta Pharmaceutica Sinice*, vol. 35, pp. 585-91.
- Zuk PA, Zhu M, Ashjian P, De Ugarte DA, Huang JI, Mizuno H, Alfonso ZC, Fraser JK, Benhaim P, Hedrick MH, 2002. Human adipose tissue is a source of multipotent stem cells, *Molecular biology of the cell*, vol. 13, pp. 4279-95.