



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development
Scopus coverage years: from 2010 to 2017 Publisher:
R.K. Sharma, Institute of Medico-Legal Publications
ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:
Public Health, Environmental and Occupational Health
CiteScore 2015- 0.02
SJR 2015-0.105
SNIP 2015-0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof Vidya Surwade

Associate Professor, Dr Baba Saheb Ambedkar, Medical College & Hospital, Rohinee, Delhi

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan B. Md Jagar Din**, (*Associate Professor*)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar** (*Consulting Physician*)
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum**,
Botany and Microbiology Deptt, College of Science, King Saud University,
Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar** (*Associate Professor*)
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy** (*Senior Lecturer*)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyasa** (*Clinical Researcher*)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel** (*Community Medicine*)
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade** (*Associate Professor*)
Navodaya Medical College, Raichur, Karnataka
2. **Dr. E. Venkata Rao** (*Associate Professor*) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
3. **Dr. Amit K. Singh** (*Associate Professor*) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R G Viveki** (*Professor & Head*) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Dr. Santosh Kumar Mulage** (*Assistant Professor*)
Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy** (*Associate Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Dr. Ritu Goyal** (*Associate Professor*)
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar** (*Associate Professor*)
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Dr. Md. Amirul Hassan** (*Associate Professor*)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish** (*Associate Professor*) Microbiology, VIMS&RC, Bangalore
11. **Dr. BR Hungund** (*Associate Professor*) Pathology, JNMC, Belgaum.
12. **Dr. Sartaj Ahmad** (*Assistant Professor*),
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni** (*Associate Professor*)
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava** (*Medical Biochemistry*)
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu** (*General Medicine*) GSL Medical College, Rajahmundry,
Andhra Pradesh
4. **Prof. Pankaj Datta** (*Principal & Prosthodontist*)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro** (*Pediatrician*)
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra** (*Additional Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi** (*Pathologist*) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
8. **Prof. G S Meena** (*Director Professor*)
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna** (*Community Medicine*)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra** (*Paediatrician & Executive Director*)
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr. Shailendra Handu**, *Associate Professor*, Pharma, DM (Pharma, PGI
Chandigarh)
12. **Dr. A.C. Dhariwal**: *Directorate of National Vector Borne Disease*
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245 **Electronic-ISSN:** 0976-5506, **Frequency:** Quarterly
(Four issues per volume)

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Published at

Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

321. The Effect of Different Concentrations of the Blue-Green Algae Water Extract *Anabaenaoryzae* on the Adult Beetle (*Calosomainquisitor*) Order: Coleoptera for 24 Hours and 48 Hours after Treatment in Samarra, Iraq 1639
Athraa Hamid Jasim Al-rahmanny, Harith Ahmed Mustafa
322. Learning Cardiopulmonary Resuscitation (CPR) Using an Interactive Method for Students of an Elementary School in Indonesia 1642
Bintari Ratih Kusumaningrum, Ika Setyo Rini, Nikma Alfi Rosida
323. Concentrations of Lead, Cadmium and Mercury in Cord Blood and Prematurity in the Sidi Bel Abbes Region (West of Algeria) 1647
Bouhadiba Hadjer, DEMMOUCHE Abbassia, Menadi Norreddine, Mai Hichem, Ferrag Dalila, Zine Charaf Khalloua, Bekhadda Hadjer
324. The Potency of Private Practitioners on Tuberculosis Cases Finding and the Factors Associated in Surabaya City, East Java, Indonesia 1654
Chatarina Umbul Wahyuni, I Wayan Gede Artawan Eka Putra, Hari Basuki Notobroto, Abdul Rohim Tualeka
325. The Effect of Family Assessment of Less Chronic Energy (LCE) Pregnant on Nutritional Status of Pregnant and Baby Birth Size 1659
Demsa Simbolon, Antun Rahmadi, Jumiati
326. Safe Concentration of Mercury (Hg) Exposure in Fish Consumed by the Residents of Bulawa Subdistrict, Bone Bolango District, Gorontalo Province, Indonesia 1665
Deviyanti Wahyu Izati, Abdul Rohim Tualeka, Siprianus Singga, Pudhi Rahmawati, Syamsiar S Russeng, Atjo Wahyu, Ahsan
327. Evaluation of Abutment Health Related to Two Different Retention Methods of Maxillary Obturator (Randomized Control Trial) 1670
Doaa Tawfik Hassan, Hamzawi AG, Mohamed Mohamed Fata, Gehan Fekry Mohamed
328. Antibacterial Activity of Silver-Nanoparticles Synthesized Using *Trichoderma Harzianum* 1675
Ehab Y. Jabber, Mohamed M. Abdul Hussein
329. Incidence of Bacterial Vaginitis and Aerobic Vaginitis before and after Pessary Insertion in Indonesia 1680
Eighty Mardiyani Kurniawati, Faria Toma, Harry Parathon, Gatut Hardianto, Azami Azinar Denas, Kartuti Deborah, Firas Farisi Alkaff
330. Mother's Independence Model within Caring for Low Birth Weight Babies at Home after Hospital Care Based on Mother Factors, Family Support, and Social Support 1685
Erlina Suci Astuti, Nursalam Nursalam, Shrimarti Rukmini Devy, Risa Etika
331. The Evaluation of My Home My Village Method to Support the Complete Basic Immunization Programme in Surabaya, Indonesia 1691
Fariani Syahrul, Hario Megatsari, Ratna Dwi Wulandari, Arief Hargono, Kurnia Dwi Artanti
332. Safe Concentration of Benzene on Workers at Public Gas Stations around Diponegoro University Semarang, Indonesia 1697
Fatin Zuhra, Abdul Rohim Tualeka, Dimas Triyadi, Pudji Rahmawati, Syamsiar S Russeng, Atjo Wahyu, Ahsan
333. Operations Strategy as a Strategic Reconciliation towards a World Class University in Indonesia 1703
Febriana Wurjaningrum, Armanu, Fatchur Rohman, Mintarti Rahayu

The Potency of Private Practitioners on Tuberculosis Cases Finding and the Factors Associated in Surabaya City, East Java, Indonesia

Chatarina Umbul Wahyuni¹, I Wayan Gede Artawan Eka Putra^{1,2}, Hari Basuki Notobroto¹, Abdul Rohim Tualeka³

¹Faculty of Public Health, Universitas Airlangga; ²School of Public Health, Faculty of Medicine, Universitas Udayana; ³Department of Occupational Health and Safety, Faculty of Public Health, Universitas Airlangga

ABSTRACT

Private practitioners (PPs) have an important role for TB cases findings. The study aimed to assess the characteristic and potency PPs on TB cases finding. This was an operational research conducted from April until August 2018. PPs is general practitioners who have valid license and private practice session in Surabaya city. Data were collected through face to face interview using a structured questionnaire. The study succeeded interviewing 86 PPs, 35(40.7%) found presumptive TB at the last 3 months, 30(85,7%) of them refer the presumptive TB for smear examination and 21(60.0) manage the presumptive until definite the TB status. PPs who have average patients a day 15 persons and more have higher potency to found presumptive TB. This study shows the high potency and good willingness of PPs on presumptive TB and cases finding. First priority is involving PPs who have average patients a day 15 persons or more.

Keywords: *Private practitioners, tuberculosis cases finding, public-private mix.*

Introduction

The National Tuberculosis Control Program (NTP) in Indonesia faces the low of tuberculosis (TB) cases finding. In 2016, the national Cases Notification Rate (CNR) were 128 per 100.000 persons and 2017 were 152 per 100.000 persons. There is a large gap between CNR compare to the TB prevalence (660 per 100.0000 persons). The National Cases Detection Rate also did not reach the target. In 2016 the CDR were 33% and 2017 were 40%^{1,2}. One of important strategy to increase TB cases finding is to involve all private health care providers in NTP (NTP Private Practitioners partnership)³. Private practitioners (PPs) have an important role for TB cases findings and Directly Observe Treatments Short-course (DOTS) expansion⁴.

The involvement of PPs in health services delivery is an important strategy in health systems strengthening⁵. The engagement of PPs on NTP were roled in *public-private mix* (PPM) scheme. Base on national TB prevalence survey more than 40% people with TB symtoms sought treatment to PPs. Hence, the contribution of PPs to presemptive and TB cases finding relatively low compare to public health center (PHC) and hospital. A study involving telephone interviews of 25% of the private practitioners in Jogjakarta Province another Indonesian city shows that most PPs (63%) reported to have seen TB presumptive in their private practice⁶.

Improving the contribution of PPs will effect to increase and accelerate TB cases finding. TB cases that found by PPs mostly at earlier stage and this is give good effect to prevent TB infection⁷. Many related studies show the involvement of PPs on TB control program, factors associated and intervention supporting but in the setting of Surabaya a comprehensive study is need to assess the characteristic and potency PPs on TB cases finding in Surabaya City on DOTS implementation. This study aimed to assess the potency of PPs on tuberculosis cases finding and identify the factors associated to their potency.

Corresponding Author:

Abdul Rohim Tualeka
Department of Occupational Health and Safety,
Public Health Faculty, Airlangga University
Kampus C, Jalan Mulyorejo, Surabaya, 60115, Indonesia
Phone: +62 81 333 519 732
Email: inzut.tualeka@gmail.com

Method

Design: This was an operational research for evidence support to improve program performance. The design is cross-sectional study.

Study Settings: The study conducted for 5 months since from April until August 2018 in Surabaya City, the capital of East Java Province, Indonesia. Surabaya city is the second biggest city in Indonesia with population in 2017 was 3.057.766 persons.

Study population and sampling design: The study population private practitioners in Surabaya City, East Java Province. PPs were general practitioner who registered in the medical care unit of health office. We were selecting and visiting all PPs in East Surabaya as the samples of this study. We divided the interviewers in certain area and responsible visiting all PPs (sweeping) in that area. Through this strategy we could validating the PPs data from the register and minimize the sampling bias.

Data Collection and Analysis: PPs are general practitioners who have valid license and private practice session in Surabaya city. Potency defined as presumptive TB found at last 3 months at the practice site. Data were collected through face to face interview using a structured questionnaire. We collected information about the characteristics of the private practitioners (age, sex, qualification, type of practice), number of patient a day, opening hours per day and potency of private practitioners.

We trained 16 interviewers to perform the data collection. They trained to approach private practitioners and make an interview. Data was analysed descriptively and delivered in table and narration. Logistic regression was performed to identify characteristic and factor that associated the potency. The ethical clearance was obtained from The Ethical Committee of Faculty of Public Health, Universitas Airlangga.

Results

The Characteristic of Private Practitioners: The study succeeded interviewing 86 PPs with average of age 36.7 years old, 44 (51.2%) were male, 74(86.0%) were full time private practice and 57(66.3%) were practice with associate (group). Average length of practice were 4.5 years and average patients load a day were 14,2 persons (Table 1).

Table 1: The Characteristic of Private Practitioners

Characteristic		(n = 86)
Age	means (SD)	36.7 (11.1)
	< 35 years old	47 (54.7)
	≥ 35 years old	39 (45.3)
Sex	Male	44 (51.2)
	Female	42 (48.8)
Main occupation	Full time private Practitioners	74 (86.0)
	Government employee	12 (14.0)
Type of practice	Group	57 (66.3)
	Individual	29 (33.7)
Length of practice (year)	Means (SD)	4.5 (5.3)
	< 5 years	28 (32.6)
	≥ 5 years	58 (67.4)
Average number patient a day	Means (SD)	14.2 (11.8)
	<15 persons	52 (60.5)
	≥15 persons	34 (39.5)

SD = Standard deviation

The Potency of Private Practitioners on TB Cases Findings: PPs who found presumptive TB at the last 3 months were 35(40.7%), most of them, 30(85.7%) refer the presumptive TB for smear examination) to PHC or private laboratory. 18 (51.4%) PPs say that they also refer the presumptive TB for chest X-Ray. Most of PPs who found presumptive TB, 21(60.0) managed the presumptive until definite the TB status (Table 2).

Table 2: The Potency of Private Practitioners on TB Cases Findings

Variables		(n = 86)
Found presumptive TB cases at last 3 months	No	51 (59.3)
	Yes	35 (40.7)
Refer the patient to PHC or Lab for smear examination? (n = 35)	No	5 (14.3)
	Yes	30 (85.7)
Refer the patient for chest X-ray? (n = 35)	No	17 (48.6)
	Yes	18 (51.4)
The examination complete until the TB diagnosis confirmation (n = 35)	No	35 (40.0)
	Yes	21 (60.0)

The Factors Associated to Presumptive TB Cases Finding By Private Practitioners: The characteristic of PPs who found more presumptive TB were male and have average number of patients ≥ 15 persons a day. The PPs who practice with their associate (groups) and have good knowledge regarding TB controls program also tend to found more presumptive TB. PPs with average patients a day 15 persons and more have higher potency to found presumptive TB (AOR: 10.5; 95%CI: 3.4-32.4). Male PPs have higher potency to found presumptive TB compare to female (AOR: 3.3; 95%CI: 1.04-10.6) Table 3.

Table 3: Factors Associated to Presumptive TB Cases Finding By Private Practitioners

Variables	Found Presumptive TB Cases At Last 3 Months		Simple Logistic Regression		Multiple Logistic Regression	
	No	Yes	OR (95%CI)	p value	AOR (95%CI)	p value
Age, means (SD)						
< 35 years old	29(61.7)	18(38.3)	ref		ref	
≥ 35 years old	22(56.4)	17(43.6)	1.2(0.5-3.0)	0.619	0.8(0.2-3.4)	0.728
Sex						
Female	30(71.4)	12(28.6)	ref		ref	
Male	21(47.7)	23(52.3)	2.7(1.1-6.7)	0.027	3.3(1.1-10.7)	0.041
Main occupation						
Full time Private Practitioners	43(58.1)	31(41.9)	ref		ref	
Government employee	8(66.7)	4(33.3)	0.7(0.2-2.5)	0.577	0.6(0.1-3.3)	0.585
Type of practice						
Group	30(52.6)	27(47.4)	ref		ref	
Individual	21(72.4)	8(27.6)	0.4(0.2-1.1)	0.081	0.4(0.1-1.5)	0.176
Length of practice (year)						
< 5 years	18(64.3)	10(35.7)	ref		ref	
≥ 5 years	33(56.9)	25(43.1)	1.4(0.5-3.5)	0.514	2.0(0.4-8.6)	0.376
Average number patient a day						
<15 persons	30(71.4)	12(28.6)	ref		ref	
≥ 15 persons	21(47.7)	23(52.3)	11.7(4.2-32.6)	<0.001	10.4(3.4-32.1)	<0.001
Knowledge regarding TB program						
Lack	39(66.1)	20(33.9)	ref		ref	
Good	12(44.4)	15(55.6)	2.4(1.0-6.2)	0.061	2.8(0.8-9.6)	0.099

Discussion

This study assessed the PPs potency on presumptive and TB cases findings. Results show that most of the PPs were found presumptive TB cases on their practice site. The high potency shows the important partnerships TB program with PPs. A good model collaboration should be built to involve and improve the TB cases finding from PPs. For the first step, the program may priorities to certain characteristic of PPs. PPs who have more patients a day, practice in group should be involved first on collaboration scheme. Prioritization will make the collaboration more feasible and cost effective ⁸ to impact the TB incidence world over, there is an urgent need to address and accelerate TB control activities in the country. Nearly, half of the TB patients first seek TB care

in private sector. However, the participation of private practitioners (PPs. This finding is similar compare to study regarding public private mix (PPM) in Pakistan that found and suggest selection criteria for PPs involved in the TB program. The selection criteria were important to prevent failure and increase the cost effective ^{9,10}.

The potency of PPs in presumptive TB finding are quite high because we identify 59.3% of PPs found presumptive TB cases at last 3 month and most of them (85.7%) refer presumptive TB cases to the public health center for smear examination. The results of potency are lower compare to previous study in Jogjakarta that found 63.4%, this because we assess only in 3 months. In term of referring presumptive TB our result are higher compare to Jogjakarta only 41.5% ⁶. Their potency and

good willingness should be followed and accommodated by sustain response and care. The engagement of PPs on TB case finding will reduce diagnostic delay and TB transmission¹¹.

High potency of PPs on TB presumptive and cases finding also indicate opportunity involve them on TB reporting. PPs who found presumptive and TB cases should report to the program. This will decrease TB under reporting in private sector. An innovative collaboration model should be arranged to involving PPs. Previous study that conducted in several countries including Indonesia found that PPM intervention not yet succeeded to increase TB cases finding¹². Lesson learn from previous study and intervention regarding PPM is needed.

Innovative approach through good communication and trust building strategies should be arrange to encourage PPs involve on TB program¹³. Training and education for PPs should focus on achieving the minimum standard of TB diagnosis, drug adherence guaranty, prevent the using of inappropriate second line drugs that could lead to more cases of drug resistant TB and simple reporting to program¹⁴. Furthermore, intervention to engage PPs should consider the notification and reporting mechanisms. Notification mechanisms should address their perceived barriers, improve the efficiency and simplicity of the notification process. Internet and mobile phones may be use in reporting presumptive and TB cases by PPs to the program¹⁵.

Previous study in Bandung, the other city in Indonesia, learn about the effectiveness of face to face education using catharsis education action (CEA) method in improving the adherence of PPs to national guideline on management of TB. The study found that this method as effective as brief reminder with provision of pamphlet in improving the adherence of PPs to national guideline on management of TB. This finding indicate a psychoeducational strategies is important in influencing physician decision and behaviour¹⁶. This method is a counselling technique that brings out the psychological concerns that result from wrong perception of reality and hinder appropriate behaviour¹⁷. This method may an alternative method to inform and encourage PPs to participate on TB program.

These study findings have important policy implications. First, the study shows as the high potency of PPs involved in TB control program. Second, the

prioritization should be arranged for first step program to involve PPs in TB control program in Surabaya. The priority is involving the male PPs and who have average patients a day 15 persons or more, practice in group and increase their knowledge regarding TB control program. The involvement of PPs should be arrange in a bold collaboration and good evaluation. The collaboration program should be support for increase the factor associated to PPs contribution on referring the presumptive TB such as supervising or visiting PPs practice site, providing referral forms of presumptive TB and give the feedback of the examination results¹⁸.

Conclusions

This study shows the high potency and good willingness of PPs on presumptive and TB cases finding. The involvement of PPs in Surabaya City on TB cases finding should be arrange in a bold collaboration. First priority is involving the male PPs and who have average patients a day 15 persons or more and increase their knowledge regarding TB control program. The method and materials for PPs training and education should be arrange in a workshop involving their professional organization.

Acknowledgements

The authors would like to thank Research Department of Universitas Airlangga for funding this research. The authors are grateful to thank all private practitioners who kindly consented to participate in this study. We sincerely thank to Health Office of Surabaya City for their support and provision of TB related epidemiological information.

Ethical Clearance: Taken from Ethics Committee of the Public Health Faculty of Airlangga University with ethics number 189-KEPK

Source of Funding: The article "The Potency of Private Practitioners on Tuberculosis Cases Finding and The Factors Associated in Surabaya City, East Java, Indonesia" was supported by Faculty of Public Health, Airlangga University, Indonesia, 2019.

Conflict of Interest: The authors declare that we have no competing interests.

REFERENCES

1. Balitbangkes Kemenkes RI. *Laporan Survei Prevalensi Tuberkulosis 2014*. Jakarta; 2015.
2. Pusdatin Kemenkes RI. *Tuberkulosis: Temukan Obat Sampai Sembuh 2016*. Jakarta; 2016. doi:24442-7659
3. Ministry of Health Republic of Indonesia. *National Strategy for TB Control in Indonesia 2010-2014*. Jakarta; 2011.
4. Kemenkes RI. *Rencana Strategis Kementerian Kesehatan 2015 - 2019*.; 2015.
5. Kaboru BB. Uncovering the potential of private providers' involvement in health to strengthen comprehensive health systems: a discussion paper. *Perspect Public Health*. 2012;132(5):245-252. doi:10.1177/1757913911414770
6. Mahendradhata Y, Utarini A, Lazuardi U, Boelaert M, Stuyft P Van Der. Private practitioners and tuberculosis case detection in Jogjakarta, Indonesia : actual role and potential. *Trop Med Int Heal Vol*. 2007;12(10):1218-1224. doi:10.1111/j.1365-3156.2007.01946.x
7. Ahmed J, Ahmed M, Laghari A, Lohana W, Ali S, Fatmi Z. Public private mix model in enhancing tuberculosis case detection in District Thatta, Sindh, Pakistan. *J Pak Med Assoc*. 2009;59(2):14-16.
8. Anand T, Babu R, Jacob AG, Sagili K, Chadha SS. Enhancing the role of private practitioners in tuberculosis prevention and care activities in India. *Lung India*. 2017;34(6):538-544. doi:10.4103/0970-2113.217577
9. Pethani A, Zafar M, Khan AA, Rabbani Sana U, Ahmed S, Fatmi Z. Engaging General Practitioners in Public-Private Mix Tuberculosis DOTS Program in an Urban Area in Pakistan: Need for Context-Specific Approach. *Asia-Pacific J Public Heal*. 2015;27(2):NP984-NP992. doi:10.1177/1010539513483828
10. Pantoja A, Floyd K, Unnikrishnan KP, et al. Economic evaluation of public-private mix for tuberculosis care and control, India. Part I. Socio-economic profile and costs among tuberculosis patients. *INT J TUBERC LUNG DIS*. 2009;13(December 2008):698-704.
11. Uplekar M. 10. Involving Private Health Care Providers in Delivery of Tb Care : Global Strategy. <http://medind.nic.in/haa/t08/i1/haat08i1p38.pdf>.
12. Hinderaker SG, Rusen ID, Chiang C-Y, Yan L, Heldal E, Enarson D a. The FIDELIS initiative: innovative strategies for increased case finding. *Int J Tuberc Lung Dis*. 2011;15(1):71-76. <http://www.ncbi.nlm.nih.gov/pubmed/21276300>.
13. Philip S, Isaakidis P, Sagili KD, Meharunnisa A, Mrithyunjayan S, Kumar AM V. "They Know, They Agree, but They Don't Do" - The Paradox of Tuberculosis Case Notification by Private Practitioners in Alappuzha District, Kerala, India. *PLoS One*. 2015;10(4):1-14. doi:10.1371/journal.pone.0123286
14. Bharaswadkar S, Kanchar A, Thakur N, Shah S, Patnaik B. Tuberculosis Management Practices of Private Practitioners in Pune Municipal Corporation, India. 2014;9(6). doi:10.1371/journal.pone.0097993
15. Yeole RD, Khillare K, Chadha VK, Lo T, Kumar AM V. Tuberculosis case notification by private practitioners in Pune, India: how well are we doing? *Public Heal Action*. 2015;5(3):173-179. doi:10.5588/pha.15.0031
16. Arisanti N. The effectiveness of face to face education using catharsis education action (CEA) method in improving the adherence of private general practitioners to national guideline on management of tuberculosis in Bandung, Indonesia. *Asia Pac Fam Med*. 2012;11(1):2. doi:10.1186/1447-056X-11-2
17. Lukens EP, Mcfarlane WR. Psychoeducation as Evidence-Based Practice : Considerations for Practice, Research, and Policy. *Br Treat Cris Interv*. 2004;4(3):205-225. doi:10.1093/brief-treatment/mhh019
18. Putra IWGAE, Utami NWA, Suarjana IK, et al. Factors associated to referral of tuberculosis suspects by private practitioners to community health centres in Bali Province, Indonesia. *BMC Health Serv Res*. 2013;13(1):1. doi:10.1186/1472-6963-13-445