

The Influence of Age And Coping Mechanism on The Resilience of Cancer Patients Undergo Chemotherapy

by Kusnanto Kusnanto

Submission date: 15-Mar-2019 11:27AM (UTC+0800)

Submission ID: 1093640995

File name: 1..pdf (380.81K)

Word count: 5625

Character count: 30788

The Influence of Age And Coping Mechanism on The Resilience of Cancer Patients Undergo Chemotherapy

Muhammad Saleh Nuwa¹, Kusnanto², Sri Utami³

¹Master Student Faculty of Nursing, Airlangga University Surabaya

²Lecturer of nursing department , Airlangga University, Surabaya

³Lecturer in Poltekkes Kemenkes Surabaya

{(muhammad.saleh.nuwa-2016@fkip.unair.ac.id) (kusnanto_ners@yahoo.com) (bu.sri.utami67@gmail.com)}

Keywords: Resilience, Coping Mechanism, Cancer, Chemotherapy, Ag

Abstract: **Background :** Resilience is the dynamic capacity of individuals to successfully retain their mental health in the face of life difficulties. Chemotherapy makes patients who have been diagnosed with cancer have a sense of worry, anxiety and fear of facing death threats. Resilience may depend on positive emotions, flexible thinking, age and spirituality. This study aims to determine the effect of age and coping mechanism on the resilience of cancer patients undergo chemotherapy. **Method:** The study was conducted on 60 respondents. The coping mechanism was measured using 21 items of The Cancer Coping Questionnaire, and the resilience was measured by The 14 item Resilience Scale. Multivariate multiple linear regression analysis was used to predict the effect of age factor and coping mechanism on the resilience of cancer patients undergo chemotherapy. **Results :** The result showed that 60% of respondents had maladaptive coping mechanism 83,33% had low enough resilience and only 5% had high resilience. Pearson correlation shows an association between age and resilience, as well as coping mechanisms with resilience. The effect size of age and coping mechanism on resilience is 44, 9%. **Conclusion :** Age, coping mechanism and resilience are interrelated factors in cancer patient. Adaptive coping mechanism in cancer patients can increase their resilience undergo chemotherapy.

1 INTRODUCTION

Cancer is still a major health problem in the world and the second leading cause of death after heart and blood vessel disease. A total of 8.2 million people died from cancer each year (Komiya et al. 2017). According to data from GLOBOCAN, the International Agency for Research on Cancer (IARC) found that in 2012 there were 14,090,100 new cases of cancer and 8,201,600 deaths from cancer worldwide. A total of 5.3 million deaths occurred in developing countries and 2.9 million occurs in developed countries (American Cancer Society 2015). This number increased to 8.8 million deaths by 2015. Globally, almost 1 in 6 deaths are caused by cancer and approximately 70% of deaths caused by cancer occur in poor and developing countries.

The number of cancer cases has been increased in Indonesia, slowly began to replace the position of heart attack as the main caused of death. Basic Health Research Reported (2003) the prevalence of

cancer was 1.4 per 1,000 residents or approximately (347,000) people and the cause of death number 7 (5.7%) from all death in Indonesia (KEMENKES RI 2013).

The current efforts undertake to manage cancer, carried out in various way including surgery followed by radiotherapy, treatment with three combinations of surgery, radiation and chemotherapy. Chemotherapy is the treatment of antineoplastic preparation for an attempt to kill tumor cells by interfering with cellular function and the reproduction (Yusof et al. 2016). According to Jacobson (2009), the problem of chemotherapy is a very harmful and adverse side effects. The side effect of chemotherapy on physical problems has been clearly illustrated by Chan & Ismail (2014);Lorusso et al. (2016), the most common signs and symptoms are nausea and vomit, decreased appetite, hair loss, bone marrow damage, neuropathy, gastrointestinal disorders and skin damage. Chemotherapy also greatly affects the sexual life, daily activities and work.

According to Sana et al. (2016) cancer treatment can cause many health problems and treatment-related symptoms may last longer and can add great pressure on patients to cope with demands and treatment of disease. In addition, the accompanying consequences on chemotherapy make the majority of patients that have been diagnosed with cancer filled with worry, anxiety and fear of facing the threat of death and pain while undergo therapy. They also experience problems of interpersonal relationship so that cancer patients need individual psychological adjustment. Another opinion by Partridge et al (2007) depression and stressors that appear during treatment, length of therapy, poor communication between health worker and patient, high medical costs and poor side effects are the contribution for patients to not comply chemotherapy.

Beside that, at the time patient being diagnosed with cancer these circumstances was very difficult to accept. The long-term effects of cancer among adolescents and young adults have been known to disrupt normal developmental processes, poor mental health, impaired quality of life, social isolation, and impaired education (Rosenberg et al. 2017). In a large-scale survey 73% of cancer patients who undergo treatment two years after diagnosed with cancer, Wolff (2007) found that over 70% of respondents reported being depressed due to the disease, 60% reported relationship problems, and 83% reported declines on income. More than a quarter indicates that they have insufficient resources to fulfil the emotional needs.

According to Loprinzi et.al (2011) one of the possible ways for newly diagnosed cancer patients to control emotional stress are through good coping mechanism. Krohne (2002) says coping strategy is an individual adaptation mechanism that is done consciously and directed towards overcoming pain or facing stressor. According Yunitasari (2016) adaptive coping in cancer patients can be achieved by minimizing and even eliminating the cause of stressor. A good coping mechanism in cancer patients undergo chemotherapy will improve resilience.

According to Hodges et al (2010) resilience is a transformation, a deliberate desire to withstand environmental complexity and unfavorable uncertainty. Resilience is a dynamic process that includes positive adaptation in the context of significant difficulties, hazardous, and may change with time and in different environments (Fletcher & Sarkar 2013). Resilience may depends on several factors including positive emotions, flexible

thinking such as acceptance, active coping and spirituality. (Portzky et al. 2010). Wagnild (2003) identify five characteristics of resilience that become the main component of a person in responding to the difficulties they might experiences. The five components are meaningfulness, Equanimity, Self Reliance, Perseverance and Existential Aloneness ((Boyle et al. 2015)

Some studies related to resilience in cancer patients showed that most cancer patients have less resilience to the diagnosis of cancer and chemotherapy they undergo (Pertiwi et al. 2011; Dubey et al. 2015; Proyogi & Agung 2016).

Individuals diagnosed with cancer in younger ages (before 45) has been found to be at high risk of psychological problems, which can survive in the development of life (Hoffman et al. 2013). Research Gaffey et al (2016) on stress and resilience in the elderly showed that resilience sources could modulate cortisol in elderly health. This study aimed to determine the effect of age factor and coping mechanism on the resilience of cancer patients undergo chemotherapy.

2 METHODS

2.1 Study Design

This research is an analytic descriptive research, with cross sectional design. In this design, researchers wanted to know the effect of age and coping mechanisms on the resilience of cancer patients who undergo chemotherapy at the General Hospital of East Nusa Tenggara province Indonesia in patients with cancer diagnosis from 2016 to January 2018.

2.2 Sampling

A total of 60 respondents that were undergo chemotherapy were taken as a research sample of 93 cancer patients treated in oncology ward and outpatient department of general hospital area with purposive sampling technique.

Purposive sampling is a technique of determining samples by selecting the population sample in accordance with the researcher desired so that the sample can represent the characteristics of the previously known population (Nursalam 2017)

Samples were selected with the following inclusion criteria: (1) Patients with cancer

diagnose, who undergo chemotherapy aged 21-70 years, (2) able to read and written in Indonesian, (3) level awareness is composmentis, (4) cancer patients who undergo maximum 5 sessions of chemotherapy, (6) wanted to be respondents by signing the informed consent. While the exclusion criteria of this study were: (1) pediatric patients, (2) cancer patients undergo chemotherapy who had comorbidities more than 2, (3) cancer patients with psychiatric disorders.

2.3 Procedure

The study protocol was approved by the research ethics commission of the University of Airlangga Surabaya Indonesia. Prior to the data collection the researcher gave the respondent information conducted by 3 enumerators with the qualification professional of nursing. Data were collected in the beginning of February 2018 on respondents who were undergo chemotherapy.

Patient demographic data were collected through questionnaires while clinical data were collected from patient medical records, Patient completed coping questionnaire and resilience in the outpatient department and chemotherapy ward prior to scheduled chemotherapy sessions or after chemotherapy. Questionnaire filling takes about 10 minutes for each respondent and accompanied by an enumerator.

2.4 Research Instrument

The measuring tool used in this study was questionnaire with aim to measure coping mechanism and resilience. While age is the basic data obtained directly from the patient through the demographic data questionnaire.

2.4.1 Resilience Measurement

Resilience was measured using the Resilience Scale Resource Kit developed by Wagnild & Young (1993). The selection of this measuring instrument because this tool is the widest use until now. This measuring tool has also been used in adolescents, adults, and elderly for assessed according to the characteristics of respondents. Initially this measuring instrument consist of 50 items, after the analysis, the item was reduced to 25 items and repaired into 14 items reflecting the five components of Resilience: Equanimity,

Meaningfulness, Perseverance, Self-reliance and Existential aloneness.

This measuring instrument has a high reliability coefficient of 0.84 until 0.94. (Pinheiro et al. 2015; Ntountoulaki et al. 2017). According to Kaplan and Sacuzzo (2005), one of the requirements of a good measuring instrument is to have a reliability coefficient of 0.7 until 0.8.

2.4.2 Coping Measurement

The respondents coping mechanism were measured using *The Cancer Coping Questionnaire* developed by Moorey et al. (2003). This *Self Rating Scale* measuring instrument consists of 21 items is a special measuring tool used to measure coping mechanism in cancer patients that have been tested in 201 cancer patients. This questionnaire has 2 general questions related to stress and anxiety about cancer, the next items question 1-14 about the individual scales consisting of sub-coping scale (items 2, 6, 7, 11, 12), positive focus (Items 1, 9, 14), Transfer (item 3, 4, 8), and Planning (Item 5, 10, 13). For items 15-21 is an assessment for interpersonal scale. This instrument has an internal consistency of 0.87 for individual scale and 0.82 for interpersonal scale. Patients coping mechanism then divided into adaptive and maladaptive based on the mean and standard deviation of the respondents.

2.5 Data Analysis

The categorical data is presented as the sum and percentage. The age data of coping mechanism and resilience are presented in the form of mean calculation. The Pearson product moment coefficient correlation was determined to evaluate the linear relationship between age and resilience, as well as coping mechanism and resilience. Multiple linear regression analysis was conducted to determine the factors independently affect the resilience factor of value $P < 0.25$ on Pearson correlation test will be incorporated into multiple linear regression model. Statistical analysis was performed using IBM software statistics 21. $P < 0.05$ on a two-tailed test was statistically significant.

3. RESULT

19

3.1 Demographic and Clinical Characteristics

Table 1. Summary of demographic and clinical characteristics of 60 cancer patients that undergo chemotherapy

Characteristics		N= 60
Age	Mean ± SD	49,12 ± 10.37
	Adolescent	2 (3,3 %)
	Young Adult	4(6,7 %)
	Adult	13 (21,7 %)
	Young Elderly	26 (43,3 %)
Elderly		15 (25 %)
Gender	Male	26 (43,3 %)
	Female	34 (56,7 %)
Education level	Primary School	13 (21,7 %)
	Junior High School	23 (38,3 %)
	Senior High School	17 (28,3 %)
	College/ above	7 (11,7 %)
Marital status	Married	52 (86,7 %)
	Single	5 (8,3 %)
	Widow /Widower	3 (5 %)
Occupation	Civil Servant	7 (11,7 %)
	Housewife	25 (41,7 %)
	Private sector	5 (8,3 %)
	Farmer	23 (38,3 %)
Income	High salary	19 (31,7 %)
	Low salary	41 (68,3 %)
Type of cancer	SC/PC/OsC/TC	1(1,7%)1(1,7%)1
	/PaC/PrC/	(1,7)1(1,7%)1(1,
	EC/CoC/CR/N	7)1(1,7%)2(3,3%
	C/GSCC/OvC/)2(3,3%)2(3,3%)
	LNH/CeC/BC	/3(5%)4(6,7%)4(6,7%)6(10%)7(11,7%)24 (40%)
Stadium of cancer	Stadium II	24 (40 %)
	Stadium III	36 (60 %)
Chemotherapy sessions	1 Session	2 (3,3 %)
	2 Sessions	16 (26,7 %)
	3 Sessions	14(23,3 %)
	More than 3 sessions	28 (46,7 %)
Body mass index	<18,5	37 (61,7 %)
	18,5-25	18 (30 %)
	>25	5 (8,3%)
Comorbidity	Yes	14 (26,3 %)
	No	46 (76,7 %)

SC=scapula cancer, PC=Penis Cancer, OsC=Osteosarcoma Cancer, TC=Tongue Cancer, PaC

=Parotid Cancer, PrC= peritoneal cancer, EC= Endometrium Cancer, CoC= Colon cancer, CR= Carcinoma Rectum, NC=Nasofaring Cancer,GSCC= Gingival Squamous Cell Carcinoma, OvC= Ovarian Cancer, LNH= Limfoma Non Hodgkin, CeC= Cervical Cancer, BC= Breast Cancer

A total of 60 cancer patients who undergo chemotherapy had an average age of 49.12 years (SD = 10.37 years), and 43% were young elderly, 25% of elderly, 21.7% adult, 6.7% young adult, and 3.3% are teenagers.

Patients diagnosed with cancer on stage III were 60%, the remaining 20% were stage II, with the most common type of breast cancer being 24 (40%) patients. Other details of the patient's demographic and clinical characteristics are listed in Table 1. Coping and resilience mechanism in Table 2.

Table 2. Measuring instrument scores from 60 cancer patients undergo chemotherapy

Resilience	Mean ± SD	68,62 ± 5,415
	High Resilience	3 (5%)
	Average Resilience	50 (83,3%)
	Low Resilience	7 (11,7%)
Coping Mechanism	Mean ± SD	56,33± 5,332
	Adaptive	30 (50%)
	Maladaptive	30 (50%)
Individual Scale of Coping Mechanism		
Positive focus	Mean ± SD	8,100± 1,03
	High	44 (73,3%)
	Low	16 (26,7%)
Coping	Mean ± SD	12,87± 1,96
	High	32 (53,3%)
	Low	28 (46,7%)
Diversion	Mean ± SD	7,53± 1,112
	High	28 (46,70%)
	Low	32 (53,3%)
Planning	Mean ± SD	7,43± 1,14
	High	29 (48,3%)
	Low	31 (51,7%)
Interpersonal Scale of Coping Mechanism		
	Mean ± SD	20,48± 2,05
	High	31 (51,7 %)
	Low	29 (48,3%)

Adaptive coping mechanism = score > 56,33, maladaptive coping mechanism = score < 56,33. Total score resilience >90 = very high resilience, 82-90 high resilience, 65-81 moderate resilience, 57-64 low resilience and score resilience <57 = very low resilience.

3.2 Relationship between age and coping mechanism with resilience

13

Table 3. Summary of Pearson correlation coefficients of age and coping mechanism with the resilience of cancer patients undergo chemotherapy

	Resilience
Age	0,274*
Coping mechanism All score	0,654**
Positive focus score	0,481**
Coping score	0,586**
Diversion score	0,524**
Planning score	0,513**
Interpersonal scale of Coping mechanism	0,457**

*Correlation is significant at the 0.05 level (2-tailed).

**Correlation is significant at the 0.01 level (2-tailed).

The result showed a positive correlation between age and resilience $\rho = 0.34$, meanwhile for coping mechanism showed a positive correlation between coping mechanism and resilience $\rho = 0.000$. Furthermore, for individual scale the coping mechanism of 60 respondents indicates a positive correlation for each sub item positive focus $\rho = 0.000$, coping $\rho = 0.000$, diverse $\rho = 0.000$, and planning $\rho = 0.000$. On the interpersonal scale also shows a positive correlation between resilience $\rho = 0.000$ (table 3).

3.3 The Effect of age and coping mechanism on resilience

Multiple multivariate linear regression showed a large contribution of age influence and coping mechanism to the resilience of cancer patients undergo chemotherapy in this model was 44.9% ($\rho = 0.000$), while the remaining 55.1% was influenced by other variables than age and coping mechanism. Partially, the contribution of age to resilience is 2.19% (correlation part = 0.148, $\rho = 0.139$) while for coping mechanism is 37.45% (correlation part = 0.612, $\rho = 0.000$) (table 4). From this analysis it is known that age has no significant effect on resilience whereas opposite coping mechanism significantly affect the resilience of cancer patients undergo chemotherapy. The large prediction of age variable

with coping mechanism to resilience can be seen from the following regression equation.

$$\text{Resilience} = 29.052 + 0.079 (\text{age}) + 0.634 (\text{coping mechanism}).$$

The following equation can be explained as:

- The constant value about 29,052 means that the magnitude of resilience when the age value, and the coping mechanism is 0.
- An increase in 1 unit age score can increase the resilience value by 0.079 with age constant.
- An increase of 1 unit of coping mechanism score may increase resilience by 0.634 with a constant age.

Table 4. Summary of regression model, resilience on cancer patients undergo chemotherapy

Model	Coefficient	Classic assumption
R = 0,670	Constant = 29.052	Existency: mean residual= 0,000
R square = 0,449	β age = 0.079	Independency : Durbin-Watson=1,230
Adjusted R square = 0,430	β coping=0,634	Linearity: anova: p=0,000
Anova (p=0,000)	Correlations Part age : 0,148 Part coping mechanism :0,612	Multikolinearity : variance inflation factor : VIF age =1,041 VIF coping mechanism =1,041

Normality: age ($p=0.232$), coping mechanism ($p=0.602$), resilience ($p=0.472$).




Figure1. Classic assumption : Homocedasticity




Figure2. Classic Assumption: Multivariat Normality

4. DISCUSSION

Cancer and chemotherapy are very distressing life events, and this is the real case for cancer patients undergo chemotherapy at the General Hospital Prof. Dr. W.Z. Johannes Kupang, East Nusa Tenggara Province, Indonesia. Cancer patients mostly experience stress and anxiety because of nature of the disease and the treatment. Patients said anxious and think about the disease for several times 29 (48,3%), almost whole time 27 (45%) and the rest 4 (6,7%) patients said worried and thinking about the disease everytime. Regard to question what stress felt in the last week, the patient said less stress as much as 30 (50%) patients, quite stressful 26 (43.3%) and very stressful 4 (6.7%) patients. The impact of cancer and chemotherapy on these psychological problems lead to a worse disease progression for the individual (Zhang et al. 015). One effort that can be done is by build a good coping mechanism. According to Baqutayan (2015) there are two main ways in which people cope with stress. The first approach, one can decide to follow or reject the stress experienced. This is a passive approach. Alternatively, the other one can decide to face the reality of the stress experienced and to clarify the problem through negotiation. The second approach is an active approach.

The result showed 50% of patients still have maladaptive coping mechanism against cancer and chemotherapy. This can be seen from the score of coping mechanism on individual and interpersonal scale. The findings of this study are also known that patients use several actions to overcome the problems faced in the form of diversion efforts, coping, positive focus and planning that all of them is a kind of emotional focus coping. The result of

this study is accordance with Ahadi et al. (2014) in 80 cancer patients has found that the average value of coping in cancer patients lower than non-cancer. This study also known the cancer patients used more emotional focus coping to overcome problems related to cancer.

Dunkel et la. (1992) is known in general, cancer patients make a variety of coping choices. In the face of cancer-causing symptoms that cause pain, they usually choose to use problem focused on coping strategies, such as seeking alternative medicine or taking drugs, while facing future uncertainty they tend to use emotional focused on coping strategies such as dodge or denial. Furthermore, Faye et al. (2006) found that emotional focused coping is more commonly used by cancer patients to address existential problems, while problem focused on coping is more commonly used to overcome physical problems.

Researchers see the tendency of selecting coping problem in cancer patients to overcome the existing problems due to disease conditions that have entered the advanced stage and the success rate of low disease healing, where the patient has no other efforts to recover, in addition to continue to survive and fight with the disease cancer. Despite being in a stressful situation, cancer patients actually still have an inner strength that can help them to adapt to the conditions and make sense of life.

A good coping mechanism in cancer patients is needed so the patient can undergo the disease and can survive despite the downturn. This condition is called resilience. The result showed that as many as 83.3% of patients had average resilience scores. The result of the analysis revealed the effect of coping mechanism on the resilience of cancer patients undergo chemotherapy. The result of this study is accordance to Haase (2004) study that resilience in cancer patients and chronic diseases is the result of an interaction between three protective factors and two risk factors. The three protective factors in question are individual protective factors (coping courageous in the face of stressful situations and meaningful situations), family protective factors (family atmosphere and family support or resources), and social protective factors (health care and social integration resources). The two risk factors faced by the study subjects include individual risk factors (defensive coping) and disease-related risk factors.

Furthermore, Peterson & Bredow (2013) describes risk factors as a factor that directly magnifies the potential risk for individuals which

can increase the likelihood of developing maladaptive behaviors and lifestyles while the protective factor is a healthy skill and ability possessed by individuals, which promotes resilience. The low resilience scores in cancer patients in this study may be influenced by low cognitive capacity, stage of cancer, chemotherapy series treatment, body mass index, comorbidities and other demographic characteristics. This condition results in low of confidence to recover, lack of confidence and lack of optimism. This assumption accordance to the opinion of Portzky et al. (2010) explained the high level of resilience in a person is usually positively correlated with high self esteem, self confidence, and discipline, courage and optimism in the face of failure, above average cognitive capacity, and greater possibilities for free from disease.

A study of 60 women with ovarian cancer, it is known that those who have a greater tendency to attribute negative meanings to illness are more likely to show poor adjustment. Some patients may withdraw socially in response to diagnosis or treatment and treatment measures (Deshields et al. 2016)

The study also found that age had no significant effect on resilience ($p = 0.139$). Resilience is not a static trait that a person possesses from birth or automatically persists in a person once he or she achieves it. Resilience is a dynamic process that includes positive adaptation in the context of significant difficulties, hazardous, and may change with time and in different environment environments (Fletcher & Sarkar 2013). This view is in line with Galli & Vealey (2008) research on resilience among the top athletes and concludes that an important aspect of resilience is the agitation process, in which the individual uses multiple coping strategies to deal with unpleasant emotional combinations and struggles mentally. Many athletes report that positive adaptation occurs gradually, often requiring a lot of shifting thoughts. These findings support that the resilience is a capacity that develops over time in the context of people's interactions with the environment, regardless of age.

The greater effect of age and coping mechanism both in the resilience of cancer patients undergo chemotherapy in this study about 44.9% while the remaining 55.1% is influenced by other variables than age and coping mechanism out of this model. Pentz research (2005) on resilience in elderly patients with cancer is known that the

aspects that contribute to resilience are social support and spirituality aspect (ie belief in God and their hope). Another study was conducted by Duan porter et al. (2016) on factors that contribute to physical resilience in the elderly with cancer is known the majority of older cancer sufferer show physical resilience. This is related to basic health, physical function, self efficacy, and high social support.

Some of the research findings above in line with the opinions of (Vanderbilt-Adriance & Shaw 2009). There are several protective factors that make up resilience such as: (1) individual characteristics, such as gender, level of intelligence, personal characteristics, (2) family characteristics, such as warmth and family structure (3) availability of social support system outside the individual and family environment eg peers. Furthermore, according to Woodgate (1999) there are five categories of specific stressors that exist in cancer patients as risk factors affecting the resilience of cancer patients that is loss, disruption of relationships, events that change family status, events that require social adaptation, and acute negative event such as physical trauma. This stress may be direct or indirect result of major developmental changes occurring in the individual, or the consequences of the disease itself.

From the various opinion and result above it can be conclude that in principle resilience is influenced by internal and external factors of individuals, both protectives and risks. Age and coping mechanism are some of the factors that contribute to patient resilience. Both are mutually related internal protective factors in which the existing research evidence suggests that age maturity making individual coping mechanism more adaptive.

5. CONCLUSION

Age and coping mechanism are some of the protective factors that can affect the resilience of cancer patients. Increased age does not necessarily increase patient resilience or coping mechanism. A positive adaptation of cancer patients undergo chemotherapy is a key requirement for cancer patients to remain resilient in their lives.

REFERENCES

- Ahadi, H., Delavar, A. & Rostami, A.M., 2014. Comparing Coping Styles in Cancer Patients and Healthy Subjects. *Procedia - Social and Behavioral Sciences*, 116, pp.3467–3470.
- American Cancer Society, 2015. *Global Cancer Facts & Figures* 3rd ed., Atlanta: American Cancer Society.
- Baqutayan, S.M.S., 2015. Stress and Coping Mechanisms: A Historical Overview. *Mediterranean Journal of Social Sciences*, 6(2), pp.479–488.
- Barbara J. De faye, Keith G. Wilson, Susan Chater, Raymond A. Viola, P.H., 2006. Stress and coping with advanced cancer. *Palliative and Supportive Care*, 4, pp.239–249.
- Boyle, G.J., Saklofske, D.H. & Matthews, G., 2015. *Measures of Personality and Social Psychological Constructs*, Elsevier Inc. Available at: <http://dx.doi.org/10.1016/B978-0-12-386915-9.00011-5>.
- Chan, H. & Ismail, S., 2014. Side Effects of Chemotherapy among Cancer Patients in a Malaysian General Hospital: Experiences, Perceptions and Informational Needs from Clinical Pharmacists. *Asian Pacific Journal of Cancer Prevention*, 15(13), pp.5305–5309.
- Deshields L. Teresa, Heiland F. Mark, Kracen C. Amanda, D.P., 2016. Resilience in adults with cancer: Development of a conceptual model. *Psycho-Oncology*, 25(1), pp.11–18.
- Dubey C, Maria J. De, Hoepli C, Betticher D C, E.M., 2015. Resilience and unmet supportive care needs in patients with cancer during early treatment: A descriptive study. *European Journal of Oncology Nursing*, 19(5), pp.582–588.
- Dunkel C. Schetter, Feinstein L G, Taylor S E, F.R.L., 1992. Patterns of coping with cancer. *Health psychology: official journal of the Division of Health Psychology, American Psychological Association*, 11(2), pp.79–87.
- Fletcher, D. & Sarkar, M., 2013. Psychological Resilience A Review and Critique of Definitions, Concepts, and Theory. *European Psychologist*, 18(1), pp.12–23.
- GaffeyE. Allison, B.C.S. & Clark LA, W.M.M., 2016. Aging and the HPA axis: Stress and resilience in older adults. *Neuroscience and Biobehavioral Reviews*, 68, pp.928–945. Available at.
- Galli, N. & Vealey, R.S., 2008. “ Bouncing Back ” From Adversity: Athletes’ Experiences of Resilience. *The Sport Psychologist*, 22(2002), pp.316–335.
- Haase, J.E., 2004. The Adolescent Resilience Model as a Guide to Interventions. *Journal of Pediatric Oncology Nursing*, 21(5), pp.289–299.
- Hodges F. Helen, Troyan J. Patricia, K.C.A., 2010. Career Persistence in Baccalaureate-Prepared Acute Care Nurses. *Journal of nursing scholarship*, 42(1), pp.83–91.
- Hoffman, M.A., Lent, R.W. & Raque-Bogdan, T.L., 2013. A Social Cognitive Perspective on Coping With Cancer. *The Counseling Psychologist*, 41(2), pp.240–267.
- Jacobson O. Joseph, Polovich Martha, Mcniff K. Kristen, Lefebvre B. Kristine, Cummings Charmaine, GaliotoM ichele, Bonelli R. Katherine, M.R.M., 2009. American Society of Clinical Oncology / Oncology Nursing Society Chemotherapy Administration Safety Standards. *American Society of Clinical Oncology*, 27(32), pp.5469–5475.
- KEMENKES RI, 2013. *Riset Kesehatan Dasar (RISKESDAS 2013)*, JAKARTA: Badan LITBANG KEMENKES RI.
- Komiya, T., Mackay, C.B. & Chalise, P., 2017. Who treats lung cancer? Results from a global. *Respiratory Investigation*, 55(5), pp.308–313.
- Krohne, H.W., 2002. *Stress and Coping Theories*, Germany.
- Loprinzi E. Caitlin, Prasad Kavita, Schroeder R. Darrell, S.A., 2011. Stress Management and Resilience Training (SMART) Program to Decrease Stress and Enhance Resilience Among Breast Cancer Survivors: A Pilot Randomized Clinical Trial. *Clinical Breast Cancer*, 11(6), pp.364–368.
- Lorusso Domenica, Bria Emilio, Costantini Anna, Massimo Di Maio, Rosti Giovanni, M.A., 2016. Patients’ perception of chemotherapy side effects: Expectations, doctor – patient communication and impact on quality of life – An Italian survey. *European Journal of Cancer Care*, (November), pp.1–9.
- Moorey, S., Frampton, M. & Greer, S., 2003. The Cancer Coping Questionnaire: a self-rating scale for measuring the impact of adjuvant psychological therapy on coping behaviour. *Psycho-oncology*, 12(4), pp.331–44.
- Ntountoulaki Elisavet, Paika Vassiliki, Kotisis Konstantinos, Papaioannou Dimitra, Andreoulakis Elias, Fountoulakis N. Konstantinos, Carvalho F. Andre, H.T., 2017. The Greek Version of the Resilience Scale (RS-14): Psychometric Properties in three Samples and Associations with Mental Illness, Suicidality, and Quality of Life. *Journal of Psychology and Clinical Psychiatry*,

- 7(5).
- Nursalam, 2017. *Metodologi Penelitian Ilmu Keperawatan: Pendekatan Praktis* 4th ed., Jakarta: Salemba Medika.
- Partridge H. Ann, Ades Terri, Spicer Patricia, Englander Linda, W.D.L., 2007. Helping breast cancer patients adhere to oral adjuvant hormonal therapy regimens. *Community Oncology*, 4(12), pp.725–731.
- Pertiwi, M.P., Karini, S.M. & Agustin, R. widya, 2011. Relationship Between Resiliency and Knowledge about Treatment Breast Cancer With Patient Compliance in RSUD DR. Moewardi Surakarta.
- Peterson, S.J. & Bredow, T.S., 2013. *Middle range theories: application to nursing research* 3rd ed., philadelphia new york: Wolters Kluwer Health | Lippincott Williams & Wilkins.
- Pinheiro Maria do Rosário, Paula AM. Soares, Pestana AP Catia, Atalaia O. Sara, J.J.C., 2015. The Resilience Scale: A study in a Portuguese adult sample. In *The European Proceedings of Social & Behavioural Sciences*. portugal: a Faculty of Psychology and Educational Sciences, Research Centre of Cognitive and Behavioural Studies and Intervention.
- Portzky Michael, Wagnild Gail, Bacquer Dirk De, A.K., 2010. Psychometric evaluation of the Dutch Resilience Scale RS-nl on 3265 healthy participants: a confirmation of the association between age and resilience found with the Swedish version. *Scandinavian Journal of Caring Sciences*, 24, pp.86–93.
- Proyogi, A.S. & Agung, G.A.K., 2016. The Relationship Between Resilience and The Anxiety Level of Cancer Patients. *jurnal penelitian kesehatan suara forikes*, VII(3), pp.149–155.
- Rosenberg R. Abby, Bona Kira, Ketterl Tyler, Wharton M. Claire, Wolfe Joanne, Baker, K.S., 2017. Intimacy, Substance Use, and Communication Needs During Cancer Therapy: A Report From the “Resilience in Adolescents and Young Adults” Study. *Journal of Adolescent Health*, 60(1), pp.93–99.
- Sana et al., 2016. Assessment of psychological impact of breast cancer in women. *International Journal of Applied Pharmaceutical Sciences and Research* 2016, 1(3), pp.119–126.
- Vanderbilt-Adriance, E. & Shaw, D.S., 2009. Protective Factors and the Development of Resilience in the Context of Neighborhood Disadvantage. *Journal Abnorm Child Psychol*, 36(6), pp.887–901.
- Wagnild, G., 2003. Resilience and Successful Aging Comparison Among Low and High Income Older Adults. *JOURNAL OF GERONTOLOGICAL NURSING*, 29(12), pp.42–49.
- Wagnild, G.M. & Young, H.M., 1993. Development and psychometric evaluation of the Resilience Scale. *Journal Of Nursing Measurement*, 1(2).
- Wolff, S.N., 2007. The burden of cancer survivorship. In M. Feuerstein, ed. *Handbook of cancer survivorship*. New York: Springer, pp. 7–18.
- Woodgate, R.L., 1999. Conceptual Understanding of Resilience in the Adolescent With Cancer: Part I INDIVIDUALS. *journal of pediatric oncology nursing*, 16(1), pp.35–43.
- Yunitasari, E., 2016. *The Development of Coping-Roy's Adaptation Nursing Care Model To Increase The Resiliency of Post Radikal Hysterectomy + BSO Cervical Cancer Patients Who Received Chemotherapy*. Universitas Airlangga.
- Yusof Suzana, Nadzirah Fatin, Keterina Nina, D.R., 2016. Depressive Symptoms among Cancer Patients Undergoing Chemotherapy. *Procedia - Social and Behavioral Sciences*, 234, pp.185–192.
- Zhang, Mei-fen Zheng, Mei-chun Liu, Wei-yan Wen, Y. & Wu, Xiao-dan Liu, Q., 2015. The influence of demographics, psychological factors and self-efficacy on symptom distress in colorectal cancer patients undergoing post-surgical adjuvant chemotherapy. *European journal of oncology nursing: the official journal of European Oncology Nursing Society*, 19(1), pp.89–96.

The Influence of Age And Coping Mechanism on The Resilience of Cancer Patients Undergo Chemotherapy

ORIGINALITY REPORT

11%

SIMILARITY INDEX

7%

INTERNET SOURCES

9%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** DebraR. Hanna. "The Richness or Poverty of Nurses' Professional Selves", Research and Theory for Nursing Practice, 2017
Publication 1%
- 2** Shadiya Mohamed Saleh Baqutayan. "The Importance of Religious Orientation in Managing Stress", International Journal of Psychological Studies, 05/31/2011
Publication 1%
- 3** Woodgate, R.L.. "Conceptual understanding of resilience in the adolescent with cancer: Part I", Journal of Pediatric Oncology Nursing, 199901
Publication 1%
- 4** Mary Ann Hoffman, Robert W. Lent, Trisha L. Raque-Bogdan. "A Social Cognitive Perspective on Coping With Cancer", The Counseling Psychologist, 2012
Publication 1%
- 5** www.phdernegi.org
Internet Source 1%

6	www.ncbi.nlm.nih.gov Internet Source	1%
7	etd.fcla.edu Internet Source	<1%
8	docshare.tips Internet Source	<1%
9	www.apa.org Internet Source	<1%
10	Enas M. F. El Houby. "Framework of Computer Aided Diagnosis Systems for Cancer Classification Based on Medical Images", <i>Journal of Medical Systems</i> , 2018 Publication	<1%
11	openaccess.city.ac.uk Internet Source	<1%
12	Abby R. Rosenberg, Kira Bona, Tyler Ketterl, Claire M. Wharton, Joanne Wolfe, K. Scott Baker. "Intimacy, Substance Use, and Communication Needs During Cancer Therapy: A Report From the "Resilience in Adolescents and Young Adults" Study", <i>Journal of Adolescent Health</i> , 2017 Publication	<1%
13	Mei-fen Zhang, Mei-chun Zheng, Wei-yan Liu, Yong-shan Wen, Xiao-dan Wu, Qian-wen Liu.	<1%

"The influence of demographics, psychological factors and self-efficacy on symptom distress in colorectal cancer patients undergoing post-surgical adjuvant chemotherapy", *European Journal of Oncology Nursing*, 2015

Publication

14

media.proquest.com

Internet Source

<1%

15

Michael Portzky. "Psychometric evaluation of the Dutch Resilience Scale RS-nl on 3265 healthy participants: a confirmation of the association between age and resilience found with the Swedish version : Psychometric evaluation of RS-nl on healthy participants", *Scandinavian Journal of Caring Sciences*, 12/2010

Publication

<1%

16

www.panaxea.com

Internet Source

<1%

17

www.macrothink.org

Internet Source

<1%

18

Eshel, Yohanan, and Shaul Kimhi. "Postwar Recovery to Stress Symptoms Ratio as a Measure of Resilience, Individual Characteristics, Sense of Danger, and Age", *Journal of Loss and Trauma*, 2015.

Publication

<1%

19

journals.sagepub.com

Internet Source

<1%

20

"XXIst Symposium Neuroradiologicum",
Neuroradiology, 2018

Publication

<1%

21

ajrccm.atsjournals.org

Internet Source

<1%

22

rd.springer.com

Internet Source

<1%

23

seatvet.seameo.org

Internet Source

<1%

24

file.scirp.org

Internet Source

<1%

25

essay.utwente.nl

Internet Source

<1%

26

helda.helsinki.fi

Internet Source

<1%

27

Wei-Wen Wu, Joanne T. Chang, Shao-Yu Tsai,
Shu-Yuan Liang. "Assessing Self-concept as a
Mediator Between Anger and Resilience in
Adolescents With Cancer in Taiwan", Cancer
Nursing, 2018

Publication

<1%

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On

The Influence of Age And Coping Mechanism on The Resilience of Cancer Patients Undergo Chemotherapy

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
