

DAFTAR PUSTAKA

- Ai L, Mu S, Hu Y. (2018), Prognostic role of RDW in hematology malignancies: a systematic review and meta-analysis, primary research., *BMC, Cancer Cell International*, 18:61, <https://doi.org/10.1186/s12935-018-0558-3>
- Bennet, M. C. (2005). Immunophenotyping of acute leukemia, 98, 9-21.
- Bogar L, Molnar Z, Kenyeres P dan Tarsoly P, (2006), Sedimentation characteristic of leucocytes can predict bacteraemia in critical care patients. *J Clin Pathol* ;59:523-525. doi 10.1136/jcp.2005.033035
- Borowitz, M. J., Wood, B. L., Devidas, M., Loh, M. L., Raetz, E. A., Salzer, W. L., Larsen, E. (2015). Prognostic significance of minimal residual disease in high risk B-ALL: A report from Children's Oncology Group study AALL0232. *Blood*, 126(8), 964–971. <https://doi.org/10.1182/blood-2015-03-633685>.
- Bridgen, M. L. (1999). Clinical utility of the erythrocyte sedimentation rate. *American Family Physician*.
- Bulloch, B., Vyles, D., Gnagi, F., Hu, C., & Muenzer, J. (2016). Procalcitonin as a Marker of Bacteremia in Patients With Fever and Acute Lymphoblastic Leukemia. *Pediatric Emergency Care*, 32(9), 590–593.
- Cooper SL, B. P. (2015). Treatment of Pediatric Acute Lymphoblastic Leukemia. *Pediatr Clin North Am. National Comprehensive Cancer Network*, (62(1):), 61–73.
- Data divisi hematologi onkologi Departemen/SMF Ilmu Kesehatan Anak RSUD DR. Soetomo tahun 2013 (Data tidak dipublikasikan).
- Dahlan. (2010). Besar Sampel dan Cara Pengambilan Sampel Dalam Penelitian Kedokteran dan Kesehatan. Penerbit Salemba, 76.
- Devina T, Lubis M, Mutiara E. (2016), *Red cell distribution width and mortality in pediatric sepsis. Paediatric Indonesia.*;56(6):320-324
- Erber WN. (2010). Immunocytochemistry. In: Wendy N. Erber (ed). *Diagnostic Techniques in Hematological Malignancies*, Cambridge University Press, New York, 51-70.
- Fianza, P. I. (2009). Leukemia limfoblastik akut, 5, 1266-75.

- Forsythe A, Breland T, Majumdar S, Elkin D, Johnson D, Megason G. (2010). Gender Differences in Incidence Rates of Childhood B-Precursor Acute Lymphocytic Leukemia in Mississippi, *Journal of Pediatric Oncology Nursing*, 27(3)164-167. doi:10.1177/1043454209357919, <http://jopon.sagepub.com>
- Gabay C, Kushner I. (1999). Acute-phase proteins and other systemic responses to inflammation. *N. Engl. J. Med.* Feb 11; 340(6):448-54
- Gaynon, P. S., Desai, A. A., Bostrom, B. C., Hutchinson, R. J., Lange, B. J., Nachman, J. B., Reaman, G. K., Sather, H. N., Steinherz, P. G., Trigg, M. E., Tubergen, D. G., Uckun, F. M. (1997). Early response to therapy and outcome in childhood acute lymphoblastic leukemia: a review. *Cancer*, 80, 1717-26.
- Gil, H., Rabaud, C., Estavoyer, J., Khayat, N., Schuhmacher, H., Hoen, B., Chirouze, C. (2002). Low Serum Procalcitonin Level Accurately Predicts the Absence of Bacteremia in Adult Patients with Acute Fever. *Clinical Infectious Diseases*, 35(2), 156–161. <https://doi.org/10.1086/341023>
- Greaves, M. 2002. Childhood leukemia. *BMJ*, 324, 283-7.
- Gundy, C. M., Mostert, S., Sutaryo, Sitaresmi, M. N., & Veerman, A. J. P. (2006). Influence of Socioeconomic Status on Childhood Acute Lymphoblastic Leukemia Treatment in Indonesia. *Pediatrics*, 118(6), e1600–e1606. <https://doi.org/10.1542/peds.2005-3015>.
- Hannisdal, E., Grøttum, K. A., & Langmark, F. (1986). Erythrocyte sedimentation rate as a prognostic factor in chronic lymphocytic leukaemia. *Scandinavian Journal of Haematology*, 36(3), 253–257. <https://doi.org/10.1111/j.1600-0609.1986.tb01730.x>
- Hatzistilianou, M. (2010). Diagnostic and prognostic role of procalcitonin in infections. *TheScientificWorldJournal*, 10, 1941–1946. <https://doi.org/10.1100/tsw.2010.181>
- Hatzistilianou, M., Rekliti, A., Athanassiadou, F., & Catriu, D. (2010). Procalcitonin as an early marker of bacterial infection in neutropenic febrile children with acute lymphoblastic leukemia. *Inflammation Research*, 59(5), 339–347. <https://doi.org/10.1007/s00011-009-0100-0>
- Haque Z, Alam S, Noman M., Azhar M. (2007). Clinical Study Of Patients with Grossly Elevated Erythrocyte Sedimentation Rate, *J Medicine*; 8:64-68
- Hiroto Inaba, MD, Prof. Mel Greaves, PhD, and Charles G. Mullighan, M. (2014). NIH

- Public Access, 381(9881), 1–27. [https://doi.org/10.1016/S0140-6736\(12\)621874](https://doi.org/10.1016/S0140-6736(12)621874).
- Hunger, S. P., & Mullighan, C. G. (2015). Acute Lymphoblastic Leukemia in Children. *New England Journal of Medicine*, 373(16), 1541–1552. <https://doi.org/10.1056/nejmra1400972>.
- Hunger S P, Sung L, Howard S C. (2009). Treatment strategies and regimen graduated intensity for childhood acute lymphoblastic leukemia in low income countries : A proposal. *Pediatr Blood Cncer*;52: 559-65
- Hus, M., Halicka, D., Chocholska, S., Macheta, A., Darzynkiewicz, Z., Podhorecka, M., & Szymczyk, A. (2016). Assessment of red blood cell distribution width as a prognostic marker in chronic lymphocytic leukemia. *Oncotarget*, 7(22). <https://doi.org/10.18632/oncotarget.9055>.
- Hwan Cha C, Park C, Cha Y, Kim H, Kim D, Hoonghoon, Bae J, Jung J, Jang S, Chi H, Lee D, Cho H, (2009) Erythrocyte Sedimentation Rate Measurements by Test 1 Better Reflect Inflammation Than Do Those by The Westergreen Method in Patients With Malignancy, Autoimmune Disease, or infection. *Am J Clin Pathol* ;131:189-194. doi:10.1309/AJCPOIJ1ASTLRANIJ
- IM, W., Permono, B., IDG, U., & Ratwita, M. (2017). Luaran Pengobatan Fase Induksi Pasien Leukemia Limfoblastik Akut pada Anak di Rumah Sakit Umum Dr. Soetomo Surabaya. *Sari Pediatri*, 12(2), 128. <https://doi.org/10.14238/sp12.2.2010.128-34>
- Jandial A, Kumar S, Bhala A, Navneet S, Varma N, Varma S, (2017). Elevated Red Cell Distribution Width as Prognostic marker in Severe Sepsis: A Prospective Observational Study;21:552-62, DOI 10.4103/ijccm.IJCCM_208_17
- Julia, M. (2009). *Paediatrica Indonesiana*, 49(4), 229–233. <https://doi.org/10.14238/pi>
- Jurado R. (2001), Why Shouldn't We determine the Erythrocyte Sedimentation rate?, *Viewpoints Clinical Infectious Disease*;33:548-9. doi 1058-4838/2001/3304.0019\$03.00
- Kapoor R, Simalti A., Roy S, Agarwal P., Hematological Profile of Febrile Neutropenia in Acute Leukemia and Utility of Serum Procalcitonin Levels in Neutropenic Patients
- Kirana Kamima, Djajadisman Gatot, Sri Rezeki S. Hardinegoro (2009) Profil

- Antioksidan dan Oksidan Pasien Anak dengan Leukemia Limfoblastik Akut pada Kemoterapi Fase Induksi, Peditriacia Indonesia,
- Khalid S, Moiz B, Adil S N, Khurshid M. 2010. Retrospective review of pediatric patients with acute lymphoblastic leukemia : A single center experience. *Indian J Pathol and Microbiol*; 534 (4) : 704-10.
- Kiriu T, Yamamoto M, Nagano T, Koyama K, Katsurada M, tamura D, Nakata K, Tachihara M, Kobayashi K and Nishimura Y. 2019. Prognostic value of Red Blood Cell Distribution Width in Non small Cell Lung Cancer Treated with Anti programme Cell Death-1 Antibody. *Invivo* 33: 213-220.
- Lee, H. (2013). Procalcitonin as a biomarker of infectious diseases. *Korean Journal of Internal Medicine*, 28(3), 285–291. <https://doi.org/10.3904/kjim.2013.28.3.285>
- Lee, W. S. M., & Kim, T. M. (2010). Relation Between Red Blood Cell Distribution Width and Inflammatory Biomarkers in Rheumatoid Arthritis, *133*(April), 628–633. Retrieved from <https://www.archivesofpathology.org/doi/pdf/10.1043/1543-2165-134.4.505.c>
- Li B, You Z, Xiong X Z, Zhou Y, Wu S J, Zhou R X, Lu J and Cheng N S. (2017). Elevated red cell distribution width predict poor prognosis in hilar cholangiocarcinoma. *Oncotarget*, 8 (65), 109468-109477.
- Liu, Y., Zhang, Q., Wang, J., Wang, X., Jiang, J., Qu, Z., Fang, B. (2017). Significance of procalcitonin in bacterial infections among acute leukemia patients with post-chemotherapy agranulocytosis. *Srpski Arhiv Za Celokupno Lekarstvo*, 145(7–8), 382–386. <https://doi.org/10.2298/sarh160811066q>
- Litao M, Kamat D. (2018), Red Blood Cell Distribution Width Clinical Use Beyond Hematology, Back to Basics, *Pediatrics in Review*, April, Vol 39 No 4.
- Meisner, M. (2014). Update on Procalcitonin Measurements. Procalcitonin (PCT) as a marker for the diagnosis of sepsis and to guide antibiotic therapy. *Ann Lab Med*, 34, 263–273. <https://doi.org/10.3343/alm.2014.34.4.263>
- Mohanty J, Nagababu E, Rifkind J. (2014), Red blood cell oxidative stress impairs oxygen delivery and induces red blood cell aging, Review article, pub 28 february 2014, doi:10.3389/fphys.2014.00084
- Monig H, Marquardt, T Arendt T and Kloehn S. 2002. Limited value of elevated

erythrocyte sedimentation rate as an indicator of malignancies.

- Montagnana, M., & Danese, E. (2016). Red cell distribution width and cancer. *Annals of Translational Medicine*, 4(20), 399–399. <https://doi.org/10.21037/atm.2016.10.50>
- Neuenschwander L, Bittencourt H, Riberio A, Lucio A, Teixeira M, Teixeira J, Nobre V., Plasma levels of procalcitonin and eight additional inflammatory molecules in febrile neutropenic patients, *Clinical Science, Clinics*, 2011; 66(10):1699-1705. doi:10.1590/s1807-59322011001000006
- Ociepa, T., Sawka, E., & Panek, P. (2013). The value of concomitant measurement of white blood count, C-reactive protein and procalcitonin in predicting the course of infection in children with acute lymphoblastic leukemia and central venous catheter. *Ocena przydatności jednoczasowego oznaczania I*, (9), 634–639.
- Perlstein, T. S. (2009). NIH Public Access. *Archives of Internal Medicine*, 169(6), 588–594. <https://doi.org/10.1001/archinternmed.2009.55>. Red.
- Pui, C-H., Sandlund, J. T., Pei, D., Campana, D., Rivera, G. K., Ribeiro, R. C., Rubnitz, J. E., Razzouk, B. I., Howard, S. C., Hudson, M. M., Cheng, C., Kun, L. E., Raimondi, S. C., Behm, F. G., Downing, J. R., Relling, M. V., Evans, W. E. 2004. Improved outcome for children with acute lymphoblastic leukemia: result of total therapy study XIIIB at St Jude children's research hospital. *Blood*, 104, 2690-6.
- Pui C H, Pei D, Sandlund J T, Riberio R C, Rubnitz J E, Raimondi S C. 2010. Long term results of St Jude Total Therapy studies 11, 12, 13A, 13B, and 14 for childhood acute lymphoblastic leukemia. *Leukemia*, 24 (2): 371-82.
- Pui, C-H., Carroll, W. L., Meshinchi, S., Arceci, R. J. 2011. Biology, risk stratification, and therapy of pediatric acute leukemias: an update. *J Clin Oncol*, 29, 5551-65.
- Pulte D, Gondos A, Brenner H. 2008. Trend in 5 and 10 years survival after diagnosis with childhood hematologic malignancies in The United States, 1999-2004. *J Nat cancer Inst*; 100: 1301-1309.
- Puspitasari, D., Cahyadi, A., Larasati, M. C. S., Andarsini, M. R., Ugrasena, I. D., Permono, B. 2010. Risk factor of relapse in childhood acute lymphoblastic leukemia treated with Indonesian acute lymphoblastic leukemia protocol 2006 in Dr. Soetomo hospital. *Paediatr Indonesiana*, 51, 182 (Abstrak).

- Poernomo B, Ugrasena IG, Supriyadi, Buku Ajar Hematologi-Onkologi Anak Edisi Revisi. 2018, Bab 7 (276-285)
- Qu, Z., Fang, B., Ma, G., Jiang, J., Wang, X., Wang, J., Zhang, Q. (2017). Significance of procalcitonin in bacterial infections among acute leukemia patients with post-chemotherapy agranulocytosis. *Srpski Arhiv Za Celokupno Lekarstvo*, 145(7–8), 382–386. <https://doi.org/10.2298/SARH160811066Q>
- Rafsanjani, K. A., Fallahi, M., Kiumarsi, A., Falahati, V., & Moghaddam, S. (2018). The Association between Red Cell Distribution Width and Mortality in Pediatric Acute Lymphoblastic Leukemia. *Journal of Neoplasms*, 02(02), 1–3. <https://doi.org/10.21767/2576-3903.100011>
- Renaud T. 2011. Leukemias. In : Lankowsky P, editor Manual of Pediatric Hematology and Oncology. 5 nd Ed. New York: Elsevier. p 518-60
- Rhee, C. (2017). Using procalcitonin to guide antibiotic therapy. *Open Forum Infectious Diseases*, 4(1), 1–10. <https://doi.org/10.1093/ofid/ofw249>
- Riedl, J., Eigenbauer, E., Marosi, C., Posch, F., Pabinger, I., Reitter, E.-M., Königsbrügge, O. (2014). Red Cell Distribution Width and Other Red Blood Cell Parameters in Patients with Cancer: Association with Risk of Venous Thromboembolism and Mortality. *PLoS ONE*, 9(10), e111440. <https://doi.org/10.1371/journal.pone.0111440>
- Rocha, J. C. C., Cheng, C., Liu, W., Kishi, S., Das, S., Cook, E. H., Sandlund, J. T., Rubnitz, J., Ribeiro, R., Campana, D., Pui, C.-H., Evans, W.E., Relling, M. V. 2005. Pharmacogenetics of outcome in children with acute lymphoblastic leukemia. *Blood*, 105, 4752-8.
- Rule, J. A., Hynan, L. S., Attar, N., Sanders, C., Korzun, W. J., Lee, W. M., Hassanein, T. (2015). Procalcitonin identifies cell injury, not bacterial infection, in acute liver failure. *PLoS ONE*, 10(9), 1–13. <https://doi.org/10.1371/journal.pone.0138566>.
- Rytting M, Jabbour E, O'Brien S, Kantarjian H, Review Article Acute Lymphoblastic Leukemia in Adolescents and Young Adults(2017) Wiley online library.doi:10.1002/cncr.30624.
- Satyanarayana.L, Asthana S, Labani P. 2014. Childhood cancer incidence in India : A review of population based cancer registries. *Indian Pediatric*; 59 : 218-20.

- Seibel, N. L. 2008. Treatment of acute lymphoblastic leukemia in children and adolescent peak and pitfalls. *Hematology Am soc Hematol Educ Program*, 1, 374-80.
- Simanjorang, C., Kodim, N., & Tehuteru, E. S. (2013). Perbedaan Kesintasan 5 Tahun Pasien Leukemia Limfoblastik Akut dan Leukemia Mieloblastik Akut pada Anak di Rumah Sakit Kanker “Dharmais.” *Indonesian Journal of Cancer*, 7(1), 15–21.
- Singer, K., Subbaiah, P., Hutchinson, R., Odetola, F., & Shanley, T. P. (2011). Clinical course of sepsis in children with acute leukemia admitted to the pediatric intensive care unit. *Pediatric Critical Care Medicine*, 12(6), 649–654. <https://doi.org/10.1097/PCC.0b013e31821927f1>.
- Sousa J., Santos M, Magna L,Oliveira C.(2010), Validation of fractional model for erythrocyte sedimentation rate, MSC
- Suryawan, N., Idjradinata, P., & Reniarti, L. (2017). Hubungan Subtipe Sel Limfosit dengan Tingkat Remisi Pascakemoterapi Fase Induksi Leukemia Limfoblastik Akut. *Sari Pediatri*, 18(6), 448. <https://doi.org/10.14238/sp18.6.2017.448-52>.
- Tas F, Erturk K. 2017. Elevated erythrocyte sedimentation rate is associated with metastatic disease and worse survival in patients with cutaneous malignant melanoma. *Molecular and Clinical Oncology* 7: 1142-1146.
- Teachey, D. T., & Hunger, S. P. (2013). Predicting relapse risk in childhood acute lymphoblastic leukaemia. *British Journal of Haematology*, 162(5), 606–620. <https://doi.org/10.1111/bjh.12442>.
- Tehuteru, E. (2011). Gambaran Tingkat Remisi pada Leukemia Gambaran Tingkat Remisi pada Leukemia Limfoblastik Akut setelah Fase Induksi di Bangsal Kanker Anak RS Kanker Dharmais. *Indonesian Journal of Cancer*, 5(4), 159–162.
- Terwilliger, T., & Abdul-Hay, M. (2017). Acute lymphoblastic leukemia: a comprehensive review and 2017 update. *Blood Cancer Journal*, 7(6), e577. <https://doi.org/10.1038/bcj.2017.53>.
- Tschan, C., Moricke, A., Stanulla, M., Koehler, R., Goudeva, L., Bartram, C. R., Schrappe, M. (2011). Low platelet counts after induction therapy for childhood acute lymphoblastic leukemia are strongly associated with poor early response to treatment as measured by minimal residual disease and are prognostic for treatment

outcome. *Haematologica*, 97(3), 402-409.

<https://doi.org/10.3324/haeamtol.2011.045229>.

Ugrasena, G. (2016). Penyakit Leukemia Akut Banyak Jangkiti Anak-anak.

Unit Koordinasi Kerja Hematologi-onkologi Ikatan Dokter Anak Indonesia (UKK Hematologi-Onkologi IDAI) . 2013. Panduan Protokol Leukemia Limfoblastik Akut Indonesia 2018 (Indonesian ALL-2018 Protocol).

Widiaskara, I. M., Permono, B., Ugrasena, I. D. G., Ratwita, M. 2010. Luaran pengobatan fase induksi pasien leukemia limfoblastik akut pada anak di rumah sakit umum Dr. Soetomo Surabaya. *Sari Pediatri*, 12, 128-34.

Widijanti A, 2019. Pengukuran prokalsitonin serum dengan Test Cepat FIA untuk deteksi Sepsis, Materi PK Update Malang, FK Unbraw.

Williams, M. D., Braun, L. A., Cooper, L. M., Johnston, J., Weiss, R. V, Qualy, R. L., & Linde-Zwirble, W. (2004). Hospitalized cancer patients with severe sepsis: analysis of incidence, mortality, and associated costs of care. *Crit Care*, 8(5), 291–298. <https://doi.org/10.1186/cc2893>.

Yakin, Rahimul; Syarif, Syahrizal; Tehuteru, E. S. (2017). Perbandingan Kesintasan pada Anak Leukemia Limfoblastik akut antara Protokol Pengobatan 2006 dan 2013, *11*(3), 111–117.

Yousuf, M., Akhter, J., Al-Khairiy, K., Al-Saadon, M. A., & Bin-Salih, S. (2010). Extremely elevated erythrocyte sedimentation rate: Etiology at a tertiary care center in Saudi Arabia. *Saudi Medical Journal*, 31(11), 1227–1231.