


PROCEEDING



international seminar

STRATEGY TO MANAGE BIO-ECO-HEALTH SYSTEM FOR STABILIZING ANIMAL HEALTH & PRODUCTIVITY TO SUPPORT PUBLIC HEALTH



Surabaya-Indonesia, 19-20 June 2012
JW Marriott Hotel Surabaya

EDITORS:

Michael P. Ward (Australia)

Faouzi Kechrid (Africa)

Montip Gettayacamin (Thailand)

Fedik Abdul Rantam (Indonesia)

Suzanita Utama (Indonesia)

FACULTY OF VETERINARY MEDICINE - UNIVERSITAS AIRLANGGA
I-MHERE SUB-COMPONENT B.2.C PERFORMANCE BASED CONTRACT

PRODUCTION OF WHOLE SERUM PMSG (PREGNANT MARE SERUM GONADOTROPIN) WITH SEPADEX OF PREGNANT LOCAL MARE SERUM TO IMPROVE GESTATION AND NUMBER OF FAT TAILED SHEEP STRAIN IN SAPUDI ISLAND.....	27
<i>Herry Agoes Hermadi</i>	
EXPRESSION OF TOLL LIKE RECEPTOR ON RABBITS IMMUNIZED WITH ANTIGENIC PROTEINS OF SARCOPTES SCABIEI VAR.CAPRAE.....	32
<i>Nunuk Dyah Retno Lastuti</i>	
THE EFFECT OF THORACO-VAGOTOMIZED CALVES ON RUMEN DEVELOPMENT BY PGP 9.5 IMMUNOHISTOCHEMISTRY	35
<i>R. Harry Soehartono and Dwi Dian Vitasari</i>	
THE EFFECT OF BACTERIOCIN TO REDUCE THE NUMBER OF ESCHERICHIA COLI ISOLATED FROM BEEF SOULD AT ABATTOIR.....	39
<i>Nenny Harijani, Luviana Kristianingtyas, Hario Puntodewo, Soelih Estoepangestie</i>	
THE EFFECT OF BACTERIOCIN AS AN ANTIBACTERIA ON THE TOTAL BACTERIAL COUNT OF CHICKEN MEAT STORED AT 4° C.....	43
<i>Nenny Harijani, Dara Recardsari Casarus, Romziah Sidik</i>	
GROWTH ASPECTS OF BROILER AT AGE CONSTANT VS WEIGHT CONSTANT	48
<i>Andoyo Supriyantono</i>	
ULTRASONOGRAPHY INTERPRETATION OF LIVER ABNORMALITIES IN THE DOGS	52
<i>Deni Noviana, Budhy Jasa Widyananta, I Wayan Widi Parnayoga</i>	
SENSITIVITY ANALYSIS OF LAYER CHICKEN FARMS IN SUB-DISTRICT KEDUNGPRING LAMONGAN.....	56
<i>Sunaryo Hadi Warsito</i>	
PIG HUSBANDRY AND MANAGEMENT ADOPTED BY FARMERS AND THEIR IMPACTS TO CSF TRANSMISSION IN WEST TIMOR, INDONESIA	60
<i>Petrus Malo Bulu, Ian Robertson, Jenny-Ann Toribio, Maria Geong</i>	
ANTIBACTERIAL SUSCEPTIBILITY OF BACILLUS SUBTILIS ISOLATED FROM SOIL AND FISHPOND SEDIMENT.....	64
<i>Erni Rosilawati Sabar Iman, Lina Susanti, Sri Subekti</i>	
HAEMOGREGARINE CASE IN PYTHON SNAKE.....	68
<i>Mufasirin</i>	
HISTOPATHOLOGY OF HEPATOCYTE NUCLEUS DEGENERATION EXPOSED BY CURCUMA AERUGINOSA	70
<i>Eka Pramyrtha Hestianah</i>	
CORRELATION ANALYSIS MODEL OF HEMATOLOGY EXAMINATION, INFLAMMATORY CELLS AND BLOOD CHEMICAL PROFILE OF KAMBING KACANG AT DESA MOJOSARIREJO DRIYOREJO GRESIK.....	73
<i>Hana Eliyani, Soeharsono, Retno Bijanti</i>	
PREVALENCE OF OBESITY AND RISK FACTORS IN DOGS IN SURABAYA.....	76
<i>Nusdianto Triakoso</i>	



VETERINARY ANTIBIOTICS IN ANIMAL PRODUCTION AND THE ENVIRONMENT.....	80
<i>Saleha A.A.</i>	
MICROBIOLOGICAL ANALYSIS OF DRINKING WATER AND SOYBEAN MILK.....	83
<i>Lucia R.W. Muslimin and FikaYulizaPurba</i>	
THE EFFECTS OF HYPERBARIC OXYGEN ON THE NUMBER OF EOSINOPHILS AND THE PICTURES OF SPLEEN WHITE PULP DIAMETERS IN WHITE RATS GIVEN HEAVY SWIMMING EXERCISES.....	86
<i>Setianingsih, H.</i>	
CORRELATION OF SERUM ALP ACTIVITY WITH THE HEALING PROCESS OF FEMORAL FRACTURES IN RATS USED CISSUS QUADRANGULARIS EXTRACT AS THERAPY.....	90
<i>Ira Sari Yudaniyanti, Lianny Nangoi, Julien Soepraptini</i>	
IMMUNOHISTOCHEMICAL ANALYSIS ON THE DISTRIBUTION OF ADENOHYPHYSIAL CELLS IN THE PITUITARY PARS DISTALIS OF THE OSTRICH (STRUTHIO CAMELUS).....	94
<i>Dwi Kesuma Sari, Lucia Muslimin, Fika Yuliza Purba, I Ketut Mudite Adnyane, Kazuhide Adachi, Yasuhiro Tsukamoto</i>	
CORRELATION BETWEEN DURATION TIMES OF CRYOPROTECTANT TOWARD MICE EMBRYO DEVELOPMENT.....	96
<i>Bambang Poernomo S., Soeharsono, Trianto Nur Abdullah</i>	
DEVELOPMENT OF THE FIVE ELEMENTS MODEL ON INTERACTION LIVER AND KIDNEY FUNCTION THROUGH BLOOD AS MEDIATOR USING EQUALLY PARAMETER.....	100
<i>Soeharsono, RTS Adikara, E. Widjajanto, Bambang Poernomo S.</i>	
CHARACTERIZATION OF IMMUNOGLOBULIN Y AGAINST SOLUBLE PROTEIN OF <i>TOXOPLASMA GONDII</i>	104
<i>Lucia Tri Suwanti, Marek Yohana Kurniabudhi, Hani Plumeriastuti, Suwarno, Fedik Abdul Rantam</i>	
FROZEN SEMEN OF MERINO RAM PRODUCTION IN CENTRAL ARTIFICIAL INSEMINATION DISTRICT OF FACULTY OF VETERINARY MEDICINE UNIVERSITAS AIRLANGGA FOR IMPROVEMENT POPULATION OF SHEEP IN EAST JAVA.....	107
<i>Abdul Samik, Herry Agoes Hermadi, Sri Pantja Madyawati, Trilas Sardjito</i>	
CHARACTERIZATION OF <i>BRUCELLA ABORTUS</i> VACCINE STRAIN S-19 AND LOCAL ISOLATE WITH CONVENTIONAL BACTERIOLOGY METHODS AND MULTIPLEX POLYMERASE CHAINS REACTION (PCR).....	110
<i>Nunung Aji Wibowo, Didik Handijatno, Ratih Ratnasari</i>	
THE EFFECT OF EGGS YOLK SKIM AND EGG YOLK TRIS ON MOTILITY AND VIABILITY OF MERINO SHEEP SEMEN POST-THAWING.....	115
<i>Yossi Aris Munandar, Abdul Samik, Rudy Sukamto, Wurlina Meles</i>	
ARTIFICIAL INSEMINATION PROGRAM FOR BEEF CATTLE IN MADURA ISLAND "TARGETS, REALIZATION AND PROBLEMS".....	118
<i>Mas'ud Hariadi</i>	



See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/327529707>

Histopathology of Hepatocyte Nucleus Degeneration Exposed by Curcuma Aeruginosa

Conference Paper · June 2012

CITATIONS

0

READ

1

1 author:



Eka Pramytha Hestianah

Airlangga University

12 PUBLICATIONS 4 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



retno_biotek@yahoo.com [View project](#)

HISTOPATHOLOGY OF HEPATOCYTE NUCLEUS DEGENERATION EXPOSED BY CURCUMA AERUGINOSA

Eka Pramytha Hestianah

Department of Veterinary Anatomy
Faculty of Veterinary Medicine - Universitas Airlangga
E-mail: eka_1964@yahoo.co.id

ABSTRACT

This research is aimed at finding out the effect of *Curcuma Aeruginosa* given to the histopathology hepatocyte of 40 male mice, weight 25-30 g. The mice are divided into 5 treatment groups namely 1 control group, 2 infused groups and 2 squeezed groups. *Curcuma Aeruginosa* is given once a day and on day 11, the livers of the mice are taken and made into histopathology preparation with HE staining. The data taken are analyzed by manovafactorial and anova one way. The result shows the occurrence of hepatocyte nucleus degeneration, the highest means of karyopicnosis and karyorhexis on infused group of 10 mg/25 g body weight, are 10.22% and 49.6% ($\alpha = 0.05$). The highest karyolysis on infused and squeezed groups of 10 mg/25 g body weight, is 45.86% and 45.70% ($\alpha = 0.05$). The conclusion is that infused and squeezed *Curcuma Aeruginosa* given orally to male mice has caused the hepatocyte nucleus degeneration.

Keywords: Hepatocyte, nucleus degeneration, curcuma aeruginosa, mice

INTRODUCTION

Curcuma Aeruginosa is a useful herb which has been widely used as the appetite booster, helminthes medicine, etc (Santoso, 1989; Jantanet.al., 2003). It is also often used as traditional medicine in rural area in the form of infused (boiled) and squeezed. The people who use this herb sometimes do not pay attention on the dosage and how long they should consume the herb therefore they do not realize the side effect that might happen.

The WHO states that inconsistent and toxic compound found in herb could give toxic effect (Handa, 1995; Li et.al., 2008). The two reported toxics are, first, it is caused by the intrinsic factor of the herb itself related to over dosage, allergic responses such as vomiting, inflammation and diarrhea. Second, it is caused by extrinsic factor related to the mistake during preparation making such as mistakenly identifying the herbs, failure in the process of medicine making, and the occurrence of contamination (Calixto, 2000).

MATERIALS AND METHODS

The samples used were 40 male mice Balb C weight 25-30 g which were divided into 5 treatment groups namely P0 (control, given CMCNa and aquadest), P1 (consists of 2 treatment groups given 2 different dosages, infused *Curcuma Aeruginosa*), and P2 (consists of 2 treatment groups given 2 different dosages, squeezed *Curcuma Aeruginosa*). *Curcuma Aeruginosa* rhizome taken from Jombang was made into infused and squeezed and then dried using Freeze Dry method. Dosages given were 10 mg/25 g and 15 mg/25 g body weight in accordance with the group orally per day for 10 days.

On day eleventh, the livers of the male mice were taken to be made into histopathological study by using HE staining. Histopathological examination was done by counting the amount percentage of the hepatocytes undergone nucleus degeneration (karyopicnosis, karyorhexis, and karyolysis) from 1500 hepatocytes (Ekicinet.al., 2005). The data gained from the analysis were analyzed by using manovafactorial and then continued with anova one way.

RESULT AND DISCUSSION

Necrosis is the process of cell disfunction which is marked with the nucleus degeneration during the early stage of karyopicnosis, where the nucleus clumped or solidified and looked brighter. Next, the nucleus undergoes karyorhexis, it is when the nucleus is broken into two or more and then it undergoes karyolysis. It is when the nucleus dissolves or disappears and loses its form (McGavin et al., 2007).

The result of histopathology preparation toward hepatocyte nucleus degeneration of each group is as follows:

Table 1. Mean percentage of hepatocytes nucleus degeneration and SD as the result of Curcuma Aeruginosa exposition in the form of infused and squeezed.

Treatment groups	Dosage	Karyopicnosis	Karyorhexis	Karyolysis
Control	0 mg	0.39 ± 0.18 ^a	0.93 ± 0.27 ^a	4.04 ± 0.70 ^a
Infused	10 mg	7.93 ± 0.92 ^{c,d}	30.59 ± 1.33 ^d	45.86 ± 2.12 ^c
	15 mg	10.22 ± 1.32 ^c	49.66 ± 2.32 ^e	29.36 ± 2.39 ^b
Squeezed	10 mg	5.78 ± 0.90 ^b	16.03 ± 1.54 ^b	45.70 ± 2.77 ^c
	15 mg	5.93 ± 0.55 ^{b,c}	21.05 ± 1.36 ^c	53.05 ± 2.56 ^d

Superscript: Different letter on each variable shows the significant difference of each group on $\alpha = 0.05$.

The result shows the occurrence of necrosis or hepatocyte nucleus degeneration as a result of Curcuma Aeruginosa treatment, it proves that there is a real difference between the control and treatment group ($\alpha = 0.05$) as seen in table 1. Chan's research result proved that there is a disadvantage effect of using traditional medicine which often endangers people. Other research shows that most of herbal plant have toxic effect to specific organ and it is little known about the chronic toxic effect of a long term use, dosage used, and the lateness of herbs use termination of traditional medicine which brings about acute and chronic intoxication (Ezejioret al., 2008; Sunita et al., 2008; Dewi & Saraswati, 2009).

Infused shows a higher hepatocyte nucleus degeneration mean value than squeezed. It is caused by the heating on infused stock. The heating activates toxic substances contained in Curcuma Aeruginosa therefore those toxics come out of the Curcuma Aeruginosa cell, and the active toxic cell is able to go through the hepatocyte's membrane and the nucleus membrane. It causes chromatin to clump inside nucleus and it is related to RNA synthesis decrease. If this condition continues to happen, the cell will undergo nucleus degeneration called karyopicnosis (Mc Gavin et al., 2007).

CONCLUSION

The infused and squeezed Curcuma Aeruginosa given orally to male mice has caused hepatocyte nucleus degeneration and the 10 mg squeezed Curcuma aeruginosa per 25 g body weight was safe to be used in mice.

REFERENCES

- Calixto, JB., 2000. Efficacy, Safety, Quality Control, Marketing and Regulatory Guidelines for Herbal Medicines (Phytotherapeutic Agents). Brazilian, Med Biol Res, 33: 179–189.
- Chan, K., 2003. Review: Some Aspect of Toxic Contaminants in Herbal Medicine. Chemosphere (52): 1361–1371.



- Dewi, UK dan Saraswati, TR., 2009. Efek Rebusan Daun Tapak Dara pada Dosis dan Frekuensi yang Berbeda Terhadap Kerusakan dan Akumulasi Glikogen pada Hepar Mencit (*Mus musculus*). *Bioma*, 11(1): 1–5.
- Ekici, Y., Tez, M., Sökmenster, C., Baskan, S., 2005. Mechanism of Cell Death After Extensive Liver Resection: Apoptosis or Necrosis? *Internet Pathol* 3(2): 1528–8307.
- Ezejiolor, NA., Maduagwunan., Onyiaorah, VI., Hussaini, DC and Orisakwe, OE., 2008. Multiple Organ Toxicity of a Nigerian Herbal Supplement (U & D Sweet Bitter) in Male Albino Rats. *Pak. J. Pharm. Sci*, 21(4): 426–429.
- Handa, SS., 1995. Quality Control and Standardization of Herbal Raw Materials and Traditional Remedies. *Pharmatimes*, 13–21.
- Jantan, I., Yassin, M., Chin, CB., Chen., CL and Sim, NL., 2003. Antifungal Activity of The Essential Oil of Nine Zingiberaceae Species. *Pharmaceutical Biol*, 41 (5): 392–397).
- Li, S., Han, Q., Qiao, C., Song, J., Cheng, CL., and Xu, H., 2008. Chemical Marker for The Quality Control of Herbal Medicines: An Overview. *Chinese Medicine, Hongkong SAR, PR China*, 1–25.
- McGavin, MD and Zachary, JF., 2007. *Pathology Basic of Veterinary Disease*. 4th Ed. Mosby Elsevier, St. Louis, Missouri, 3–34.
- Santoso, SO., 1989. Penggunaan Obat Tradisional Secara Rasional. *Cermin Dunia Kedokteran*, 59: 3–6.
- Sunita, P., Rathore, KS., Sisodia, SS., Meenakshi, B., Nema, RK., 2008. An Overview on Herbal Toxicity. *Herbal Toxicol* (2): 35–37.

