

**DAFTAR PUSTAKA**

1. Weiss SL, Fitzgerald JC, Pappachan J, Wheeler D, Jaramillo-Bustamante JC, Salloo A, et al. Global Epidemiology of Pediatric Severe Sepsis: The Sepsis Prevalence, Outcomes, and Therapies Study. 2015 May 15; 191(10): p. 1147-1157.
2. Priyatiningsih LAPA. Karakteristik sepsis di pediatric intensive care unit RSUPN dr. Cipto Mangunkusumo. 2016.
3. Supit P, Mandei JM, Rampengan NH. Profil anak dengan sepsis dan syok sepsis yang dilakukan kultur darah periode Januari 2010 – Juni 2015 di RSUP Prof. Dr. R. D. Kandou Manado. 2015; 4(1).
4. Chong SL, Ong GY, Venkataraman A, Chan YH. The Golden Hours in Paediatric Septic Shock—Current Updates and Recommendations. 2014; 43: p. 267-74.
5. Davis AL, Carcillo JA, Aneja RK, Deymann AJ, Lin JC, Nguyen TC. American College of Critical Care Medicine Clinical Practice Parameters for Hemodynamic Support of Pediatric and Neonatal Septic Shock. 2017; 45(6): p. 1061-1092.
6. Carcillo JA, Han YY, Kisson N. Sepsis guidelines and the global pediatric sepsis initiative: implications for treatment. 2008; 5(4): p. 391-394.
7. Maitland K, Kiguli S, Opoka RO, Engoru C, Olupot-Olupot P, Akech SO, et al. Mortality after Fluid Bolus in African Children with Severe Infection. 2011; 364: p. 2483-2495.
8. Gelbart B, Glassford NJ, Bellomo R. Fluid Bolus Therapy-Based Resuscitation for Severe Sepsis in Hospitalized Children: A Systematic Review. 2015; 16(8): p. 297-307.
9. Frazier WJ, Muszynsk JA. Fluid resuscitation in pediatric sepsis: lack of data versus lack of equipoise. 2015; 16(8): p. 789-790.
10. EL-Nawawy AA, Abdelmohsen AM, Hassouna HM. Role of echocardiography in reducing shock reversal time in pediatric septic shock: a randomized controlled trial. Sociedade Brasileira de Pediatria. 2017 February 20; 528: p. 1-9.
11. Ranjit S, Aram G, Kisson N, Ali K, Natraj R, Shresti S, et al. Multimodal Monitoring for Hemodynamic Categorization and Management of Pediatric Septic Shock: A Pilot Observational Study. 2013; 15(1): p. 17-26.

12. Seif D, Perera P, Mailhot T, Riley D, Mandavia D. Bedside Ultrasound in Resuscitation and the Rapid Ultrasound in Shock Protocol. 2012.
13. Ranjit S, Kissoon N. Bedside echocardiography is useful in assessing children with fluid and inotrope resistant septic shock. 2013; 17(4).
14. Saraswati DD, Pudjiadi AH, Djer MM, Supriyatno B, Syarif DR, Kurniati N. Faktor resiko yang berperan pada mortalitas sepsis. 2014 Februari; 15(5): p. 281-8.
15. Rady HI, Mohamed SA, Mohssen NA, ElBaz M. Application of Different Scoring Systems and Their Value in Pediatric Scoring Systems. 2014; 62: p. 59-64.
16. Singer M, Deutschman CS, Seymour CW, Shankar-Hari M, Annane D, Bauer M, et al. The Third International Consensus Definitions for Sepsis and Septic Shock (Sepsis-3). 2016; 315(8): p. 801-810.
17. Gogia P, Koreti S, Patel GS. SOFA (Sequential Organ Failure Assessment) and PELOD (Pediatric Logistic Organ Dysfunction). 2015 July; 3(4a): p. 1645-1648.
18. Plunkett A, Tong J. Sepsis in children. 2015; 350(h3017).
19. Gan H, Cannesson M, Chandler JR, Ansermino JM. Predicting fluid responsiveness in children: a systematic review. 2013; 117(6).
20. Lukito V, Djer MM, Pudjiadi AH, Munasir Z. The role of passive leg raising to predict fluid responsiveness in pediatric intensive care unit patients. *Pediatric Critical Care Medicine*. 2012 May; 13(3): p. 155-160.
21. Neto EPdS, Grousson S, Duflo F, Ducreux C, Joly H, Convert J, et al. Predicting fluid responsiveness in mechanically ventilated children under general anaesthesia using dynamic parameters and transthoracic echocardiography. *British Journal of Anaesthesia*. 2011; 106(6): p. 856-864.
22. Osborn TM. Severe Sepsis and Septic Shock Trials (ProCESS, ARISE, ProMISe) What is Optimal Resuscitation? 2017; 33: p. 323-344.
23. Jones AE, Shapiro NI, Trzeciak S, Arnold RC, Claremont HA, Kline JA. Lactate Clearance vs Central Venous Oxygen Saturation as Goals of Early Sepsis Therapy. 2010; 303(8): p. 739-746.
24. Sankar J, Das R, Jain A, Dewangan S, Khilnani P, Yadav D, et al. Prevalence and outcome of diastolic dysfunction in children with fluid refractory septic shock--a prospective observational study. 2014; 15(9): p. 370-378.

25. Hadinegoro SRS, Chairulfatah A, Latief A, Pudjiadi AH, Malisie RF, Alam A. *Konsensus Diagnosis dan Tatalaksana Sepsis pada Anak Jakarta: Ikatan Dokter Anak Indonesia*; 2016.
26. Bentzer P, Griesdale DE, Boyd J, MacLean K, Sirounis D, Ayas NT. Will This Hemodynamically Unstable Patient Respond to a Bolus of Intravenous Fluids? *Journal of American Medical Association*. 2016; 316(12): p. 1298-1309.
27. Maitland K, Kiguli S, Opoka RO, Engoru C, Olupot-Olupot P, Akech SO. Mortality after Fluid Bolus in African Children with Severe Infection. *The New England Journal of Medicine*. 2011; 364(26): p. 2483-2495.
28. Ventura AMC, Shieh HH, Bousso A, Góes PF, Fernandes IdCFO, Souza DCd, et al. Double-Blind Prospective Randomized Controlled Trial of Dopamine Versus Epinephrine as First-Line Vasoactive Drugs in Pediatric Septic Shock. *Critical Care Medicine*. 2015; 43(11): p. 2292–2302.
29. Deep A, Goonasekera CDA, Wang Y, Brierley J. Evolution of haemodynamics and outcome of fluid-refractory septic shock in children. *Intensive Care Medicine*. 2013; 39: p. 1602–1609.
30. Gaspar HA, Morhy SS, Lianza AC, Carvalho WBd, Andrade JL, Prado RRd, et al. Focused cardiac ultrasound: A training course for pediatric intensivists and emergency physicians. *BMC Medical Education*. 2014; 14(25).
31. Gaspar HA, SaadyMorhy S. The Role of Focused Echocardiography in Pediatric Intensive Care: A Critical Appraisal. *BioMed Research International*. 2015.
32. Ünlüer EE, Karagöz A, Akoğlu H, Bayata S. Visual Estimation of Bedside Echocardiographic Ejection Fraction by Emergency Physicians. 2014; XV(2): p. 221-226.
33. Cecconi M, Backer DD, Antonelli M, Beale R, Bakker J, Hofer C, et al. Consensus on circulatory shock and hemodynamic monitoring. Task force of the european society of intensive care medicine. *Intensive Care Medicine*. 2014; 40: p. 1795-1815.
34. O. Raux\* ASRFGMARSBKLXCd. Intraoperative use of transoesophageal Doppler to predict response to volume expansion in infants and neonates. *British Journal of Anaesthesia*. 2011; 108 (1): 100–7(1): p. 100-107.
35. Kutty S, Li L, Hasan R, Peng Q, Rangamani S, Danford DA. Systemic Venous Diameters, Collapsibility Indices, and Right Atrial Measurements in Normal Pediatric Subjects. 2014; 27(2).

36. Amoozgar H, Zare K, Ajami G, Borzoe M, Abtahi S. Estimation of Right Atrial Pressure from the Inspiratory Collapse of the Inferior Vena cava in Pediatric Patients. 2010; 20(2): p. 206-210.
37. Otto CM, Schwaegler RG, Freeman RV. Left and Right Ventricular Systolic Function. In *Echocardiography Review Guide*. Philadelphia: Saunders; 2016.
38. Goodman A, Perera P, Mailhot T, Mandavia D. The role of bedside ultrasound in the diagnosis of pericardial effusion and cardiac tamponade. 2012; 5(1): p. 72-75.
39. Perera P, Mailhot T, Riley D, Mandavia D. The RUSH Exam: Rapid Ultrasound in SHock in The Evaluation of Critically Ill. 2010; 28: p. 29-56.
40. Kawasaki T. Update on pediatric sepsis: a review. 2017; 5(47).
41. Sankar J, Das RR, Jain A, Dewangan S, Khilnani P, Yadav D, et al. Prevalence and Outcome of Diastolic Dysfunction in Children With Fluid Refractory Septic Shock—A Prospective Observational Study. 2014; 15(9).
42. Raj S, Killinger JS, Gonzalez JA, Lopez L. Myocardial Dysfunction in Pediatric Septic Shock. 2014; 164: p. 72-77.
43. Priyatiningih DR LAPA. Karakteristik sepsis di pediatric intensive care unit RSUPN dr. Cipto Mangunkusumo. 2016.
44. Vincent JL, Rhodes A, Perel A, Martin GS, Rocca GD, Vallet B, et al. Clinical review: Update on hemodynamic monitoring - a consensus of 16. *Critical Care*. 2011; 15(229): p. 1-8.
45. Chaiyakulsil C, Anantasit N, Katanyuwong P. Comparison of Three Non Invasive Hemodynamic Monitoring Methods in Critically Ill Pediatric Patients. *PLOS One*. 2017; 13(6): p. 1-12.
46. Malik V, Subramanian1 A, Chauhan S, Hote M. Correlation of Electric Cardiometry and Continuous Thermodilution Cardiac Output Monitoring Systems. *World Journal of Cardiovascular Surgery*. 2014; 4(1): p. 101-108.