

DAFTAR PUSTAKA

- A. T. Vu, V. M. (2003). Alternative T1-weighted In-phase/Out-of-phase Imaging For Body MRI at 3T. 1326.
- Ahmet K. Poyraz, M. R. (2012). Diffusion-Weighted MRI of Fatty Liver. *JOURNAL OF MAGNETIC RESONANCE IMAGING*, 1108–1111.
- Andrea Agostini, M. F. (2016). Magnetic Resonance Imaging of the Liver (Including Biliary Contrast Agents) Part 2: Protocols for Liver Magnetic Resonance Imaging and Characterization of Common Focal Liver Lesions. *Semin Roentgenol*, 317–333.
- António P Matos, F. V. (2015). Focal liver lesions: Practical magnetic resonance imaging approach. *World Journal Hepatology*, 1987-2008.
- B. M. Araujo, F. R. (2015). Fat-containing lesion in the liver - What can it be? *European Society of Radiology*, 1-21.
- Barry E. Kenneally, C. J. (2015). Utility of opposed-phase magnetic resonance imaging in differentiating sarcoma from benign bone lesions. *Journal of Bone Oncology*, 110–114.
- Bernd Hamm, G. P. (2010). *MR Imaging of the Abdomen and Pelvis*. Jerman: Thieme Medical Publishers.
- C. Santos Montón, S. d. (2014). Fat-containing lesions of the liver: What do we need to know about them? *European Society of Radiology*, 10.
- Ceyla Basaran, M. K. (2005). Fat-Containing Lesions of the Liver: Cross-Sectional Imaging Findings with Emphasis on MRI. *American Journal of Roentgenology*, 1103–1110.
- Cieszanowski, A. (2006, Agustus 1). MR imaging ensures clarity in focal liver lesions. *Diagnostic Imaging*, pp. 1-5.
- Daniele Marin, A. F. (2009). Imaging Approach for Evaluation of Focal Liver Lesions. *CLINICAL GASTROENTEROLOGY AND HEPATOLOGY*, 624-634.
- Daniella Braz Parente, J. A. (2018). Fat-containing liver lesions: a pictorial review. *Radiol Bras*, 52-57.

- Eric K Outwater, R. B. (1998). Detection of Lipid in Abdominal Tissues with Opposed-Phase Gradient-Echo Images at 1.5 T: Techniques and Diagnostic Importance. *RadioGraphics*, 1465-1480.
- Fiona Hughes Cassidy, T. Y. (2009). Fatty Liver Disease: MR Imaging Techniques for the Detection and Quantification of Liver Steatosis. *RadioGraphics RSNA Journal*, 231–260.
- Henrique Donato, M. F.-A. (2017). Liver MRI: From basic protocol to advanced techniques. *European Journal of Radiology*, 30–39.
- Jorge A. Marrero, M. J. (2014). ACG Clinical Guideline: The Diagnosis and Management of Focal Liver Lesions. *The American Journal of GASTROENTEROLOGY*, 1-20.
- Kennet L. Bontrager, J. P. (2005). *Textbook of Radiographic Positioning and Related Anatomy*. Elsevier Mosby 6 Edition.
- Kessel, C. S. (2012). Optimization of imaging and treatment of patients with focal liver lesions. *Thesis by Bayer Healthcare Nederland*, 10-17.
- Lan N Vu, J. N. (2018). Basic MRI for the liver oncologists and surgeons. *Journal of Hepatocellular Carcinoma*, 37–50.
- Leupold, M. M. (2012). Gradient Echo Imaging. *JOURNAL OF MAGNETIC RESONANCE IMAGING*, 1274–1289.
- Lumongga, F. (2008). *Struktur Liver*. Medan: USU Repository.
- M. A. A. Vieira, M. B. (2012). Imaging of focal liver lesions. *European Society of Radiology*, 1-26.
- Ma, J. (2004). Breath-Hold Water and Fat Imaging Using a Dual-Echo Two-Point Dixon Technique With an Efficient and Robust Phase-Correction Algorithm. *Magnetic Resonance in Medicine*, 415-419.
- Michael A. Fischer, O. F. (2013). Two versus three-dimensional dual gradient-echo MRI of the liver: a technical comparison. *European Society of Radiology*, 408-416.
- Muqmiroh, L. (2015, November 16). *Anatomi Organ Abdomen*. Surabaya.
- Naglaa L. Debees, M. F. (2016). Assessment of hepatic focal lesions on top of cirrhotic liver using dynamic and diffusion weighted magnetic resonance imaging. *The Egyptian Journal of Radiology and Nuclear Medicine*, 1221–1230.

- Numminen, K. (2004). *MR Imaging of the Liver: Studies on the Detection and Characterization of Focal Liver Lesions and Liver Cirrhosis*. Finland: Medical Imaging Center and Transplantation and Liver Surgery Clinic University of Helsinki.
- Ph. Soyer, Y. R. (1997). T1-weighted spoiled gradient-echo MR imaging of focal hepatic lesion: comparison of in-phase vs opposed-phase pulse sequence. *European Radiology*, 1048–1053.
- R.C, O. (2016). Radiological anatomy of the liver. *Journal of Medicine and Medical Sciences*, 72-78.
- Sajal S. Pokharel, K. J. (2013). Current MR Imaging Lipid Detection Techniques for Diagnosis of Lesions in the Abdomen and Pelvis. *RSNA*, 681-702.
- Sibulesky, L. (2013). Normal liver anatomy. *Clinical Liver Disease*, 1-3.
- Sigit Wijokongko, J. A. (2016). *Protokol Radiologi CT Scan dan MRI*. Magelang: Inti Medika Pustaka.
- Srinivasa R. Prasad, H. W. (2005). Fat-containing Lesions of the Liver: RadiologicPathologic Correlation. *RadioGraphics*, 321-331.
- Sudhakar K. Venkatesh, V. C. (2014). Liver Masses: A Clinical, Radiological and Pathological Perspective For: Perspectives in Clinical Gastroenterology and Hepatology. *Clin Gastroenterol Hepatol*, 1414–1429.
- Vasco Here´dia, M. R.-H. (2011). Comparison of a Single Shot T1-Weighted In- and Out-of-Phase Magnetization Prepared Gradient Recalled Echo With a Standard Two-Dimensional Gradient Recalled Echo: Preliminary Findings. *JOURNAL OF MAGNETIC RESONANCE IMAGING*, 1482–1490.
- W.K. Erly, E. O. (2006). The Utility of In-Phase/Opposed-Phase Imaging in Differentiating Malignancy from Acute Benign Compression Fractures of the Spine. *American Journal of Neuroradiology*, 1183-1188.
- Westbrook, C. (2014). *Handbook of MRI Technique Fourth Edition*. UK: Wiley Blackwell.
- Wolfgang Schima, C. K.-S. (2012). Non-invasive diagnosis of focal liver lesions: an individualized approach. *Cancer Imaging*, 365-372.
- Yasemin Bilgili, Z. F. (2006). Focal liver lesions evaluated by MR imaging. *Diagnostic and Interventional Radiology*, 129-135.