

DAFTAR PUSTAKA

- Alfaro-Acha, A., Snih, S. A., Raji, M. A., Kuo, Y. F., Markides, K. S., & Ottenbacher, K. J. (2006). Handgrip strength and cognitive decline in older Mexican Americans. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 61(8), 859-865.
- Bainbridge, D., Seow, H., Sussman, J., Pond, G., Martelli-Reid, L., Herbert, C. and Evans, W., 2011. Multidisciplinary health care professionals' perceptions of the use and utility of a symptom assessment system for oncology patients. *Journal of Oncology Practice*, 7(1), pp.19-23.
- Chen, L.K., Lee, W.J., Peng, L.N., Liu, L.K., Arai, H., Akishita, M. and for Sarcopenia, A.W.G., 2016. Recent advances in sarcopenia research in Asia: 2016 update from the Asian Working Group for Sarcopenia. *Journal of the American Medical Directors Association*, 17(8), pp.767-e1.
- Chen, L.K., Liu, L.K., Woo, J., Assantachai, P., Auyeung, T.W., Bahyah, K.S., Chou, M.Y., Chen, L.Y., Hsu, P.S., Krairit, O. and Lee, J.S., 2014. Sarcopenia in Asia: consensus report of the Asian Working Group for Sarcopenia. *Journal of the American Medical Directors Association*, 15(2), pp.95-101.
- Cruz-Jentoft, A.J., Baeyens, J.P., Bauer, J.M., Boirie, Y., Cederholm, T., Landi, F., Martin, F.C., Michel, J.P., Rolland, Y., Schneider, S.M. and Topinková, E., 2010. Sarcopenia: European consensus on definition and diagnosis Report of the European Working Group on Sarcopenia in Older People. *A. J. Cruz-Gentoft et al. Age and ageing*, 39(4), pp.412-423.
- Chien, M.Y., Huang, T.Y. and Wu, Y.T., 2008. Prevalence of sarcopenia estimated using a bioelectrical impedance analysis prediction equation in community-dwelling elderly people in Taiwan. *Journal of the American Geriatrics Society*, 56(9), pp.1710-1715.
- Clegg, A., Young, J., Iliffe, S., Rikkert, M. O., & Rockwood, K. (2013). Frailty in elderly people. *The lancet*, 381(9868), 752-762.
- Cheung, C. L., Nguyen, U. S. D., Au, E., Tan, K. C., & Kung, A. W. (2013). Association of handgrip strength with chronic diseases and multimorbidity. *Age*, 35(3), 929-941.
- Cappola, A. R., Bandeen-Roche, K., Wand, G. S., Volpato, S., & Fried, L. P. (2001). Association of IGF-I levels with muscle strength and mobility in older women. *The Journal of Clinical Endocrinology & Metabolism*, 86(9), 4139-4146.

- Chumlea, W. C., Guo, S. S., & Steinbaugh, M. L. (1994). Prediction of stature from knee height for black and white adults and children with application to mobility-impaired or handicapped persons. *Journal of the American Dietetic Association*, 94(12), 1385-1391.
- Druzhyna NM, Wilson GL, LeDoux SP. Mitochondrial DNA repair in aging and disease. *Mechanisms of ageing and development*. 2008 Jul 1;129(7-8):383-90.
- Drey, M., 2011. Sarcopenia–pathophysiology and clinical relevance. *Wiener Medizinische Wochenschrift*, 161(17-18), pp.402-408.
- Da Câmara, S.M.A., Alvarado, B.E., Guralnik, J.M., Guerra, R.O. and Maciel, Á.C.C., 2013. Using the Short Physical Performance Battery to screen for frailty in young-old adults with distinct socioeconomic conditions. *Geriatrics & gerontology international*, 13(2), pp.421-428.
- Essen-Gustavsson, B. and Borges, O., 1986. Histochemical and metabolic characteristics of human skeletal muscle in relation to age. *Acta Physiologica Scandinavica*, 126(1), pp.107-114.
- Fried, L.P., Tangen, C.M., Walston, J., Newman, A.B., Hirsch, C., Gottdiener, J., Seeman, T., Tracy, R., Kop, W.J., Burke, G. and McBurnie, M.A., 2001. Frailty in older adults: evidence for a phenotype. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 56(3), pp.M146-M157.
- Frontera, W.R., Hughes, V.A., Lutz, K.J. and Evans, W.J., 1991. A cross-sectional study of muscle strength and mass in 45-to 78-yr-old men and women. *Journal of applied physiology*, 71(2), pp.644-650.
- Frontera, W.R., Meredith, C.N., O'Reilly, K.P., Knuttgen, H.G. and Evans, W.J., 1988. Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *Journal of applied physiology*, 64(3), pp.1038-1044.
- Fiatarone, M.A. and Evans, W.J., 1993. 11 The Etiology and Reversibility of Muscle Dysfunction in the Aged. *Journal of Gerontology*, 48(Special_Issue), pp.77-83.
- Goodpaster, B.H., Park, S.W., Harris, T.B., Kritchevsky, S.B., Nevitt, M., Schwartz, A.V., Simonsick, E.M., Tylavsky, F.A., Visser, M. and Newman, A.B., 2006. The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 61(10), pp.1059-1064.

- Grimby, G., & Saltin, B. (1983). The ageing muscle. *Clinical physiology*, 3(3), 209-218.
- Hughes, V.A., Frontera, W.R., Wood, M., Evans, W.J., Dallal, G.E., Roubenoff, R. and Singh, M.A.F., 2001. Longitudinal muscle strength changes in older adults: influence of muscle mass, physical activity, and health. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 56(5), pp.B209-B217.
- Halliwell, B. and Gutteridge, J.M., 1985. Free radicals in biology and medicine.
- Haider, S., Luger, E., Kapan, A., Titze, S., Lackinger, C., Schindler, K.E. and Dorner, T.E., 2016. Associations between daily physical activity, handgrip strength, muscle mass, physical performance and quality of life in prefrail and frail community-dwelling older adults. *Quality of Life Research*, 25(12), pp.3129-3138.
- Houston, D.K., Cesari, M., Ferrucci, L., Cherubini, A., Maggio, D., Bartali, B., Johnson, M.A., Schwartz, G.G. and Kritchevsky, S.B., 2007. Association between vitamin D status and physical performance: the InCHIANTI study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 62(4), pp.440-446.
- Hudon, C., Fortin, M. and Vanasse, A., 2005. Cumulative Illness Rating Scale was a reliable and valid index in a family practice context. *Journal of clinical epidemiology*, 58(6), pp.603-608.
- Ito, K. and Barnes, P.J., 2009. COPD as a disease of accelerated lung aging. *Chest*, 135(1), pp.173-180.
- Iannuzzi-Sucich, M., Prestwood, K.M. and Kenny, A.M., 2002. Prevalence of sarcopenia and predictors of skeletal muscle mass in healthy, older men and women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57(12), pp.M772-M777.
- Janssen, I., Heymsfield, S.B. and Ross, R., 2002. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *Journal of the American Geriatrics Society*, 50(5), pp.889-896.
- Kim, Y.H., Kim, K.I., Paik, N.J., Kim, K.W., Jang, H.C. and Lim, J.Y., 2016. Muscle strength: A better index of low physical performance than muscle mass in older adults. *Geriatrics & gerontology international*, 16(5), pp.577-585.

- Kim, K.E., Jang, S.N., Lim, S., Park, Y.J., Paik, N.J., Kim, K.W., Jang, H.C. and Lim, J.Y., 2012. Relationship between muscle mass and physical performance: is it the same in older adults with weak muscle strength?. *Age and Ageing*, 41(6), pp.799-803.
- Kamo, T. and Nishida, Y., 2014. Direct and indirect effects of nutritional status, physical function and cognitive function on activities of daily living in Japanese older adults requiring long-term care. *Geriatrics & gerontology international*, 14(4), pp.799-805.
- Limpawattana, P., Kotruchin, P. and Pongchaiyakul, C., 2015. Sarcopenia in Asia. *Osteoporosis and Sarcopenia*, 1(2), pp.92-97.
- Lau, E.M., Lynn, H.S., Woo, J.W., Kwok, T.C. and Melton III, L.J., 2005. Prevalence of and risk factors for sarcopenia in elderly Chinese men and women. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 60(2), pp.213-216.
- Liu, L.K., Lee, W.J., Liu, C.L., Chen, L.Y., Lin, M.H., Peng, L.N. and Chen, L.K., 2013. Age-related skeletal muscle mass loss and physical performance in Taiwan: Implications to diagnostic strategy of sarcopenia in Asia. *Geriatrics & gerontology international*, 13(4), pp.964-971.
- Lexell, J., Taylor, C. C., & Sjöström, M. (1988). What is the cause of the ageing atrophy?: Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15-to 83-year-old men. *Journal of the neurological sciences*, 84(2-3), 275-294.
- Legrand, D., Adriaensen, W., Vaes, B., Matheï, C., Wallemacq, P. and Degryse, J., 2013. The relationship between grip strength and muscle mass (MM), inflammatory biomarkers and physical performance in community-dwelling very old persons. *Archives of gerontology and geriatrics*, 57(3), pp.345-351.
- Landi, F., Calvani, R., Tosato, M., Martone, A.M., Fusco, D., Sisto, A., Ortolani, E., Saveria, G., Salini, S. and Marzetti, E., 2017. Age-related variations of muscle mass, strength, and physical performance in community-dwellers: results from the Milan EXPO survey. *Journal of the American Medical Directors Association*, 18(1), pp.88-e17.
- Lamberts, S. W. (2002). The endocrinology of aging and the brain. *Archives of neurology*, 59(11), 1709-1711.
- Metter, E.J., Talbot, L.A., Schrager, M. and Conwit, R., 2002. Skeletal muscle strength as a predictor of all-cause mortality in healthy men. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57(10), pp.B359-B365.

- Moreira, V.G. and Lourenço, R.A., 2013. Prevalence and factors associated with frailty in an older population from the city of Rio de Janeiro, Brazil: the FIBRA-RJ Study. *Clinics*, 68(7), pp.979-985.
- Mijnarends, D.M., Meijers, J.M., Halfens, R.J., ter Borg, S., Luiking, Y.C., Verlaan, S., Schoberer, D., Jentoft, A.J.C., van Loon, L.J. and Schols, J.M., 2013. Validity and reliability of tools to measure muscle mass, strength, and physical performance in community-dwelling older people: a systematic review. *Journal of the American Medical Directors Association*, 14(3), pp.170-178.
- Mendes, J., Amaral, T.F., Borges, N., Santos, A., Padrão, P., Moreira, P., Afonso, C. and Negrão, R., 2017. Handgrip strength values of Portuguese older adults: a population based study. *BMC geriatrics*, 17(1), p.191.
- Morie, M., Reid, K.F., Miciek, R., Lajevardi, N., Choong, K., Krasnoff, J.B., Storer, T.W., Fielding, R.A., Bhasin, S. and LeBrasseur, N.K., 2010. Habitual physical activity levels are associated with performance in measures of physical function and mobility in older men. *Journal of the American Geriatrics Society*, 58(9), pp.1727-1733.
- Newman, A.B., Kupelian, V., Visser, M., Simonsick, E.M., Goodpaster, B.H., Kritchevsky, S.B., Tylavsky, F.A., Rubin, S.M. and Harris, T.B., 2006. Strength, but not muscle mass, is associated with mortality in the health, aging and body composition study cohort. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 61(1), pp.72-77.
- Ogawa, S., Yakabe, M. and Akishita, M., 2016. Age-related sarcopenia and its pathophysiological bases. *Inflammation and regeneration*, 36(1), p.17.
- Ong, H.L., Abdin, E., Chua, B.Y., Zhang, Y., Seow, E., Vaingankar, J.A., Chong, S.A. and Subramaniam, M., 2017. Hand-grip strength among older adults in Singapore: a comparison with international norms and associative factors. *BMC geriatrics*, 17(1), p.176.
- Peters, P. G., Alessio, H. M., Hagerman, A. E., Ashton, T., Nagy, S., & Wiley, R. L. (2006). Short-term isometric exercise reduces systolic blood pressure in hypertensive adults: possible role of reactive oxygen species. *International journal of cardiology*, 110(2), 199-205.
- Pusat Data Dan Informasi Kementerian Kesehatan, R., 2013. Gambaran Kesehatan Lanjut Usia di Indonesia. *Pusat Data dan Informasi Kementerian Kesehatan RI, Jakarta*.
- Parmelee, P.A., Thuras, P.D., Katz, I.R. and Lawton, M.P., 1995. Validation of the Cumulative Illness Rating Scale in a geriatric residential population. *Journal of the American Geriatrics Society*, 43(2), pp.130-137.

- Reed, R.L., Pearlmutter, L., Yochum, K., Meredith, K.E. and Mooradian, A.D., 1991. The relationship between muscle mass and muscle strength in the elderly. *Journal of the American Geriatrics Society*, 39(6), pp.555-561.
- Roppolo, M., Mulasso, A., Gobbens, R.J., Mosso, C.O. and Rabaglietti, E., 2015. A comparison between uni-and multidimensional frailty measures: prevalence, functional status, and relationships with disability. *Clinical interventions in aging*, 10, p.1669.
- Siparsky, P.N., Kirkendall, D.T. and Garrett Jr, W.E., 2014. Muscle changes in aging: understanding sarcopenia. *Sports Health*, 6(1), pp.36-40.
- Setiati , Harimurti K, & R Govinda Arya, 2014. Proses Menua dan Implikasi Kliniknya. In: Setiati S, Alwi I, Sudoyo A, Simadibrata M, Setiyohadi S, dan Syam A (Editors). *Buku Ajar Ilmu Penyakit Dalam Jilid III Edisi VI* Chapter 40, pp. 3669-3685.
- Taylor AC, McCartney N, Kamath MV, Wiley RL (2003) Isometric training lowers resting blood pressure and modulates autonomic control. *Med Sci Sports Exerc* 35:251–256
- Van Lummel, R.C., Walgaard, S., Pijnappels, M., Elders, P.J., Garcia-Aymerich, J., van Dieën, J.H. and Beek, P.J., 2015. Physical performance and physical activity in older adults: associated but separate domains of physical function in old age. *PLoS One*, 10(12), p.e0144048.
- Van den Akker, M., Buntinx, F., & Knottnerus, J. A. (1996). Comorbidity or multimorbidity: what's in a name? A review of literature. *The European Journal of General Practice*, 2(2), 65-70.
- World Health Organization. 2004. Health of Elderly in South East Asia: a profile. World Health Organization Regional Officer fo South East Asia.
- World Health Organization. 2011. World Population Prospect. The 2010 Revision United Nations
- Wu, S.W., Wu, S.F., Liang, H.W., Wu, Z.T. and Huang, S., 2009. Measuring factors affecting grip strength in a Taiwan Chinese population and a comparison with consolidated norms. *Applied ergonomics*, 40(4), pp.811-815.
- Yamada, Y., Watanabe, Y., Ikenaga, M., Yokoyama, K., Yoshida, T., Morimoto, T. and Kimura, M., 2013. Comparison of single-or multifrequency bioelectrical impedance analysis and spectroscopy for assessment of appendicular skeletal muscle in the elderly. *Journal of applied physiology*, 115(6), pp.812-818.

Yeh, C., Chen, Y.J., Lai, L.Y., Jang, T.R., Chiang, J., Chen, Y.Y. and Hsieh, K.C., 2012. Bioelectrical impedance analysis in a mathematical model for estimating fat-free mass in multiple segments in elderly Taiwanese males. *International Journal of Gerontology*, 6(4), pp.273-277.