

Specific Gyrb Sequence of Mycrobacterium Tuberculosis Clinical Isolated From Sputum of Pulmonary Tuberculosis Patients in Indonesia

by Ni Made Mertaniasih

Submission date: 28-Nov-2019 08:51PM (UTC+0800)

Submission ID: 1223264309

File name: From_Sputum_of_Pulmonary_Tuberculosis_Patients_in_Indonesia.pdf (771.61K)

Word count: 4667

Character count: 25633

SPECIFIC GYRB SEQUENCE OF MYCOBACTERIUM TUBERCULOSIS CLINICAL ISOLATED FROM SPUTUM OF PULMONARY TUBERCULOSIS PATIENTS IN INDONESIA

^{1,3}Ni Made Mertaniasih, ^{1,3}Nurul Wiqoyah, ^{1,3}Deby Kusumaningrum, ²Soedarsono, ³Agnes Dwi Sis Perwitasari, and ⁴Wayan Tunas Artama

¹Department of Clinical Microbiology, Faculty of Medicine Airlangga University/Dr. Soetomo Academic Hospital, Surabaya-Indonesia

²Department of Pulmonology and Medical Respiratory, Faculty of Medicine Airlangga University/ Dr. Soetomo Academic Hospital, Surabaya-Indonesia

³Department of Tuberculosis, Institute of Tropical Diseases, Airlangga University, Surabaya-Indonesia

⁴Department of Biochemistry, Faculty of Veterinary Medicine Gadjah Mada University, Yogyakarta-Indonesia

Background: Indonesia have many different geographic areas which could be various on the variant strains of *Mycobacterium tuberculosis*. The *gyrB* gene codes GyrB protein as sub unit compound of Gyrase enzyme that functioning in multiplication of bacteria. Detection of *gyrB* gene could be a marker of active multiplication of viable bacteria in the specimen from patients, and some of the DNA sequence regions were conserved and specific in the strain of *Mycobacterium tuberculosis* that would be a marker for identification. This research aims to analyze the sequence of *gyrB* gene of *Mycobacterium tuberculosis* clinical isolates from sputum of pulmonary TB patients in Indonesia, and determine the specific region. **Method:** *Mycobacterium tuberculosis* clinical isolates have been collected from sputum of the patients with pulmonary TB that live in some area in Indonesia. Isolation and identification of *Mycobacterium tuberculosis* clinical isolates using standard culture method; sequence analysis using PCR-direct sequencing of the part bases region of *gyrB*. **Results:** this study revealed that nucleotide sequence on a fragment 764 bases of *gyrB* gene *Mycobacterium tuberculosis* strains among clinical isolates almost identically to a wild type strain *Mycobacterium tuberculosis* H37Rv and subspecies member of *Mycobacterium tuberculosis* complex (MTBC), with a little difference of SNPs; there are many difference nucleotide sequence with MOTT and Gram positive or negative bacteria, except *Corynebacterium diphtheriae* identically with MTBC. **Conclusion:** the *gyrB* sequence in *Mycobacterium tuberculosis* strains among these clinical isolates from sputum of pulmonary TB patients in Indonesia have the conserved specific DNA region that almost identically with wild type strain H37Rv and MTBC.

Keywords: sequence; *gyrB* gene; *Mycobacterium*; tuberculosis; sputum.

INTRODUCTION

Tuberculosis (TB) is a major health problem in Indonesia and in the world. The WHO global TB control report of 2009, mentions in Indonesia noticed 528.063 new TB cases and 102 new sputum smear positive cases per 100.000 population, with the total population 226 million people in 2007 as reported by Indonesian Ministry of Health; the prevalence of TB in Indonesia differs per geographical areas, Java-Bali having the lowest prevalence (67 per

100.000 population), and Eastern Indonesia the highest (198 per 100.000 populations).¹ The other period reported at 2011 in Indonesia there are AFB positive 197.797 cases which the highest cases in Java; on the other hands at 2010 estimated the prevalence of pulmonary TB 725 per 100.000 population with the highest TB prevalence in Papua province 1.441 per 100.000 population, Batan 1.282/ 100.000, north Sulawesi 1.221/ 100.000, and lower in Lampung (270/ 100.000), Bali (306/ 100.000), DI. Yogyakarta (311/ 100.000) (Indonesia MOH, 2012).² At 2014, WHO reported the estimates of TB disease problem in Indonesia at 2012 noticed in 246.864.000 population with incidence rate 185 (153 – 220) per 100.000 population (95% CI); prevalence rate 297 (144 – 506); and mortality rate 27 (12 – 48). In South

Address of Correspondence:

Ni Made Mertaniasih

Department of Clinical Microbiology, Faculty of Medicine Airlangga University/Dr. Soetomo Academic Hospital, Surabaya-Indonesia
Email: mniasih@gmail.com

East Asia at 2012 estimate TB global incidence 8.300.000–9.000.000 cases of all forms of TB. In global TB report at 2013 estimated that total TB cases 39 % prevalent in South East Asia region of all TB cases in the world, and mortality estimated 47,6 % from totally in the world, estimated global TB mortality 790.000 – 1.100.000 cases of all form of TB.³

The effective control of TB case finding is of utmost importance. Accurate and rapid diagnostic methods are needed for determined TB diagnosis. Until now the culture method is the gold standard for TB diagnosis, on the other hands the culture method constrains with takes the time consuming, more than 3 weeks up till 2 months. Nucleic acid amplification is a rapid diagnostic methods that is sensitive and specific.⁴⁻⁶ Nucleic acid amplification tests have as the target gene region for identification, i.e. insertion sequence IS 6110, 16S rRNA, *hsp60*, *gyrB*, *rpoB*, 16-23S rRNA etc., which have the conserved and specific gene regions of *Mycobacterium tuberculosis*.⁷ High copy number of *gyrB* gene have the important role to distinguish the specific nucleotide sequence with MOTT species or other Gram positive bacteria.⁸ Based on the advantages of the role for the specific region identification in *gyrB* gene, as a basis primer design, the sequence of *gyrB* gene of *Mycobacterium tuberculosis* by DNA sequencing must be determine among clinical isolates from TB patients.

This study aims to analyze the sequence of *gyrB* gene *Mycobacterium tuberculosis* clinical isolates from sputum of pulmonary TB patients that live in some area in Indonesia. Some isolates from TB patients in many different geographic areas in Indonesia could be various strains of *M. tuberculosis*, however the *gyrB* gene nucleotide sequence would have the conserved and specific nucleotide sequence in the DNA regions.

MATERIALS AND METHOD

Bacterial Strains of *Mycobacterium tuberculosis*

Mycobacterium tuberculosis strains 102 isolated from sputum of 307 suspected pulmonary TB patients that collected in 2010, and 170 from 536 suspected pulmonary TB patients at 2013, patients from some area in Indonesia. Isolation and identification of *Mycobacterium tuberculosis* were carried out using standard culture method -MGIT 960 System (BD) and conventional standard on L-J medium (WHO). All of these were conducted in TB Laboratory in Department of Clinical Microbiology, Dr Soetomo Academic Hospital, Surabaya, Indonesia.

Analysis of DNA sequence of *gyrB* gene of *Mycobacterium tuberculosis* clinical isolates

Molecular study of DNA sequencing

conducted in Institute of Tropical Diseases, Airlangga University, Surabaya, Indonesia; eight isolates of TB patients (in 2010) have been done sequencing on a part of the target the 1.020 bases, and three isolates of TB patients (in 2013) with target a part of the almost whole gene sequence. Extraction and purification of DNA from *Mycobacterium tuberculosis* clinical isolates, as is follow, one loop full of fresh colonies of *Mycobacterium tuberculosis* (3 – 4 weeks old on Lowenstein-Jensen medium) was suspended in a buffer solution, extraction and purification of DNA using DNeasy kit (QIAGEN).

PCR to detect and amplify the DNA region of *gyrB* gene *Mycobacterium tuberculosis*, 41 isolates that collected at 2010 were conducted PCR for gene target 1.020-bp of *gyrB*. The suspension of PCR mix Dream Taq™ Green PCR Master Mix (Fermentas) was added by primer target 1.020-bp region as is MTUB-f 5'-TCGGACGCGTATGCGATATC-3' and MTUB-r 5'-ACATACAGTTCGGACTTGCG-3' each 1,0 µM, and DNA template 2µl, reaction volume 50 µl. Amplification reaction is 98° C 2 minutes; 96° C 20 seconds; 58° C 20 seconds; 72° C 1 minute; and 72° C 7 minutes, and the reaction 40 cycles.^{7,8}

Optimized PCR for target the sequence next to the whole genome of *gyrB* gene were done in 3 isolates from TB patients in 2013 using the five primer settings, Using *Mycobacterium tuberculosis* H37Rv as a positive control and mix reaction suspension without DNA as a negative control.

DNA sequencing and analysis of DNA sequence, PCR product is purified using QIAquick-spin PCR purification kit (QIAGEN). Eight isolates of *Mycobacterium tuberculosis* strains were conducted sequence analysis of 1.020 base region of *gyrB*, and three isolates sequenced of the next to the whole gene using ABI PRISM 310 System Sequencer. Alignment analysis was carried out using Genetix WIN V.10 program.

RESULTS

All of the 41 isolates (100%) *Mycobacterium tuberculosis* could be detect with PCR 41 strains of clinical isolates were positive *gyrB* conserved region at 1.020-bp (Figure 1, with example of five isolates).

Isolates P-1, P-2, P-3, P-4, P-5, P-6, P-7, P-8 from eight TB patients collected in 2010 have been sequenced analysis on alignment and phylogeny tree, revealed that all of these isolates were almost identically, more than 97 % with homolog nucleotide sequence in 764 bases that have been sequenced, with strain *Mycobacterium tuberculosis* H37Rv and other member of MTBC i.e. *Mycobacterium africanum*, *Mycobacterium bovis*, *Mycobacterium canettii*, *Mycobacterium microti*, *Mycobacterium caprae*; on the other hands

also closed related to *Corynebacterium diphtheriae*. These six strains are not identical to *Mycobacterium kansasii* and other MOTT i.e. *Mycobacterium avium*, *Mycobacterium intracellulare*, *Mycobacterium scrofulaceum*, *Mycobacterium asiaticum*, *Mycobacterium smegmatis*.




Figure 1

PCR products of *gyrB* gene *Mycobacterium tuberculosis* clinical isolates from sputum of pulmonary TB patients. Amplicon of *gyrB* at 1.020-bp. lane 1, 2, 3, 4, and 5 are amplicon from clinical isolates; M: 100-bp ladder; K+: positive control is amplicon from *Mycobacterium tuberculosis* H37Rv; negative control: suspension of reaction mixture without DNA template.

These strains distance correlated with *Nocardia*, and no correlated to Gram positive bacteria *Staphylococcus aureus* and Gram negative bacteria *Klebsiella pneumonia* and *Shigella sp* were indicated as the outgroup (Figure 2). In these 764 bases sequenced found a single nucleotide polymorphism (SNP) of isolate P4 at base no. 14 A – to – T, no. 169 G – to – T, no. 423 A – to – T, no. 446 G – to – T, no. 449 A – to – T; isolate P5 at base no. 28 C – to – A, no. 32 A – to – C; and isolate P6 no. 480 G – to – C (Figure 2). The 1.020-bp region position in the whole genome, flanked with sequence of the target by next setting primers which target to almost the whole genome.

The strains PS1, PS2, PS3 isolates from patients that collected at 2013 revealed, in around the 1054 bases a part of the target sequenced of almost the whole gene of *gyrB*, the result of alignment analysis and phylogeny tree similar with the strains from TB patients in 2010, were as followed: three strains isolates from patients almost identically with *Mycobacterium tuberculosis* H37Rv and other members of MTBC; distance correlated with MOTT; and no correlated to Gram positive and Gram negative bacteria, except *Corynebacterium diphtheriae* have the same sequence gene region closed correlated to MTBC.

These study also found SNPs in sequence *gyrB* gene of isolates PS1 at base no. 192 G – to – C, no. 436 C – to – G, no. 466 C – to – A, no. 866 G – to – C; and PS3 base no. 404 A – to – C, no. 410 A – to – C, no. 852 A – to – G. Isolate PS1 and PS2 also found two bases different with strain H37Rv at no. 417-418 AT – to – CC (Figure 2).

Figure 2 indicates sequence alignment of *gyrB* gene *Mycobacterium tuberculosis* clinical isolates from sputum of pulmonary TB patients in Indonesia at 2010 and 2013 (Genetix WIN V. 10 program), *Mycobacterium tuberculosis* H37Rv, other MTBC, MOTT, *Corynebacterium diphtheriae*, *Staphylococcus aureus*, and Gram negative bacteria.

Figure 3 Phylogeny tree of *gyrB* *Mycobacterium tuberculosis* clinical isolates of pulmonary TB patients in Indonesia at 2010 and 2013, *M. tuberculosis* H37Rv, MTBC, MOTT, Gram positive and negative bacteria (Genetix WIN V.10 program).


This phylogeny tree of *gyrB* *Mycobacterium tuberculosis* from clinical isolates have revealed the closed relation among the MTBC strains, on the other hands need more investigate the other DNA region for differentiate to *Corynebacterium spp*.

DISCUSSION


All of the 41 (100%) *Mycobacterium tuberculosis* clinical isolates from pulmonary TB patients in some area in Indonesia were positive for the target gene at 1.020-bp *gyrB*. Primer MTUB-f and MTUB-r were used for amplification of the 1.020-bp fragment of the *gyrB* gene *Mycobacterium tuberculosis* similar to investigation Chimara *et al.*, 2004 and Niemann *et al.*, 2000. The DNA region of *gyrB* gene. 1.020-bp fragments is amplified with species-specific primers MTUBf and MTUBr which do not generate amplicons from other species of mycobacteria.^{7,14}

In this study have showed the similar result of alignment analysis in the sequence 1.020 fragment and a part of almost the whole genome, that all clinical isolates *Mycobacterium tuberculosis* showed almost identically with H37Rv wild type and members of MTBC, and there are a little number of SNP; however close related to *Corynebacterium diphtheriae*, that could be need to investigate the other region of DNA fragment analyzed in *gyrB* gene regions. The nucleotide sequence of polymorphism in *gyrB* gene represents a unique marker that facilitates the differentiation of the MTBC.^{4,8}


Convenient system for bacterial identification using species-specific *gyrB* gene sequences, because of *gyrB* gene rarely transmitted horizontally, is distributed ubiquitously among bacterial species, its molecular evolution rate is higher than 16S rRNA, and the




2A




2B



2C



2D



2E

Figure 2A,B,C,D,E

Sequence alignment of *gyrB* gene *Mycobacterium tuberculosis* clinical isolates from sputum of pulmonary TB patients in Indonesia at 2010 and 2013 (Genetix WIN V. 10 program), *Mycobacterium tuberculosis* H37Rv, other MTBC, MOTT, *Corynebacterium diphtheriae*, *Staphylococcus aureus*, and Gram negative bacteria.




Figure 3

Phylogeny tree of *gyrB* *Mycobacterium tuberculosis* clinical isolates of pulmonary TB patients in Indonesia at 2010 and 2013, *M. tuberculosis* H37Rv, MTBC, MOTT, Gram positive and negative bacteria (Genetix WIN V.10 program).

essential roles of the *gyrB* product in DNA replication and transcription functions against the introduction of foreign *gyrB*. The *gyrB* sequences have the high correlation between the phylogenetic distance and the total genome homology analyzed by DNA-DNA hybridization.^{17,21} DNA topoisomerase are essential for DNA replication, transcription, recombination, repair and control the level of

supercoiling by cleaving and resealing the phosphodiester bond of DNA.⁵ They are classified into type I and type II according to enzymatic properties. The bacterial DNA gyrase is a type II topoisomerase that is capable of introducing negative supercoiling into a relaxed closed circular DNA molecule. DNA gyrase consists of two subunit protein-active form enzymes in the hetero-quaternary

complex structure of A2B2, A protein (GyrA) and B protein (Gyr B) that coding by *gyrA* and *gyrB* gene, only in prokaryotes.^{16,17,19} In *M. tuberculosis* presence only single topoisomerase I and DNA gyrase as a type II DNA topoisomerase.^{19,21,23} Expression of virulence genes depend on topological status of the genome, the topoisomerase as sensor of supercoils influence the specific gene expression.¹⁹ The high validity of *gyrB* sequences as taxonomic marker was evaluated mainly: the rate of their base substitutions and the consistency of the results of *gyrB*-based analysis. Protein-coding genes evolve faster than rRNA genes because synonymous substitutions mainly at the third positions of codons in the protein-coding genes are permitted without causing any changes the amino acid sequences. The average base-substitution rate of 16S rRNA genes was 1% per 50 million years, while *gyrB* was estimated 0.7 – 0.8% per one million years, therefore some species with identical 16S rDNA sequences can be differentiated using their *gyrB* sequences.¹⁷ The two genes encoding DNA gyrase in *Mycobacterium tuberculosis* are present next to each other in the genome, with *gyrB* upstream of *gyrA*, primary transcript is dicistronic with multiple promoters that appear to fine-tune transcription, the *gyr* genes in *Mycobacterium tuberculosis* as in other species are subject to autoregulation, albeit with slower kinetics, probably reflecting the slower metabolism of organism.^{16,19,21,24} The most probable translational start codon for *M. tuberculosis* GyrB is GTG (Val) which results in translation of a protein of 674 amino acids (74 kDa) at position 5240-5242 in *M. tuberculosis* genome sequence.¹⁶

Mycobacterium tuberculosis complex (MTBC) are the causative agents of tuberculosis in humans and animals, genetic close relationship, but differ in epidemiology pathogenicity, geographic range, host preferences, and in importance for tuberculosis disease in human. MTBC includes *Mycobacterium tuberculosis*, *M. bovis* subsp. *bovis*, *M. bovis* subsp. *caprae*, *M. bovis* BCG vaccine strain, *M. africanum* (subtype I and II), *M. microti*, *M. canettii*, *M. pinnipedii*.^{6,7,9,26} Routine differentiation is still base on phenotypic characteristics are time-consuming. Species belonging to MTBC cannot be differentiated by small subunit rRNA (16S rRNA) or internal transcribed spacer (ITS) 16-23S rRNA sequencing.⁵ Niemann *et al.* 2000 showed the result of DNA sequencing on gene region 1.020 bases of *gyrB* gene *Mycobacterium tuberculosis* and *Mycobacterium africanum* subtype II have an identical *gyrB* sequence that facilitates discrimination from the other species.⁸ All

members of MTBC are identical sequence gene, closely related gene with *gyrB* discriminatory regions in 1.020 bases. Base on these revealed, DNA sequence of species-specific region on 1.020-bp of *gyrB* gene *Mycobacterium tuberculosis* could be sole as the primer design for nucleic acid amplification that purpose to detect and identificate MTBC.

The *gyrB* could be useful for differentiation of MTBC and MOTT. The *gyrB*-based phylogeny was better than ITS-based phylogeny. The *gyrB* sequence of MTBC *M. tuberculosis*, *M. bovis*, *M. africanum*, *M. microti* found substitution at four sites, at 675 in *M. microti*, *M. bovis* at 756, at 1410 of *M. bovis*, *M. tuberculosis* at 1450, these substitutions can be regarded as the result of naturally occurring divergent evolution. The quantitation analysis *gyrB*-based method more useful than 16S rDNA-based method because the copy number of *gyrB* is single, while that of 16S rDNA is variable.^{9,17}

Based on sequenced genome of MTBC and outside the MTBC (*M. leprae*, *M. ulcerans*, *M. avium*, *M. paratuberculosis*, *M. marinum*, and fast-growing *M. smegmatis*, the most distant of the sequenced mycobacterial genomes are minimally related by 60% DNA/DNA homology, and comparative genomic analysis has shown that gene loss is a significant part of the ongoing evolution of the slow-growing mycobacterial pathogens. Single nucleotide polymorphisms (SNPs) can result a silent amino acid substitution in which the protein sequence remains unchanged, synonymous or can alter the protein coding sequence (non synonymous) and act as a substrate for evolutionary selection. The identification of SNP markers for study of evolution, pathogenesis, and epidemiology in clinical *M. tuberculosis* and *M. bovis*, the ratio of SNPs type within a genome can act as a molecular clock, the high ratio of non synonymous to synonymous mutations across coding sequences suggests a recent divergence of *M. tuberculosis* and *M. bovis*. The prominent role of genomic deletion relative to *M. tuberculosis* i.e. *M. bovis* contains 66.037 bp less than *M. tuberculosis* H37Rv, genomic deletion can be used to reconstruct phylogeny trees. Genomic deletion behave as unidirectional event polymorphism which represent one time events in the evolution, can serve as robust markers of clonal organisms for determining phylogeny classification. MTBC present relatively little genomic diversity, based on SNPs similar with large-sequence polymorphism, genomic flexibility exist within MTBC for specific host adaptation. Genomic deletions seem to several regions of difference (RD) as a genetic instability at a locus.²⁰ Namouchi at 2012 reported the nucleotide substitution is a major mechanism for the

emergence of *M. tuberculosis* pathogenesis, MTBC genome exhibit significant regional variation in the density of SNPs, the region of high SNPs may harbor rapidly evolving genes which positively selected for adaptation to human environment, the extremely conserved genes essential for survival.²⁶ The other advantages of *gyrB* as a marker differentiation, there were only a little evidence showed mutations that are related to quinolone resistance.^{10,12,13,22,25}

Gutierrez *et al.*, 2005, The highly successful human pathogen *M. tuberculosis* has the extremely low level of genetic variation, incongruence among gene phylogenies as mosaic gene sequences, despite its apparent homogeneity its genome appears to be a composite assembly resulting from horizontal gene transfer events predating clonal expansion, with synonymous nucleotide variation in housekeeping genes.¹⁵ Members of MTBC are the most successful human pathogens, *M. tuberculosis*, *M. bovis*, *M. africanum*, *M. microti*, *M. pinnipedii*, and species *M. caprae*, these members display different phenotypic characteristics and specific mammalian host ranges, they represent one of the most extreme genetic homogeneity with about 0.01%-0.03% synonymous nucleotide variation, and no significant trace of genetic exchange among them, it is believed the members are clonal progeny of a single successful ancestor that occurred 20,000 to 35,000 y ago. The six housekeeping genes, *katG*, *gyrB*, *gyrA*, *rpoB*, *hsp65*, *sodA*, and sequence of 16S rRNA revealed a smooth strain *M. canettii* and other members of MTBC form a single species defined by a compact phylogenetic clade.³³

Goh KS *et al.*, 2006, MTBC has been expanded additional members such as *M. canettii*, *M. pinnipedii*, *M. caprae*, *M. vaccae*, *M. africanum*, *M. canettii* infect only human, *M. tuberculosis* infect human may also occasionally infect domestic and wild animals upon exposure. Host range *M. bovis* and *M. caprae* is very broad, causing disease among a wide range of feral, domestic mammals, and human. Analysis of *gyrB* and *hsp65* targets could be used for differentiation of the species within MTBC.¹⁴

CONCLUSION

The nucleotide sequence of *gyrB* gene *Mycobacterium tuberculosis* clinical isolates from sputum of TB patients in Indonesia that collected at 2010 and 2013 revealed almost identically to *Mycobacterium tuberculosis* H37Rv and MTBC; all of 41/ 41 (100%) clinical isolates positive PCR with target 1.020-bp as the specific and conserved gene region.

ACKNOWLEDGMENTS

The authors would like to thanks to DP3M Ditjen Dikti, Indonesia Ministry of Education; Rector of Airlangga University; Director of LPPM Airlangga University; Dean of Medical Faculty, Airlangga University, Director of Dr Soetomo Academic Hospital, Director of Institute of Tropical Diseases, for all kinds of supporting in this TB research.

REFERENCES

1. WHO. Global TB Control Report. 2009.
2. MOH (Indonesia Ministry of Health). 2012.
3. WHO. Global TB Control Report. 2014.
4. Fukushima M, Kakinuma K, Hayashi H, Nagai H, Ito K, Kawaguchi R. 2003. Detection and Identification of *Mycobacterium species* Isolates by DNA Microarray. *J. Clin. Microbiol.* 41(6): 2605 – 2615.
5. Katoch VM, Parashar D, Chauhan DS, Singh D, Sharma VD, Ghosh. 2007. Rapid identification of mycobacteria by gene amplification restriction analysis technique targeting 16S-23S ribosomal RNA internal transcribed spacer & flanking region. *Indian J Med Res.* 125: 155 – 162.
6. Nakajima C, Rahim Z, Fukushima Y, Sugawara I, Zanden AG, Tamaru A, and Suzuki Y. 2010. Identification of *Mycobacterium tuberculosis* clinical isolates in Bangladesh by a species distinguishable multiplex PCR. *BMC Infectious Diseases.* 10: 118 – 125.
7. Chimara E, Ferrazoli L, and Leao SC. 2004. *Mycobacterium tuberculosis* Complex Differentiation Using *gyrB* Restriction Fragment Length Polymorphism Analysis. *Mem Inst Oswaldo Cruz.* 99(7): 745-748.
8. Neimann S, Harmsen D, Susch-Gerdes S, and Richter E. 2000. Differentiation of Clinical *Mycobacterium tuberculosis* Complex Isolates by *gyrB* DNA Sequence Polymorphism Analysis. *J. Clin. Microbiol.* 38 (9): 3231-3234.
9. Aranaz A, Liebaba E, Gomez-Mampazo E, Galan J, Cousins D, Ortega A *et al.* 1999. *Mycobacterium tuberculosis* subsp. *caprae* subsp. Nov. a taxonomic study of a new member of *Mycobacterium tuberculosis* complex isolated from goats in Spain. *Int J Syst Bacteriol.* 49: 1263 – 1273.
10. Aubry A, Veziris N, Cambau E, Truffot-Pernot C, Jarlier V, and Fisher M. 2006. Novel Gyrase Mutations in Quinolone-Resistant and Hypersusceptible Clinical Isolates of *Mycobacterium tuberculosis*: Functional Analysis of Mutant Enzymes. *Antimicrobial Agents and Chemother.* 50(1): 104-112.
11. Brosch R, Gordon SV, Maresse M, Bordin P, Buchrieser C, Eiglmeier K, *et al.* 2002. A new evolutionary scenario for *Mycobacterium*

- tuberculosis* complex. Proc Natl Acad Sci USA. 99: 3684 – 3689.
12. Cui Z, Wang J, Lu J, Huang X, and Hu Z. 2011. Association of mutation patterns in *gyrAB* genes and ofloxacin resistance levels in *Mycobacterium tuberculosis* isolates from East China in 2009. BMC Infectious Diseases. 11: 78 – 83.
 13. Feuerriegel S, Cox HS, Zarkua N, Karimovich HA, Braker K, Rusch-Gerdes S, and Nieman S. 2009. Sequence Analysis of Just Four Genes to Detect Extensively Drug-Resistant *Mycobacterium tuberculosis* Strains in Multidrug-Resistant Tuberculosis Patients Undergoing Treatment. Antimicrobial Agents and Chemoter. 53(8): 3353 – 3356.
 14. Goh KS, Fabre M, Huard RC, Schmid S, Sola C, and Rastogi N. 2006. Study of the *gyrB* gene polymorphism as a tool to differentiate among *Mycobacterium tuberculosis* complex subspecies further underlines the older evolutionary age of *Mycobacterium canettii*. Molecular and Cellular Probes. 20: 182 – 190.
 15. Gutierrez MC, Brisse S, Brosch R, Fabre M, Omais B, Marmiesse M, Supply P, and Vincent V. 2005. Ancient Origin and Gene Mosaicism of the Progenitor of *Mycobacterium tuberculosis*. PLoS Pathog 1(1): e5: 0055 – 0061.
 16. Karkare S, Brown A, Parish T, and Maxwell. 2013. Identification of the likely translational start of *Mycobacterium tuberculosis* GyrB. BMC research notes. 6: 274 – 280.
 17. Kasai H, Watanabe K, Gasteiger E. 1997. Construction of the *gyrB* Database for the Identification and Classification of Bacteria. Nature. 25: 3389-3402.
 18. Kasai H, Ezaki T, and Harayama S. 2000. Differentiation of Phylogenetically Related Slowly Growing Mycobacteria by their *gyrB* Sequences. J. Clin. Microbiol. 38(1): 301-08.
 19. Manjunatha UH, Madhusudan K, Unniraman S, Sikder D, Chatterjee M, Bhaduri T, Radha DR, and Nagaraja V. 2006. DNA topoisomerases from *Mycobacterium tuberculosis* and *Mycobacterium smegmatis*. J. Indian Inst. Sci. 86: 751 – 761.
 20. Mostowy S, Marcel A, and Behr MD. 2005. The Origin and Evolution of *Mycobacterium tuberculosis*. Clin Chest Med. 26: 207 – 216.
 21. Nagaraja V. 2004. Regulation of DNA topology in mycobacteria. Current Science 86(1): 135 – 139.
 22. Pitaksajakul P, Wongwit W, Punprasit W, Eampokalap B, Peacock S, and Ramasoota P. 2005. Mutations in The *GYRA* and *GYRB* Genes of Fluoroquinolone-Resistant *Mycobacterium tuberculosis* from TB Patients in Thailand. Southeast Asian J. Trop. Med. Public Health. 36(suppl 4): 228-237.
 23. Takiff HE, Salazar L, Guerrero C, Philipp W, Huang WM, Kreiswirth B, Cole ST, Jacobs WR, and Telenti A. 1994. Cloning and Nucleotide Sequence of *Mycobacterium tuberculosis gyrA* and *gyrB* Genes and Detection of Quinolone Resistance Mutations. Antimicrobial Agents and Chemoter. 38 (4): 773-780.
 24. Unniraman S, Chatterji M, and Nagaraja V. 2002. DNA Gyrase Genes in *Mycobacterium tuberculosis*: a Single Operon Driven by Multiple Promoters. J. Bacteriol. 184 (19): 5449– 5456.
 25. Von Groll A, Martin A, Jureen P, Hoffner S, Vandamme P, Portaels F, Palomino JC, and da Silva PA. 2009. Fluoroquinolone Resistance in *Mycobacterium tuberculosis* and Mutations in *gyrA* and *gyrB*. Antimicrobial Agents and Chemoter. 53(10): 4498-4500.
 26. Wang TC and Chen FC. 2013. The evolutionary landscape of the *Mycobacterium tuberculosis* genome. Gene 518: 187 – 193.



This work is licensed under
a Creative Commons Attribution

Specific GyrB Sequence of Mycobacterium Tuberculosis Clinical Isolated From Sputum of Pulmonary Tuberculosis Patients in Indonesia

ORIGINALITY REPORT

21%

SIMILARITY INDEX

14%

INTERNET SOURCES

18%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	Chimara, Erica, Lucilaine Ferrazoli, and Sylvia Cardoso Leão. "Mycobacterium tuberculosis complex differentiation using gyrB-restriction fragment length polymorphism analysis", Memórias do Instituto Oswaldo Cruz, 2004. Publication	2%
2	journals.plos.org Internet Source	2%
3	newgenforex.com Internet Source	2%
4	johnhawks.net Internet Source	1%
5	seasquirt.mbio.co.jp Internet Source	1%
6	repository.ias.ac.in Internet Source	1%
7	Goh, K.S.. "Study of the gyrB gene	

polymorphism as a tool to differentiate among Mycobacterium tuberculosis complex subspecies further underlines the older evolutionary age of 'Mycobacterium canettii', Molecular and Cellular Probes, 200606/08

Publication

1%

8

preview-bmcsresnotes.biomedcentral.com

Internet Source

1%

9

Juliëtte A. Severin. "Faecal carriage of extended-spectrum β -lactamase-producing Enterobacteriaceae among humans in Java, Indonesia, in 2001-2002 : ESBL colonisation in Indonesia", Tropical Medicine & International Health, 04/2012

Publication

1%

10

Ji-Hye Han, Myoung-Ho Cho, Seung Bum Kim. "Ribosomal and protein coding gene based multigene phylogeny on the family Streptomycetaceae", Systematic and Applied Microbiology, 2012

Publication

1%

11

Tai-Chun Wang, Feng-Chi Chen. "The evolutionary landscape of the Mycobacterium tuberculosis genome", Gene, 2013

Publication

1%

12

krishikosh.egranth.ac.in

Internet Source

<1%

13

Shoeb, M., D. L. de Castilla, and P. Pottinger. "The forest for the trees: tuberculosis control efforts in west Kalimantan", *Tropical Doctor*, 2011.

Publication

<1%

14

www.sidilv.org

Internet Source

<1%

15

B. Bayraktar. "Species Distribution of the Mycobacterium tuberculosis Complex in Clinical Isolates from 2007 to 2010 in Turkey: a Prospective Study", *Journal of Clinical Microbiology*, 11/01/2011

Publication

<1%

16

D. Alland. "Modeling Bacterial Evolution with Comparative-Genome-Based Marker Systems: Application to Mycobacterium tuberculosis Evolution and Pathogenesis", *Journal of Bacteriology*, 06/01/2003

Publication

<1%

17

B. Mathema. "Molecular Epidemiology of Tuberculosis: Current Insights", *Clinical Microbiology Reviews*, 10/01/2006

Publication

<1%

18

"Molecular Identification, Systematics, and Population Structure of Prokaryotes", *Springer Nature*, 2006

Publication

<1%

19

Motoshima, M.. "Rapid and accurate detection of *Pseudomonas aeruginosa* by real-time polymerase chain reaction with melting curve analysis targeting *gyrB* gene", *Diagnostic Microbiology & Infectious Disease*, 200705

Publication

<1%

20

link.springer.com

Internet Source

<1%

21

academicjournals.org

Internet Source

<1%

22

A. Somoskovi. "Sequencing of the *pncA* Gene in Members of the *Mycobacterium tuberculosis* Complex Has Important Diagnostic Applications: Identification of a Species-Specific *pncA* Mutation in "*Mycobacterium canettii*" and the Reliable and Rapid Predictor of Pyrazinamide Resistance", *Journal of Clinical Microbiology*, 11/15/2006

Publication

<1%

23

J. R. Lopez. "*Pseudomonas baetica* sp. nov., a novel fish pathogen isolated from wedge sole, *Dicologlossa cuneata* (Moreau)", *INTERNATIONAL JOURNAL OF SYSTEMATIC AND EVOLUTIONARY MICROBIOLOGY*, 06/03/2011

Publication

<1%

24

Internet Source

<1%

25

hdl.handle.net

Internet Source

<1%

26

www.searo.who.int

Internet Source

<1%

27

"Stroke Society of Australasia, Annual Scientific Meeting, 22-24 October 1997", Journal of Clinical Neuroscience, 199901

Publication

<1%

28

www.omicsonline.org

Internet Source

<1%

29

www.patentsencyclopedia.com

Internet Source

<1%

30

www.alice.cnptia.embrapa.br

Internet Source

<1%

31

microbewiki.kenyon.edu

Internet Source

<1%

32

article.sapub.org

Internet Source

<1%

33

Khye Seng Goh, Michel Fabre, Richard C. Huard, Solveig Schmid, Christophe Sola, Nalin Rastogi. "Study of the gyrB gene polymorphism as a tool to differentiate among Mycobacterium

<1%

tuberculosis complex subspecies further underlines the older evolutionary age of 'Mycobacterium canettii"', Molecular and Cellular Probes, 2006

Publication

34

"Tuberculosis", Springer Nature, 2004

Publication

<1%

35

Nosova, E., A. Bukatina, Y. Isaeva, M. Makarova, K. Galkina, and A. Moroz. "Analysis of mutations in the gyrA and gyrB genes and their association with the resistance of Mycobacterium tuberculosis to levofloxacin, moxifloxacin and gatifloxacin", Journal of Medical Microbiology, 2012.

Publication

<1%

36

Nabin K Shrestha. "Rapid diagnostic testing for mycobacterial infections", Future Microbiology, 08/2007

Publication

<1%

37

bmcpublichealth.biomedcentral.com

Internet Source

<1%

38

Nasir Mahmood, Shahzada N. Abbas, Neelam Faraz, Saman Shahid. "Mutational analysis of gyrB at amino acids: G481A & D505A in multidrug resistant (MDR) tuberculosis patients", Journal of Infection and Public Health, 2019

Publication

<1%

39

scholarbank.nus.edu.sg

Internet Source

<1%

40

S. Unniraman. "DNA Gyrase Genes in *Mycobacterium tuberculosis*: a Single Operon Driven by Multiple Promoters", *Journal of Bacteriology*, 10/01/2002

Publication

<1%

41

Kawabata, Hiroki, Seiji Sakakibara, Yasuyuki Imai, Toshiyuki Masuzawa, Hiromi Fujita, Miyako Tsurumi, Fumio Sato, Ai Takano, Sadao Nogami, Kazuhide Kaneda, and Haruo Watanabe. "First Record of *Leptospira borgpetersenii* Isolation in the Amami Islands, Japan", *Microbiology and Immunology*, 2006.

Publication

<1%

42

"Strain Variation in the *Mycobacterium tuberculosis* Complex: Its Role in Biology, Epidemiology and Control", Springer Science and Business Media LLC, 2017

Publication

<1%

43

I. C. SHAMPUTA. "Molecular genetic methods for diagnosis and antibiotic resistance detection of mycobacteria from clinical specimens", *Apmis*, 12/2004

Publication

<1%

44

Gaby E. Pfyffer. "*Mycobacterium tuberculosis* Complex, *Mycobacterium leprae*, and Other

<1%

Slow-Growing Mycobacteria", Topley & Wilson s Microbiology and Microbial Infections, 03/15/2010

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Specific GyrB Sequence of Mycobacterium Tuberculosis Clinical Isolated From Sputum of Pulmonary Tuberculosis Patients in Indonesia

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
