

AMERICAN JOURNAL OF  
**CHEMISTRY**

Effect of Ionic Liquids on the  
Hatching of *Artemia salina* Cysts



# Editorial Structure

## Editorial Board

Acceptance of papers is undertaken by the team of Associate Editors under the guidance of the Editor-in-Chief. All papers are refereed. Editorial Board members adjudicate in the case of conflicting or adverse reports.

## Meet the Editorial Board

### Editors-in-Chief

- George Koutsantonis, University of Western Australia
- John D. Wade, University of Melbourne and Florey Institute of Neuroscience and Mental Health

### Associate Editors

- Manabu Abé, Hiroshima University
- Stuart Batten, Monash University
- Paul Bernhardt, University of Queensland
- Cinzia Chiappe, University of Pisa
- Martyn Coles, Victoria University of Wellington
- Charlotte Conn, School of Applied Science, RMIT
- Deanna M. D'Alessandro, University of Sydney
- Peter Duggan, CSIRO
- Amanda Ellis, University of Melbourne
- Richard Hoogenboom, Ghent University
- T. S. Andy Hor, The University of Hong Kong
- Craig Hutton, University of Melbourne
- Amir Karton, University of Western Australia
- Anastasios (Tash) Polyzos, CSIRO CMHT
- Jenny Pringle, Deakin University
- Greg Qiao, University of Melbourne
- Devon Shipp, Clarkson University
- Pall Thordarson, University of New South Wales
- Xinhua Wan, Peking University
- Jonathan White, University of Melbourne

### Publisher

- Jenny Foster

# AUSTRALIAN JOURNAL OF CHEMISTRY

AN INTERNATIONAL JOURNAL FOR CHEMICAL SCIENCE

Volume 64 Number 4 2011

## CONTENTS

**Special issue: A dedication to Professor Athelstan L. J. Beckwith**  
**Guest editor: Peter J. Duggan**

### Foreword

Foreword to Professor Athelstan L. J. Beckwith Special Issue  
*P. J. Duggan* 355

### Reviews

Athelstan L. J. Beckwith and the Flowering of Hex-5-enyl Radical Cyclization Chemistry.  
The Adelaide Years

*A. K. Serelis* 358

Cyclohexane-1,3-dione Oxime Ether Grass-Specific Herbicides and the Discovery of Butoxydim

*K. G. Watson* 367

Profluorescent Nitroxides as Sensitive Probes of Oxidative Change and Free Radical Reactions

*J. P. Blinco, K. E. Fairfull-Smith, B. J. Morrow, S. E. Bottle* 373

### Full Papers

Destabilization of Conjugated Systems of  $\alpha$ -Dicarbonyls and of Cyanogen

*A. A. Zavitsas, D. W. Rogers, N. Matsunaga* 390

O-H Bond Dissociation Energies

*B. Chan, M. Morris, L. Radom* 394

The Distal Effect of *N*-Electron-withdrawing Groups on the Stability of Peptide Carbon Radicals

*J. Ho, M. L. Coote, C. J. Eastor* 403

Synthetic and Computational Studies of Acyl Radical Cyclizations with  $\beta$ -Alkoxyacrylates:  
Formal Synthesis of ( $\pm$ )-Longianone

*H. M. Aitken, C. H. Schiesser, C. D. Donner* 409

Fluoroazaindolines by an Uncommon Radical *ipso*-Substitution of a C-F Bond

*Y. Laot, L. Petit, N. D. M. Tran, S. Z. Zard* 416

A Profluorescent Azaphenylene Nitroxide for Nitroxide-Mediated Polymerization

*J. M. Colwell, J. P. Blinco, C. Hulbert, K. E. Fairfull-Smith, S. E. Bottle* 426

A Potential New RAFT-Click Reaction or a Cautionary Note on the Use of Diazomethane  
to Methylate RAFT-synthesized Polymers

*M. Chen, G. Moad, E. Rizzardo* 433

An Oxidized Abasic Lesion as an Intramolecular Source of DNA Adducts

*L. Guan, M. M. Greenberg* 438

Reaction of *N*-Acyloxy-*N*-alkoxyamides with Biological Thiol Groups

*S. A. Glover, M. Adams* 443

Synthesis of 2,3-Dihydro-4(1*H*)-quinolones and the Corresponding 4(1*H*)-Quinolones  
via Low-Temperature Fries Rearrangement of *N*-Arylazetid-2-ones

*J. Lange, A. C. Bissember, M. G. Banwell, I. A. Cade* 454

Studies Directed Towards the Preparation of Probes for the Photoaffinity Labelling of Gibberellin Receptors

*J. R. Crow, P. M. Chandler, L. N. Mander* 471

**Phytochemical Study of *Fagraea* spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties**

*Suciati, L. K. Lambert, B. P. Ross, M. A. Deseo, M. J. Garson* 489

Wood Protection Properties of Quaternary Ammonium Spiroborate Esters Derived from Alkyl Tartrates

*J. M. Carr, P. J. Duggan, D. G. Humphrey, J. A. Platts, E. M. Tyndall* 495

**Bukti online**

<http://www.publish.csiro.au/paper/CH10421.htm>

## CHEMISTRY

An international journal for chemical science

## Search

This Journal

GO

Advanced Search

## Journal Home

About the Journal  
New Editor-in-Chief  
Editorial Structure  
Contacts  
For Advertisers

## Content

Online Early  
Current Issue  
Just Accepted  
All Issues  
Virtual Issues  
Special Issues  
Research Fronts  
Sample Issue  
Covers

## For Authors

General Information  
Scope  
Submit Article  
Author Instructions  
Open Access

## For Referees

Referee Guidelines  
Review an Article

## For Subscribers

Subscription Prices  
Customer Service  
Print Publication Dates

## e-Alerts

Subscribe to our Email Alert  
or RSS feeds for the latest  
journal papers.

## Connect with us




Affiliated with RACI



## Table of Contents

&lt;&lt; Previous Issue | Next Issue &gt;&gt;


**Australian Journal of Chemistry**  
**Volume 64 Number 4 2011**

A Dedication to Professor Athelstan L. J. Beckwith

## Table of Contents

PDF (80 KB)

## Foreword to Professor Athelstan L. J. Beckwith Special Issue

Peter J. Duggan  
pp. 355-357




Professor Athelstan Beckwith, FRACI, FAA, FRS, OA passed away on 15th May 2010. He was a giant of Australian Chemistry and his contributions to the field of physical organic chemistry and free radical research have had major impacts worldwide. Athel published over 200 papers which have received more than 9900 citations. He was a great supporter of the *Australian Journal of Chemistry*, publishing more articles in our journal than anywhere else. This Special Issue of the *Australian Journal of Chemistry* is dedicated to him.

Full Text | PDF (235 KB)

**Athelstan L. J. Beckwith and the Flowering of Hex-5-enyl Radical Cyclization Chemistry. The Adelaide Years**

Algirdas K. Sereilis  
pp. 358-366




The late Professor Athel Beckwith made many significant contributions to the emergence of free radicals as controllable and selective reactive intermediates in synthesis. This account revisits his early mechanistic studies on the cyclization of hex-5-enyl radical systems, considered to be his most important work and the springboard for a multitude of applications in the synthesis of polycyclic natural products and related compounds.

### Studies Directed Towards the Preparation of Probes for the Photoaffinity Labelling of Gibberellin Receptors

James R. Crow, Peter M. Chandler and Lewis N. Mander

pp. 471-488



Model studies for the preparation of photoaffinity probes designed to explore the nature of gibberellin receptor sites have provided a wide range of gibberellin derivatives that should afford useful scaffolds for incorporating auxiliary groups attached to C-2 and C-12. Compounds were screened for bioactivity in growth and barley endosperm-based bioassays.

Abstract | PDF (570 KB) - \$25.00 **BUY NOW** | Supplementary Material (1.3 MB)

### Phytochemical Study of *Fagraea* spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties

Suciati, Lynette K. Lambert, Benjamin P. Ross, Myrna A. Deseo and Mary J. Garson

pp. 489-494




A new alkaloid fagraeoside, likely derived from the condensation of secologanin with L-asparagine, was isolated from *Fagraea racemosa* JACK *ex* WALL (Loganiaceae) along with secologanoside. Known lignans, flavonoids, iridoid and secoiridoid glucosides were isolated from three other species, namely *F. blumei* G. Don, *F. auriculata* Jack, Mal, and *F. ceilanica* Thunb. The anti-inflammatory, cytotoxic and anti-cholinesterase activity of selected compounds was evaluated.

Abstract | PDF (241 KB) - \$25.00 **BUY NOW** | Supplementary Material (294 KB)

### Wood Protection Properties of Quaternary Ammonium Spiroborate Esters Derived from Alkyl Tartrates

Jenny M. Carr, Peter J. Duggan, David G. Humphrey, James A. Platts and Edward M. Tyndall

pp. 495-502



## Phytochemical Study of *Fagraea* spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties<sup>1</sup>

Suciati,<sup>A,B</sup> Lynette K. Lambert,<sup>C</sup> Benjamin P. Ross,<sup>D</sup>  
Myrna A. Deseo,<sup>E</sup> and Mary J. Garson<sup>A,F</sup>

<sup>A</sup>School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>B</sup>Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia.

<sup>C</sup>Centre for Advanced Imaging, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>D</sup>School of Pharmacy, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>E</sup>Centre for Phytochemistry and Pharmacology, Southern Cross University, PO Box 157, Lismore, NSW 2480, Australia.

<sup>F</sup>Corresponding author. Email: m.garson@uq.edu.au

Phytochemical investigation of the stem bark of *F. racemosa* JACK *ex* WALL (Loganiaceae) from East Java, Indonesia, has resulted in the isolation of a new alkaloid fagraeoside along with the iridoid glycoside secologanoside. Fagraeoside may be derived from the condensation of secologanin with L-asparagine, and represents a rare example of a terpene alkaloid in which the amino acid component is non-aromatic. Investigation of three additional species of *Fagraea* provided known lignans, iridoid or secoiridoid glycosides, and flavanol-6-C-glucosides, thus it is likely that iridoid and secoiridoid glycosides are chemotaxonomic markers for the *Fagraea* genus. Fagraeoside inhibited the production of prostaglandin E<sub>2</sub> in 3T3 murine fibroblasts (IC<sub>50</sub> ~ 5.1 μM), and was not cytotoxic to this cell line or to a P388 murine leukaemia cell line. Selected isolated compounds, including fagraeoside, showed low to moderate activity in anti-acetylcholinesterase screening.

Manuscript received: 20 November 2010.

Manuscript accepted: 6 January 2011.

### Introduction

*Fagraea* (Loganiaceae) is a genus of shrubs or small trees consisting of ~50 species, which are distributed throughout Indo-Malaya, China, and Australia. Species of this genus often have large flowers and can live as an epiphyte in host trees with roots eventually reaching to the ground.<sup>[1,2]</sup> *Fagraea* has been used as a traditional medicine in many parts of South East Asia, including in Cambodia, Malaysia, the Philippines, and in Vietnam. The leaves have been used mainly for their analgesic and antipyretic effects, while a decoction of the bark is used as an antimalarial, an antipyretic, or as an anti-inflammatory agent.<sup>[3,4]</sup>

Previous studies on the chemistry of *Fagraea* spp. have identified lignans, including (+)-pinoresinol, (+)-epipinoresinol, (+)-lariciresinol, and (+)-isolariciresinol, and phenols from the root of *F. racemosa* JACK *ex* WALL collected in Sabah, Borneo, Malaysia.<sup>[5]</sup> The methanolic extract of *F. obovata* leaves obtained from Jogfalls, Karnataka, India, contained the glycosyl flavone fagovatin.<sup>[4]</sup> Other compounds including a benzyl alcohol derivative, di-*O*-methylcarenatin, the flavone C-glucoside swertisin and some iridoid glucosides, namely blumeosides A–D, were isolated from *F. blumei* G Don,<sup>[6]</sup> while the alkaloid gentianine was isolated from the leaves and fruits of

*F. fragrans*.<sup>[7]</sup> Over 50 volatile compounds have been isolated from the white flowers of *F. berteriana*, with these compounds responsible for the characteristic odor of the flower.<sup>[8]</sup> We now report a study on constituents of stem bark from *Fagraea* spp. collected in East Java, Indonesia which gave a new alkaloid, named by us as fagraeoside, together with eleven known compounds. The structures of the isolated compounds were deduced on the basis of NMR spectroscopic data and mass spectrometry. The anti-inflammatory, cytotoxic, and anti-acetylcholinesterase effects of selected metabolites have also been investigated.

### Results and Discussion

Powdered, air-dried stem bark samples from four species of *Fagraea* were collected in East Java, Indonesia and extracted sequentially with hexanes, then DCM and finally MeOH. The methanol extract from *F. racemosa* was purified by reverse phase flash chromatography followed by C<sub>18</sub> HPLC yielding fagraeoside (1), whose structure was established by 1D and 2D NMR spectroscopy, and secologanoside (2), identified by comparison with literature data.<sup>[9]</sup>

For compound (1), HRESIMS revealed an intense ion at *m/z* 469.1468 [M – H]<sup>–</sup>, corresponding to a molecular formula of

<sup>1</sup>Dedicated to the memory of the late Professor Athel Beckwith, an outstanding Australian chemist.

Table 1. NMR data (D<sub>2</sub>O) for fagraeoside

Position	$\delta_C^A$	$\delta_H, m (J \text{ in Hz})^B$	COSY <sup>C</sup>	<sup>1</sup> H- <sup>13</sup> C HMBC <sup>D</sup>
1	99.8	5.53, d (1.7)	9	3, 5, 8, 1'
3	151.2	7.43, d (2.2)	5	1, 4, 5, 11
4	110.4	—	—	—
5	26.6	3.10, m	3, 6b	3, 6, 7, 9
6	33.2	a 2.22, td (12.8, 4.0) b 1.58, ddd (12.8, 11.5, 10.6)	6b, 7 6a, 5, 7	4 <sup>E</sup> , 5, 7 <sup>E</sup> , 9 4 <sup>E</sup> , 5, 7 <sup>E</sup> , 9
7	66.2	5.24, dd (10.6, 4.0)	6a, 6b	5, 6 <sup>E</sup> , 11 <sup>E</sup> , 2'' <sup>E</sup>
8	134.3	5.58, ddd (17.1, 10.1, 9.9)	9, 10a, 10b	1, 5, 9
9	44.8	2.80, m	1, 8	1, 4, 5, 8, 10
10	123.5	a 5.37, dd (17.1, 1.4) b 5.30, dd (10.1, 1.4)	8 8	8, 9 9
11	168.3	—	—	—
1'	100.9	4.83, d (8.0)	2'	1, 3', 5'
2'	75.4	3.28, dd (9.4, 8.0)	1', 3'	—
3'	78.2	3.50, t (9.4)	2', 4'	2', 4'
4'	72.4	3.40, dd (9.8, 9.4)	3', 5'	3', 5', 6'
5'	79.1	3.51, ddd (9.8, 6.0, 2.1)	4', 6'b	4'
6'	63.5	a 3.92, dd (12.4, 2.1) b 3.72, dd (12.4, 6.0)	6'b 6'a, 5'	4' 5'
2''	175.5	—	—	—
3''	36.0	a 2.83, dd (16.6, 7.3) b 2.70, dd (16.6, 6.2)	3''b, 4'' 3''a, 4''	2'', 4'', 5'' 2'', 4'', 5''
4''	56.3	4.89, dd (7.3, 6.2)	3''a, 3''b	7, 11, 2'', 3'', 5''
5''	178.9	—	—	—

<sup>A</sup>Data recorded at 100 MHz, referenced to TSP at 0.0 ppm.

<sup>B</sup>Data recorded at 750 MHz, referenced to H<sub>2</sub>O at 4.77 ppm.

<sup>C</sup>Data recorded at 500 MHz.

<sup>D</sup>Data recorded at 750 MHz with <sup>1</sup>J<sub>CH</sub> 8 Hz.

<sup>E</sup>Additional correlations from data recorded at 750 MHz with <sup>1</sup>J<sub>CH</sub> 4 Hz.

C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>11</sub>. The <sup>13</sup>C NMR data (Table 1) were acquired in D<sub>2</sub>O and revealed 20 carbons, including three carbonyl signals at  $\delta_C$  168.3, 175.5, and 178.9. The <sup>1</sup>H NMR data, also in D<sub>2</sub>O, showed features typical of a  $\beta$ -glucose residue, including an anomeric proton at  $\delta_H$  4.83 (H-1') with the corresponding carbon at  $\delta_C$  100.9 by HSQC, a pair of multiplets (dd) at  $\delta_H$  3.92 and  $\delta_H$  3.72 for H-6', and resonances for four methine protons between  $\delta_H$  3.51 and 3.28. A full assignment of the glucose moiety was provided from TOCSY, DQF-COSY, and NOE data.

Other structural features resembled a secoiridoid fragment, and included an alkene proton at  $\delta_H$  7.43 (H-3) with the corresponding carbons at  $\delta_C$  151.2 (C-3) and 110.4 (C-4), an acetal proton at  $\delta_H$  5.53 (H-1) correlated with a carbon at  $\delta_C$  99.8 by HSQC, and resonances for three vinylic protons at  $\delta_H$  5.58 (H-8), 5.37 (H-10a), and 5.30 (H-10b), with the corresponding alkene carbons at  $\delta_C$  134.3 (C-8) and 123.5 (C-10).<sup>2</sup> A proton signal at  $\delta_H$  5.24 (1H, dd,  $J = 10.6, 4.0$ , H-7) linked to a carbon at  $\delta_C$  66.2 (d, C-7) supported either an  $\alpha$ -aminoacyl centre or an *N,N* acetal. The H-7 signal could be linked by DQF-COSY and/or TOCSY correlations to signals at  $\delta_H$  2.22 (H-6a),  $\delta_H$  1.58 (H-6b),  $\delta_H$  3.10 (H-5), and  $\delta_H$  2.80 (H-9). HSQC correlations then located C-5, C-6, C-7, and C-9, while geHMBC correlations (Fig. 1a) from H-3 and from H-7 to an amide-like carbonyl at  $\delta_C$  168.3 identified the remaining 'secoiridoid' carbon C-11. The HMBC correlation between H-1' and C-1 confirmed the position of the glucose moiety. When the <sup>1</sup>H NMR spectrum of fagraeoside was run in DMSO-*d*<sub>6</sub>, a broad NH singlet at  $\delta_H$  7.82 was detected which gave HMBC correlations to C-7 and NOESY correlations to H-7.

<sup>2</sup>Standard secoiridoid numbering used throughout.

The four remaining carbons were two carbonyls ( $\delta_C$  175.5, and 178.9), a methylene ( $\delta_C$  36.0) and a methine ( $\delta_C$  56.3). There were three unassigned proton signals: an  $\alpha$ -aminoacyl proton at  $\delta_H$  4.89 (1H, dd,  $J = 7.3, 6.2$ , H-4'') linked to the carbon at  $\delta_C$  56.3 by HSQC, and methylene protons at  $\delta_H$  2.83 (1H, dd,  $J = 16.6, 7.3$ , H-3''a) and 2.70 (1H, dd,  $J = 16.6, 6.2$ , H-3''b), linked to the signal at  $\delta_C$  36.0 by HSQC. HMBC correlations were from H-4'' to C-3'', to C-7, and notably to the amide C-11 as well as to the two carbonyls at  $\delta_C$  175.5 (C-2'') and 178.9 (C-5''). H-3'' showed correlations to C-2'', C-5'', as well as to C-4'', and H-7 at  $\delta_H$  5.24 showed correlations to C-2'' and, as mentioned, to C-11, while the NH signal showed a correlation to C-3''. These data fitted for a tetrahydropyrimid-4(3*H*)-one ring constructed biosynthetically from the condensation of an asparagine unit onto the secoiridoid framework (Fig. 2). An alternative tetrahydropyrimid-4(3*H*)-one structure with the free carboxyl group at C-3'' instead of at C-4'' was both inconsistent with the <sup>1</sup>H and <sup>13</sup>C chemical shifts of C-3'' and C-4'' and would be biosynthetically-unprecedented. The tetrahydropyrimid-4(3*H*)-one jadomycin N with  $\delta_C$  66.95 for the *N,N* acetal centre, a value comparable to that shown by C-7 of 1, has recently been isolated from cultures of *Streptomyces venezuelae* ISP5230 grown on L-asparagine as sole nitrogen source.<sup>10</sup>

The relative configuration was deduced from NOESY data (Fig. 1b) which showed correlations from H-5 to H-7 and to H-9, and between H-7 and the diastereotopic proton at  $\delta_H$  2.83 (H-3''a). Modelling of fagraeoside was undertaken using Chem Bio 3D ultra 12.0 (Cambridge); using MM2 software for energy minimization to an RMS gradient of 0.100, and revealed a



## New Terpene Alkaloid with Anti-Inflammatory Properties




Fig. 1. Structure of fagraeoside (1) showing (a) important HMBC correlations observed in  $d_6$ -DMSO; (b) MM2 minimised model showing selected nOe or NOESY correlations observed in  $d_6$ -DMSO (double-headed arrows).




Fig. 2. Proposed biosynthesis of fagraeoside from a secologanin-derived precursor and from L-asparagine.

flattened chair for the tetrahydropyrimidinone ring. In the <sup>1</sup>H NMR in D<sub>2</sub>O, H-4'' appeared as a doublet of doublets with *J* values of 7.3 Hz to H-3''a and of 6.2 Hz to H-3''b, matching the observed dihedral angles in **1** between H-4'' and H-3''a (56.7°) and H-3''b (61.8°). When the C-4'' epimer was modelled, the measured dihedral angles were respectively 64.6° and 175.9° which would have led to one large *J* value for the H-4'' signal. Although such data must be used with care, the absence of a NOESY correlation between H-7 and H-4'' was also consistent with the proposed relative configuration. In the C-4 epimer these protons are on the same face with an inter proton distance of 2.6 Å, whereas in **1**, they are on the opposite face, and thus 3.9 Å apart.

Another metabolite isolated from *F. racemosa* was (-)-secologanoside (**2**)<sup>[9]</sup> which was easily recognised by comparison of its NMR data with the literature. The presence of a D-glucose moiety in **1** and in **2** was confirmed by acid hydrolysis, derivatization with TFAA, and enantioselective gas chromatography.<sup>[11]</sup> The absolute configuration shown for **1** is further supported by its co-isolation with secologanoside, and by the proposed role of secologanin in the biosynthetic pathway of **1**. Consequently, L-asparagine is implicated in the biosynthesis of **1**.

Three other species of *Fagraea* were investigated for their chemistry. Stem bark from the species *F. blumei* yielded first, from the methanolic fraction, di-*O*-methylcrenatin (**3**).<sup>[12]</sup>

potalioside B (**4**),<sup>[12]</sup> the iridoid glucoside adoxosidic acid (**5**),<sup>[6,13]</sup> and its methyl ester derivative adoxoside (**6**),<sup>[14]</sup> second, from the DCM extract, (+)-pinoresinol (**7**)<sup>[5,15,16]</sup> and salicifolioside (**8**).<sup>[16]</sup> Di-*O*-methylcrenatin (**3**), adoxosidic acid (**5**), and adoxoside (**6**) have all been previously reported from *F. blumei*.<sup>[6]</sup> The iridoid glucoside loganic acid (**9**)<sup>[17]</sup> and the secoiridoid glucoside sweroside (**10**)<sup>[18]</sup> were isolated from *F. auriculata* together with potalioside B (**4**). An extract from *F. ceilanica* yielded two hydroxyflavanol C-glycosides whose NMR data matched those of taxifolin-6-*C*-glucoside (**11**)<sup>[19]</sup> and aromadendrin-6-*C*-glucoside (**12**)<sup>[20]</sup> together with secologanoside (**2**) and loganic acid (**9**) (Fig. 3).

In view of the traditional usage of *Fagraea* berk as an anti-inflammatory and antipyretic agent, compound **1** was screened for cytotoxicity and anti-inflammatory activity.<sup>[21]</sup> Although **1** inhibited the production of PGE<sub>2</sub> in 3T3 murine fibroblasts (IC<sub>50</sub> 5.1 μM), an atypical dose response curve was obtained (Fig. 4), even on repeat screening. Acetyl salicylic acid (IC<sub>50</sub> 2.6 μM) was used as a positive control. The observed anti-inflammatory activity did not appear to be a cytotoxic response since the compound lacked cytotoxicity to the 3T3 cells, and also to a P388 murine leukaemia cell line, when compared with chlorambucil as a reference compound. Additional testing is required to understand these effects. Although several iridoid glucosides show useful anti-inflammatory activity, there are few such reports for seco-iridoids.<sup>[22,23]</sup> Compounds **1**, **2**, **4**, **7**–**9**, **11**,




Fig. 3. Secoiridoid glycoside, iridoid glycoside, and lignan metabolites from *Fagraea* spp.




Fig. 4. Anti-inflammatory activity of fagraeoside against 3T3 murine fibroblast cells.

and 12 were also screened for anti-acetylcholinesterase activity in an assay using Ellman's reagent, 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB),<sup>[24]</sup> with the data quantified by detection at 415 nm using a microplate reader.<sup>[24,25]</sup> The inhibition observed

for all eight metabolites at  $100 \mu\text{g mL}^{-1}$  was similar to the inhibition found for galanthamine HBr salt at  $0.184 \mu\text{g mL}^{-1}$ , which indicated that the metabolites are relatively weak inhibitors of acetylcholinesterase.

### Conclusion

One of the largest group of plant alkaloids are the indole alkaloids formed from the condensation of secologanin with the aromatic amino acid tryptophan. This paper has described the structure and stereochemistry of a novel alkaloid that may be derived in Nature from the condensation of secologanin with the amino acid *t*-asparagine. There are remarkably few reports of terpene alkaloids in the literature in which the amino acid component is non-aromatic. Machida et al. (2002)<sup>[26]</sup> have described *t*-phenylalaninosecologanin, presumably formed by enzymatic reduction of the Schiff's base product of secologanin and *t*-phenylalanine, from *Lonicera japonica* Thunb. Our investigations on the secondary metabolites of the genus *Fagraea* have revealed the presence of iridoid and secoiridoid glycosides in all four species studied, and when considered together with an earlier literature report<sup>[6]</sup> suggest that such compounds may be chemotaxonomic markers for the genus.

## Experimental

### General

Reverse phase HPLC was conducted on an Agilent 1100 series instrument with variable wavelength UV detector set at 240 or at 254 nm or on a Shimadzu LC-20AT with a low temperature-evaporative light scattering detector (EL.S.D.-LT) fitted with a semi-preparative Waters<sup>®</sup> HPLC column ( $\mu$ Bondapak C<sub>18</sub> 7.8 × 300 mm) or a Phenomenex<sup>®</sup> column (Gemini 5 micron 10 × 250 mm). Positive or negative ion electrospray mass spectra (LRESMS) were determined using a Bruker Esquire HCT instrument or (HRESMS) on a MicroTof Q instrument, each with a standard ESI source. Optical rotation  $[\alpha]_D$  measurements were taken on a Perkin-Elmer 241 MC or a JASCO P-1010 polarimeter. The <sup>1</sup>H and <sup>13</sup>C NMR data were measured on Bruker Avance 400, 500, or 750 MHz spectrometers at 298 K. Normal phase and reverse phase thin-layer chromatography were performed on Merck Art 5554 and 5559 aluminium backed plates precoated with silica gel Merk Kieselgel 60 F254, respectively. Normal phase flash column chromatography was undertaken on Scharlau silica gel 60, 0.04–0.06 mm (230–400 mesh ASTM), while reverse phase flash column chromatography was carried out on silica gel prepared using the method developed by Evans et al.<sup>[27]</sup> Acetylcholinesterase inhibition assay measurements were performed using a Bio-Rad Model 550 microplate reader at 415 nm.

### Collection, Extraction, and Isolation

Stem bark of *F. blumei* G.Don (SC0106) was collected in Malang, East Java, Indonesia on 16 November 2006. Stem bark of *F. ceilanica* Thunb (SC0206) and *F. racemosa* JACK ex WALL (voucher sample SC0306) was collected on 16 November 2006 from Purwodadi Botanic Garden, East-Java, Indonesia. A methanolic extract of the stem bark of *F. auriculata* Jack, Mail (sample collected 26 February 2001; voucher FU0101) was a generous gift from Dr Achmad Fuad, Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia. All plants were identified by staff of Purwodadi Botanic Garden, East Java, Indonesia; voucher samples are stored in the Faculty of Pharmacy, Airlangga University. Plant materials were chopped and air-dried at room temperature for 5 days and then powdered using a blender.

Powdered stem bark of *F. racemosa* (530 g) was macerated with hexanes (3 × 24 h), then with DCM, and finally by MeOH. The methanolic extract (18 g) was fractionated by vacuum liquid chromatography, with a gradient of CHCl<sub>3</sub>, MeOH, and H<sub>2</sub>O in increasing polarity, and afforded 9 fractions. Fraction 8, namely FRM8, was then partitioned by NP flash chromatography, using EtOAc, MeOH, and H<sub>2</sub>O in order of increasing polarity. The first three fractions, namely FRM801, 802, and 803 eluting in EtOAc/MeOH (9:1), EtOAc/MeOH/H<sub>2</sub>O (9:1:1), and EtOAc/MeOH/H<sub>2</sub>O (8:2:1), respectively contained secologanoside (2) (134.6 mg). The fourth fraction, FRM804, was subjected to flash chromatography on SiO<sub>2</sub> using a solvent mixture of EtOAc/MeOH/H<sub>2</sub>O (8:2:1), followed by EtOAc/MeOH/H<sub>2</sub>O (5:5:1), and yielded 9 fractions. Fractions 7 and 8, namely FRM8047 and FRM8048, were then subjected to RP-HPLC using a gradient of 0–50% MeOH in H<sub>2</sub>O for 30 min, with UV detection at 240 nm, and flowrate 1.5 mL min<sup>-1</sup>. The component eluting at 20.9 min was fagraeoside (1) (3.5 mg).

Powdered stem bark of *F. blumei* (1 kg) was macerated in hexanes, followed by DCM and MeOH using the same procedure

as *F. racemosa*. A portion of the MeOH extract (382 mg) was chromatographed on RP-silica gel using a solvent gradient from 0 to 100% MeOH in H<sub>2</sub>O. A fraction eluting in 0–20% MeOH/H<sub>2</sub>O contained di-*O*-methylcresatin (3) (3.7 mg), potalioside B (4) (1.4 mg), and a mixture of adoxosidic acid (5) (3.4 mg) and adoxoside (6), which were further purified by RP-HPLC (0–50% MeOH in H<sub>2</sub>O for 60 min, detection at 254 nm). The DCM extract (11.8 g) was chromatographed on kieselgel 60H using hexanes, DCM, and MeOH in order of increasing polarity, followed by column chromatography on Diaion HP-20, using a solvent gradient (20% H<sub>2</sub>O in MeOH to 100% MeOH) yielding 5 fractions. Fraction 1 was then subjected to RP-HPLC (50% MeOH in H<sub>2</sub>O for 50 min, detection at 254 nm) to afford (+)-pinoresinol (7) (3.8 mg) and salicifoliol (8) (1.2 mg).

Extraction of powdered stem bark of *F. auriculata* was undertaken using the same procedure employed for *F. racemosa* and *F. blumei*. The MeOH extract (368 mg) was subjected to RP-flash column chromatography with a solvent gradient from 0 to 100% MeOH in H<sub>2</sub>O, and yielded 19 fractions. Fraction 4 was further purified by RP-HPLC with a gradient of 20 to 30% MeOH in water over 60 min to afford loganic acid (9) (1.16 mg). Fraction 5, eluting in 20% MeOH/H<sub>2</sub>O contained potalioside B (4) (8.79 mg), while fraction 6 (50% MeOH/H<sub>2</sub>O) afforded sweroside (10) (33.9 mg).

Stem bark of *F. ceilanica* (488 g) was extracted in the same way. A portion of the methanolic extract (3.9 g) was partitioned by RP-flash chromatography using a solvent gradient from 9 to 100% MeOH in H<sub>2</sub>O. Repeated RP-flash chromatography was undertaken on a fraction eluting in 100% H<sub>2</sub>O, followed by semi preparative RP-TLC with MeOH/H<sub>2</sub>O (3/7) to obtain five fractions. Fractions 2 and 3 were further purified by RP-HPLC (15–50% MeOH in H<sub>2</sub>O containing 0.1% TFA for 30 min, detection at 254 nm) to give taxifolin-6-*C*-glucoside (11) (22.9 mg) from fraction 2, and aromadendrin-6-*C*-glucoside (12) (14.1 mg), loganic acid (9) (5.8 mg), and secologanoside (2) (14.1 mg) from fraction 3.

**Fagraeoside (1):** Pale yellow amorphous solid (3.5 mg);  $[\alpha]_D^{25}$  –38.6 (c 0.57, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{max}$  (log  $\epsilon$ ) 241 (3.53) nm; <sup>1</sup>H and <sup>13</sup>C NMR (D<sub>2</sub>O, 750 and 100 MHz), see Table 1; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 750 MHz)  $\delta$  7.82 (1H, br s, NH), 7.28 (1H, d, *J* = 2.4, H-3), 5.47 (1H, ddd, *J* = 17.2, 10.3, 10.0, H-8), 5.36 (1H, d, *J* = 1.7, H-1), 5.30 (1H, dd, *J* = 17.1, 2.0, H-10a), 5.22 (1H, dd, *J* = 10.3, 2.0, H-10b), 5.19 (1H, dd, *J* = 11.6, 3.0, H-7), 4.90 (1H, br d, *J* = 6.8, H-4''), 4.51 (1H, d, *J* = 8.0, H-1'), 3.68 (1H, m, H-6'a), 3.42 (1H, m, H-6'b), 3.14 (2H, m, H-3' overlapping with H-5'), 3.03 (1H, m, H-4'), 2.98 (1H, m, H-2'), 2.85 (1H, m, H-5), 2.72 (1H, d, *J* = 16.0, H-3''a), 2.62 (1H, m, H-9), 2.17 (1H, dd, *J* = 16.0, 6.8, H-3''b), 2.00 (1H, m, H-6a), 1.20 (1H, m, H-6b); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 750 MHz, partial data from HMBC)  $\delta$  161.8 (C-11), 146.6 (C-3), 132.9 (C-8), 119.8 (C-10), 107.0 (C-4), 98.6 (C-1'), 95.4 (C-1), 77.1 (C-5'), 76.7 (C-3'), 72.8 (C-2'), 69.8 (C-4'), 62.6 (C-7), 61.0 (C-6'), 50.6 (C-4''), 42.0 (C-9), 34.4 (C-3''), 32.9 (C-6), 23.8 (C-5); HRESIMS calcd. for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>11</sub>, 469.1458. Found: *m/z* 469.1468 [M – H]<sup>-</sup>.

**Secologanoside (2):** Amorphous solid (134.6 mg);  $[\alpha]_D^{20}$  –98.3 (c 0.12, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{max}$  (log  $\epsilon$ ) 241 (3.53) nm; the <sup>1</sup>H and <sup>13</sup>C NMR data were comparable with published values;<sup>[9]</sup> LRESIMS *m/z* 413.2 [M + Na]<sup>+</sup>.

### Determination of Glucose Absolute Configuration

The glucose moieties were characterised according to the method of Hayes et al.<sup>[11]</sup> A sample of either fagraeoside

(0.74 mg) or secologanoside (5.63 mg) was heated for 2 h at 100°C with 10% HCl in MeOH (0.5 mL). After cooling, the solution was concentrated under N<sub>2</sub>, then partitioned between water and CHCl<sub>3</sub> (0.5 mL), and the aqueous layer concentrated under N<sub>2</sub>. The residue was dissolved in a mixture of trifluoroacetic anhydride (TFAA)/DCM (1/1, 0.2 mL) and then analysed by enantioselective chromatography (Chirasil-L-Val capillary column (25 m × 0.32 mm × 0.2 μm), FID detection, carrier gas: helium, injector: 200°C, column flow 1.32 mL min<sup>-1</sup>, split ratio: 56/0, and pressure 40 kPa. Temperature program: initial temperature 50°C for 6 min, raised at 4°C min<sup>-1</sup> to a final temperature of 160°C for 5 min). For comparison, standards were prepared from D- and L-glucose. Under these conditions, retention times were L-Glc (26.22 and 29.88 min), D-Glc (26.31 and 30.07 min). For glycoside 1, peaks were observed at 26.30 and 30.06 min, and for glycoside 2, peaks were observed at 26.26 and 30.02 min corresponding to D-glucose. Co-injection of either glucose pentatrifluoroacetate product with the D-glucose reference standard gave in each case a single peak.

#### Evaluation of Anti-Inflammatory Activity and Cytotoxicity

The prostaglandin E<sub>2</sub> assay was undertaken using Swiss albino mouse embryo fibroblast cells 3T3 (American Type Culture Collection (ATCC), Manassas, VA) and a PGE<sub>2</sub> enzyme immuno-assay kit (Prostaglandin E<sub>2</sub> ELA kit-mono-clonal, Cayman Chemical Co., Ann Arbor, MI, catalogue no. 5141010) using acetyl salicylic acid as a positive control. Cytotoxicity in 3T3 murine fibroblasts, and in P388D1 murine lymphoblasts (ATCC) was assayed using an ATPLite kit (PerkinElmer, Waltham, MA) with chlorambucil as a reference compound. Details of both assays have recently been published.<sup>[21]</sup>

#### Evaluation of Acetylcholinesterase Inhibition

Compounds were diluted in MeOH to a concentration of 1 mg mL<sup>-1</sup>. Galanthamine HBr salt was diluted in MeOH to a concentration of 100 μM. A solution of acetylthiocholine iodide in Millipore water (15 mM, 30 μL) was placed in each well of a 96-well plate. Then in sequence, 5,5'-dithiobis-(2-nitrobenzoic acid) in Millipore water (3 mM, 150 μL) and a buffer solution of bovine serum albumin (0.1%) made up in TRIS-HCl buffer (50 mM, pH 8.0, 60 μL) were added, followed by the compound under assay (30 μL) or by galanthamine HBr salt (30 μL). A blank well contained methanol (30 μL) instead of the compound under assay. The plate was shaken carefully to homogenize the mixture and the absorbance of individual wells at 415 nm was measured 10 times at intervals of 30 s. Acetylcholinesterase solution (0.22 U mL<sup>-1</sup>, 30 μL) was then added, the individual plates shaken and the absorbance measurements repeated a further 10 times. The experiments were done in quadruplicate in three independent experiments.

#### Accessory Publication

The Accessory Publication contains copies of representative NMR spectra, including the <sup>1</sup>H and <sup>13</sup>C spectra of fagraoside (1) in D<sub>2</sub>O, together with HMBC, NOESY, and ROESY data, and the <sup>1</sup>H and <sup>13</sup>C spectra of secologanoside (2) in D<sub>2</sub>O. Also included are characterisation details and spectroscopic data of compounds 2–12. The Accessory Publication is available on the Journal's website.

#### Acknowledgements

The authors thank AusAID for the award of an Australian Development Scholarship, The Queensland Pharmacy Research Trust (Pharmaceutical Society of Australia, Queensland branch) for financial support for the acetylcholinesterase inhibition assay, and Dr Achmad Fuad (Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia) for the *F. auriculata* sample. We thank Dr Patricia Hayes for assistance with the enantioselective GC characterisation of D-glucose. Mr Graham McFarlane and Dr Tri Le assisted with MS and NMR measurements respectively.

#### References

- [1] H. N. Ridley, *The Flora of the Malay Peninsula*, Vol. 2 1967 (L. Reeve & Co., Ltd.: London).
- [2] B. J. Conn, *Handbooks of the Flora of Papua New Guinea 1995* (Melbourne University Press: Melbourne).
- [3] C. Wiart, *Medicinal Plants of the Asia-Pacific: Drugs for the Future 2006* (World Scientific Publishing Ltd.: Singapore).
- [4] M. A. Qasim, S. K. Roy, M. Kamii, M. Iyas, *Phytochemistry* 1987, 26, 2871. doi:10.1016/S0031-9422(00)83614-6
- [5] E. Okuyama, K. Suzumura, M. Yamazaki, *Chem. Pharm. Bull. (Tokyo)* 1995, 43, 2200.
- [6] M. Cucudet, K. Hostettmann, O. Potterat, W. Dyatniko, *Helv. Chim. Acta* 1997, 80, 1144. doi:10.1002/HLCA.19970800411
- [7] A. S. C. Wan, Y. L. Chow, *J. Pharm. Pharmacol.* 1964, 16, 484.
- [8] S. Hayashi, H. Kameoka, S. Hashimoto, K. Furukawa, T. Arai, *J. Essent. Oil Res.* 1995, 7, 505.
- [9] I. Calis, O. Sticher, *Phytochemistry* 1984, 23, 2539. doi:10.1016/S0031-9422(00)84094-7
- [10] C. N. Borissow, C. L. Graham, R. T. Syvitski, T. R. Reid, J. Blay, D. L. Jakeman, *ChemBioChem* 2007, 8, 1198. doi:10.1002/CBIC.200700204
- [11] P. Y. Hayes, R. Lehmann, K. Penman, W. Kitching, J. J. De Voss, *Phytochemistry* 2009, 70, 105. doi:10.1016/J.PHYTOCHEM.2008.10.019
- [12] X. C. Li, H. N. FISOHLY, L. A. Walker, A. M. Clark, *Planta Med.* 2005, 71, 977. doi:10.1055/S-2005-871233
- [13] D. R. Gardner, J. Narum, D. Zook, F. R. Sternitz, *J. Nat. Prod.* 1987, 50, 485. doi:10.1021/NP50051A024
- [14] S. R. Jensen, B. J. Nielsen, *Biochem. Syst. Ecol.* 1979, 7, 103. doi:10.1016/0305-1978(79)90016-4
- [15] N. R. Guz, F. R. Sternitz, *Phytochemistry* 2000, 54, 897. doi:10.1016/S0031-9422(00)00101-1
- [16] A. G. González, R. Estévez-Reyes, C. Mato, *J. Nat. Prod.* 1989, 52, 1139. doi:10.1021/NP50065A037
- [17] I. Calis, M. F. Lahloub, O. Sticher, *Helv. Chim. Acta* 1984, 67, 160. doi:10.1002/HLCA.19840670119
- [18] Y. Zhou, Y. T. Di, S. Gesang, S. L. Peng, L. S. Ding, *Helv. Chim. Acta* 2006, 89, 94. doi:10.1002/HLCA.200690016
- [19] J. T. Mbafor, Z. T. Fernum, R. Promsathia, D. R. Sanson, M. S. Tempesta, *J. Nat. Prod.* 1989, 52, 417. doi:10.1021/NP50062A039
- [20] Y. Hayashi, S. Ohara, T. Takahashi, *Mokuzai Gakkaishi* 1987, 33, 511.
- [21] H. Wohlmut, M. A. Desco, D. J. Brushett, D. R. Thompson, G. MacFarlane, L. M. Stevenson, D. N. Leach, *J. Nat. Prod.* 2010, 73, 743. doi:10.1021/NP900688R
- [22] R. Tundis, M. R. Loizzo, F. Menichini, G. A. Statti, F. Menichini, *Mini Rev. Med. Chem.* 2008, 8, 399. doi:10.2174/138955708783955926
- [23] B. Dinda, D. R. Chowdhury, B. C. Mohanta, *Chem. Pharm. Bull. (Tokyo)* 2009, 57, 765. doi:10.1248/CPB.57.765
- [24] G. L. Ellman, K. D. Courtney, V. Andres, R. M. Featherstone, *Biochem. Pharmacol.* 1961, 7, 88. doi:10.1016/0006-2952(61)90145-9
- [25] I. K. Rhee, M. van de Meent, K. Ingkaninan, R. Verpoorte, *J. Chromatogr. A* 2001, 915, 217. doi:10.1016/S0021-9673(01)00624-0
- [26] K. Machida, H. Sasaki, T. Iijima, M. Kikuchi, *Chem. Pharm. Bull. (Tokyo)* 2002, 50, 1041. doi:10.1248/CPB.50.1041
- [27] M. B. Evans, A. D. Dale, C. J. Little, *Chromatographia* 1980, 13, 5. doi:10.1007/BF02302709

# Phytochemical Study of Fagraea spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties

*by* Suciati Suciati

---

**Submission date:** 26-Sep-2018 02:33PM (UTC+0800)

**Submission ID:** 1008647310

**File name:** fagraea-suci\_2011.pdf (241.18K)

**Word count:** 5114

**Character count:** 24252

## Phytochemical Study of *Fagraea* spp. Uncovers a New Terpene Aikaoid with Anti-Inflammatory Properties<sup>1</sup>

Suciati,<sup>A,B</sup> Lynette K. Lambert,<sup>C</sup> Benjamin P. Ross,<sup>D</sup>  
Myrna A. Deseo,<sup>E</sup> and Mary J. Garson<sup>A,F</sup>

<sup>A</sup>School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>B</sup>Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia.

<sup>C</sup>Centre for Advanced Imaging, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>D</sup>School of Pharmacy, The University of Queensland, Brisbane, Qld 4072, Australia.

<sup>E</sup>Centre for Phytochemistry and Pharmacology, Southern Cross University, PO Box 157, Lismore, NSW 2480, Australia.

<sup>F</sup>Corresponding author. Email: m.garson@uq.edu.au

Phytochemical investigation of the stem bark of *F. racemosa* JACK ex WALL (Loganiaceae) from East Java, Indonesia, has resulted in the isolation of a new alkaloid fagraeoside along with the iridoid glycoside secologanoside. Fagraeoside may be derived from the condensation of secologanin with L-asparagine, and represents a rare example of a terpene alkaloid in which the amino acid component is non-aromatic. Investigation of three additional species of *Fagraea* provided known lignans, iridoid or secoiridoid glycosides, and flavanol-6-C-glucosides, thus it is likely that iridoid and secoiridoid glycosides are chemotaxonomic markers for the *Fagraea* genus. Fagraeoside inhibited the production of prostaglandin E<sub>2</sub> in 3T3 murine fibroblasts (IC<sub>50</sub> ~5.1 μM), and was not cytotoxic to this cell line or to a P388 murine leukaemia cell line. Selected isolated compounds, including fagraeoside, showed low to moderate activity in anti-acetylcholinesterase screening.

Manuscript received: 20 November 2010.

Manuscript accepted: 6 January 2011.

### Introduction

*Fagraea* (Loganiaceae) is a genus of shrubs or small trees consisting of ~50 species, which are distributed throughout Indo-Malaya, China, and Australia. Species of this genus often have large flowers and can live as an epiphyte in host trees with roots eventually reaching to the ground.<sup>[1,2]</sup> *Fagraea* has been used as a traditional medicine in many parts of South East Asia, including in Cambodia, Malaysia, the Philippines, and in Vietnam. The leaves have been used mainly for their analgesic and antipyretic effects, while a decoction of the bark is used as an antimalarial, an antipyretic, or as an anti-inflammatory agent.<sup>[3,4]</sup>

Previous studies on the chemistry of *Fagraea* spp. have identified lignans, including (+)-pinoresinol, (+)-epipinoresinol, (+)-lariciresinol, and (+)-isolariciresinol, and phenols from the root of *F. racemosa* JACK ex WALL collected in Sabah, Borneo, Malaysia.<sup>[5]</sup> The methanolic extract of *F. obovata* leaves obtained from Jogfalls, Karnataka, India, contained the glycosyl flavone fagovatin.<sup>[4]</sup> Other compounds including a benzyl alcohol derivative, di-*O*-methylcresatin, the flavone C-glucoside swertisin and some iridoid glucosides, namely blumeosides A–D, were isolated from *F. blumei* G Don,<sup>[6]</sup> while the alkaloid gentianine was isolated from the leaves and fruits of

*F. fragrans*.<sup>[7]</sup> Over 50 volatile compounds have been isolated from the white flowers of *F. berteriana*, with these compounds responsible for the characteristic odor of the flower.<sup>[8]</sup> We now report a study on constituents of stem bark from *Fagraea* spp. collected in East Java, Indonesia which gave a new alkaloid, named by us as fagraeoside, together with eleven known compounds. The structures of the isolated compounds were deduced on the basis of NMR spectroscopic data and mass spectrometry. The anti-inflammatory, cytotoxic, and anti-acetylcholinesterase effects of selected metabolites have also been investigated.

### Results and Discussion

Powdered, air-dried stem bark samples from four species of *Fagraea* were collected in East Java, Indonesia and extracted sequentially with hexanes, then DCM and finally MeOH. The methanol extract from *F. racemosa* was purified by reverse phase flash chromatography followed by C<sub>18</sub> HPLC yielding fagraeoside (1), whose structure was established by 1D and 2D NMR spectroscopy, and secologanoside (2), identified by comparison with literature data.<sup>[9]</sup>

For compound (1), HRESIMS revealed an intense ion at *m/z* 469.1468 [M – H]<sup>+</sup> corresponding to a molecular formula of

<sup>1</sup>Dedicated to the memory of the late Professor Athel Beckwith, an outstanding Australian chemist.

Table 1. NMR data (D<sub>2</sub>O) for fagraeoside

Position	$\delta_c^A$	$\delta_H, m (J \text{ in Hz})^B$	COSY <sup>C</sup>	<sup>1</sup> H- <sup>13</sup> C HMBC <sup>D</sup>
1	99.8	5.53, d (1.7)	9	3, 5, 8, 1'
3	151.2	7.43, d (2.2)	5	1, 4, 5, 11
4	110.4	—	—	—
5	26.6	3.10, m	3, 6b	3, 6, 7, 9
6	33.2	a 2.22, td (12.8, 4.0) b 1.58, ddd (12.8, 11.5, 10.6)	6b, 7 6a, 5, 7	4 <sup>E</sup> , 5, 7 <sup>E</sup> , 9 4 <sup>F</sup> , 5, 7 <sup>E</sup> , 9
7	66.2	5.24, dd (10.6, 4.0)	6a, 6b	5, 6 <sup>E</sup> , 11 <sup>E</sup> , 2 <sup>''E</sup>
8	134.3	5.58, ddd (17.1, 10.1, 9.9)	9, 10a, 10b	1, 5, 9
9	44.8	2.80, m	1, 8	1, 4, 5, 8, 10
10	123.5	a 5.37, dd (17.1, 1.4) b 5.30, dd (10.1, 1.4)	8 8	8, 9 9
11	168.3	—	—	—
1'	100.9	4.83, d (8.0)	2'	1, 3', 5'
2'	75.4	3.28, dd (9.4, 8.0)	1', 3'	—
3'	78.2	3.50, t (9.4)	2', 4'	2', 4'
4'	72.4	3.40, dd (9.8, 9.4)	3', 5'	3', 5', 6'
5'	79.1	3.51, ddd (9.8, 6.0, 2.1)	4', 6'b	4'
6'	63.5	a 3.92, dd (12.4, 2.1) b 3.72, dd (12.4, 6.0)	6'b 6'a, 5'	4' 5'
2''	175.5	—	—	—
3''	36.0	a 2.83, dd (16.6, 7.3) b 2.70, dd (16.6, 6.2)	3''b, 4'' 3''a, 4''	2'', 4'', 5'' 2'', 4'', 5''
4''	56.3	4.89, dd (7.3, 6.2)	3''a, 3''b	7, 11, 2'', 3'', 5''
5''	178.9	—	—	—

<sup>A</sup>Data recorded at 100 MHz, referenced to TSP at 0.0 ppm.

<sup>B</sup>Data recorded at 750 MHz, referenced to H<sub>2</sub>O at 4.77 ppm.

<sup>C</sup>Data recorded at 500 MHz.

<sup>D</sup>Data recorded at 750 MHz with <sup>2</sup>J<sub>CH</sub> 8 Hz.

<sup>E</sup>Additional correlations from data recorded at 750 MHz with <sup>2</sup>J<sub>CH</sub> 4 Hz.

C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>11</sub>. The <sup>13</sup>C NMR data (Table 1) were acquired in D<sub>2</sub>O and revealed 20 carbons, including three carbonyl signals at  $\delta_c$  168.3, 175.5, and 178.9. The <sup>1</sup>H NMR data, also in D<sub>2</sub>O, showed features typical of a  $\beta$ -glucose residue, including an anomeric proton at  $\delta_H$  4.83 (H-1') with the corresponding carbon at  $\delta_c$  100.9 by HSQC, a pair of multiplets (dd) at  $\delta_H$  3.92 and  $\delta_H$  3.72 for H-6', and resonances for four methine protons between  $\delta_H$  3.51 and 3.28. A full assignment of the glucose moiety was provided from TOCSY, DQF-COSY, and nOe data.

Other structural features resembled a secoiridoid fragment, and included an alkene proton at  $\delta_H$  7.43 (H-3) with the corresponding carbons at  $\delta_c$  151.2 (C-3) and 110.4 (C-4), an acetal proton at  $\delta_H$  5.53 (H-1) correlated with a carbon at  $\delta_c$  99.8 by HSQC, and resonances for three vinylic protons at  $\delta_H$  5.58 (H-8), 5.37 (H-10a), and 5.30 (H-10b), with the corresponding alkene carbons at  $\delta_c$  134.3 (C-8) and 123.5 (C-10).<sup>2</sup> A proton signal at  $\delta_H$  5.24 (1H, dd,  $J = 10.6, 4.0$ , H-7) linked to a carbon at  $\delta_c$  66.2 (d, C-7) supported either an  $\alpha$ -aminoacyl centre or an *N,N* acetal. The H-7 signal could be linked by DQFCOSY and/or TOCSY correlations to signals at  $\delta_H$  2.22 (H-6a),  $\delta_H$  1.58 (H-6b),  $\delta_H$  3.10 (H-5), and  $\delta_H$  2.80 (H-9). HSQC correlations then located C-5, C-6, C-7, and C-9, while geHMBC correlations (Fig. 1a) from H-3 and from H-7 to an amide-like carbonyl at  $\delta_c$  168.3 identified the remaining 'secoiridoid' carbon C-11. The HMBC correlation between H-1' and C-1 confirmed the position of the glucose moiety. When the <sup>1</sup>H NMR spectrum of fagraeoside was run in DMSO-*d*<sub>6</sub>, a broad NH singlet at  $\delta_H$  7.82 was detected which gave HMBC correlations to C-7 and NOESY correlations to H-7.

<sup>2</sup>Standard secoiridoid numbering used throughout.

The four remaining carbons were two carbonyls ( $\delta_c$  175.5, and 178.9), a methylene ( $\delta_c$  36.0) and a methine ( $\delta_c$  56.3). There were three unassigned proton signals: an  $\alpha$ -aminoacyl proton at  $\delta_H$  4.89 (1H, dd,  $J = 7.3, 6.2$ , H-4'') linked to the carbon at  $\delta_c$  56.3 by HSQC, and methylene protons at  $\delta_H$  2.83 (1H, dd,  $J = 16.6, 7.3$ , H-3''a) and 2.70 (1H, dd,  $J = 16.6, 6.2$ , H-3''b), linked to the signal at  $\delta_c$  36.0 by HSQC. HMBC correlations were from H-4'' to C-3'', to C-7, and notably to the amide C-11 as well as to the two carbonyls at  $\delta_c$  175.5 (C-2'') and 178.9 (C-5''). H-3'' showed correlations to C-2'', C-5'', as well as to C-4'', and H-7 at  $\delta_H$  5.24 showed correlations to C-2'' and, as mentioned, to C-11, while the NH signal showed a correlation to C-3''. These data fitted for a tetrahydropyrimid-4(3H)-one ring constructed biosynthetically from the condensation of an asparagine unit onto the secoiridoid framework (Fig. 2). An alternative tetrahydropyrimid-4(3H)-one structure with the free carboxyl group at C-3'' instead of at C-4'' was both inconsistent with the <sup>1</sup>H and <sup>13</sup>C chemical shifts of C-3'' and C-4'' and would be biosynthetically-unprecedented. The tetrahydropyrimid-4(3H)-one jadomycin N with  $\delta_c$  66.95 for the *N,N* acetal centre, a value comparable to that shown by C-7 of 1, has recently been isolated from cultures of *Streptomyces venezuelae* ISP5230 grown on 1-asparagine as sole nitrogen source.<sup>109</sup>

The relative configuration was deduced from NOESY data (Fig. 1b) which showed correlations from H-5 to H-7 and to H-9, and between H-7 and the diastereotopic proton at  $\delta_H$  2.83 (H-3''a). Modelling of fagraeoside was undertaken using Chem Bio 3D ultra 12.0 (Cambridge), using MM2 software for energy minimization to an RMS gradient of 0.100, and revealed a




Fig. 1. Structure of fagraeoside (1) showing (a) important HMBC correlations observed in  $d_6$ -DMSO; (b) MM2 minimised model showing selected nOe or NOESY correlations observed in  $d_6$ -DMSO (double-headed arrows).




Fig. 2. Proposed biosynthesis of fagraeoside from a secologanin-derived precursor and from *L*-asparagine

flattened chair for the tetrahydropyrimidinone ring. In the  $^1\text{H}$  NMR in  $\text{D}_2\text{O}$ ,  $\text{H-4}'$  appeared as a doublet of doublets with  $J$  values of 7.3 Hz to  $\text{H-3}'\text{a}$  and of 6.2 Hz to  $\text{H-3}'\text{b}$ , matching the observed dihedral angles in **1** between  $\text{H-4}'$  and  $\text{H-3}'\text{a}$  ( $56.7^\circ$ ) and  $\text{H-3}'\text{b}$  ( $61.8^\circ$ ). When the  $\text{C-4}'$  epimer was modelled, the measured dihedral angles were respectively  $64.6^\circ$  and  $175.9^\circ$  which would have led to one large  $J$  value for the  $\text{H-4}'$  signal. Although such data must be used with care, the absence of a NOESY correlation between  $\text{H-7}$  and  $\text{H-4}'$  was also consistent with the proposed relative configuration. In the  $\text{C-4}$  epimer these protons are on the same face with an inter proton distance of 2.6 Å, whereas in **1**, they are on the opposite face, and thus 3.9 Å apart.

Another metabolite isolated from *F. racemosa* was (-)-secologanoside (**2**)<sup>91</sup> which was easily recognised by comparison of its NMR data with the literature. The presence of a D-glucose moiety in **1** and in **2** was confirmed by acid hydrolysis, derivatization with TFAA, and enantioselective gas chromatography.<sup>111</sup> The absolute configuration shown for **1** is further supported by its co-isolation with secologanoside, and by the proposed role of secologanin in the biosynthetic pathway of **1**. Consequently, *L*-asparagine is implicated in the biosynthesis of **1**.

Three other species of *Fagraea* were investigated for their chemistry. Stem bark from the species *F. blumei* yielded first, from the methanolic fraction, di-*O*-methylerenatin (**3**).<sup>121</sup>

potalioside **B** (**4**).<sup>112</sup> The iridoid glucoside adoxosidic acid (**5**).<sup>16,133</sup> and its methyl ester derivative adoxoside (**6**).<sup>144</sup> second, from the DCM extract, (+)-pinoresinol (**7**).<sup>15,15,16</sup> and salicifolol (**8**).<sup>161</sup> Di-*O*-methylerenatin (**3**), adoxosidic acid (**5**), and adoxoside (**6**) have all been previously reported from *F. blumei*.<sup>164</sup> The iridoid glucoside loganic acid (**9**)<sup>177</sup> and the secoiridoid glucoside sweroside (**10**)<sup>181</sup> were isolated from *F. auriculata* together with potalioside **B** (**4**). An extract from *F. ceilanica* yielded two hydroxyflavanol C-glycosides whose NMR data matched those of taxifolin-6-*C*-glucoside (**11**)<sup>191</sup> and aromadendrin-6-*C*-glucoside (**12**)<sup>201</sup> together with secologanoside (**2**) and loganic acid (**9**) (Fig. 3).

In view of the traditional usage of *Fagraea* bark as an anti-inflammatory and antipyretic agent, compound **1** was screened for cytotoxicity and anti-inflammatory activity.<sup>211</sup> Although **1** inhibited the production of  $\text{PGI}_2$  in 3T3 murine fibroblasts ( $\text{IC}_{50}$  5.1  $\mu\text{M}$ ), an atypical dose response curve was obtained (Fig. 4), even on repeat screening. Acetyl salicylic acid ( $\text{IC}_{50}$  2.6  $\mu\text{M}$ ) was used as a positive control. The observed anti-inflammatory activity did not appear to be a cytotoxic response since the compound lacked cytotoxicity to the 3T3 cells, and also to a P388 murine leukaemia cell line, when compared with chlorambucil as a reference compound. Additional testing is required to understand these effects. Although several iridoid glycosides show useful anti-inflammatory activity, there are few such reports for seco-iridoids.<sup>22,231</sup> Compounds **1**, **2**, **4**, **7-9**, **11**.






Fig. 3. Secoiridoid glycoside, iridoid glycoside, and lignan metabolites from *Fagraea* spp.




Fig. 4. Anti-inflammatory activity of fagraeoside against 3T3 murine fibroblast cells.

and **12** were also screened for anti-acetylcholinesterase activity in an assay using Ellman's reagent, 5,5'-dithiobis-(2-nitrobenzoic acid) (DTNB)<sup>[24,25]</sup> with the data quantified by detection at 415 nm using a microplate reader.<sup>[24,25]</sup> The inhibition observed

for all eight metabolites at  $100 \mu\text{g mL}^{-1}$  was similar to the inhibition found for galanthamine HBr salt at  $0.184 \mu\text{g mL}^{-1}$ , which indicated that the metabolites are relatively weak inhibitors of acetylcholinesterase.

#### Conclusion

One of the largest group of plant alkaloids are the indole alkaloids formed from the condensation of secologanin with the aromatic amino acid tryptophan. This paper has described the structure and stereochemistry of a novel alkaloid that may be derived in Nature from the condensation of secologanin with the amino acid L-asparagine. There are remarkably few reports of terpene alkaloids in the literature in which the amino acid component is non-aromatic. Machida et al. (2002)<sup>[26]</sup> have described L-phenylalaninosecologanin, presumably formed by enzymatic reduction of the Schiff's base product of secologanin and L-phenylalanine, from *Lonicera japonica* Thunb. Our investigations on the secondary metabolites of the genus *Fagraea* have revealed the presence of iridoid and secoiridoid glycosides in all four species studied, and when considered together with an earlier literature report<sup>[6]</sup> suggest that such compounds may be chemotaxonomic markers for the genus.

## Experimental

### General

Reverse phase HPLC was conducted on an Agilent 1100 series instrument with variable wavelength UV detector set at 240 or at 254 nm or on a Shimadzu LC-20AT with a low temperature- evaporative light scattering detector (EL.S.D.-LT) fitted with a semi-preparative Waters HPLC column ( $\mu$ Bondapak C<sub>18</sub> 7.8 × 300 mm) or a Phenomenex column (Gemini 5 micron 10 × 250 mm). Positive or negative ion electrospray mass spectra (LRESMS) were determined using a Bruker Esquire HCT instrument or (HRESMS) on a MicroTof Q instrument, each with a standard ESI source. Optical rotation  $[\alpha]_D$  measurements were taken on a Perkin-Elmer 241 MC or a JASCO P-1010 polarimeter. The <sup>1</sup>H and <sup>13</sup>C NMR data were measured on Bruker Avance 400, 500, or 750 MHz spectrometers at 298 K. Normal phase and reverse phase thin-layer chromatography were performed on Merck Art 5554 and 5559 aluminium backed plates pre-coated with silica gel Merk Kieselgel 60 F254, respectively. Normal phase flash column chromatography was undertaken on Scharlau silica gel 60, 0.04–0.06 mm (230–400 mesh ASTM), while reverse phase flash column chromatography was carried out on silica gel prepared using the method developed by Evans et al.<sup>127</sup> Acetylcholinesterase inhibition assay measurements were performed using a Bio-Rad Model 550 microplate reader at 415 nm.

### Collection, Extraction, and Isolation

Stem bark of *F. blumei* G. Don (SC0106) was collected in Malang, East Java, Indonesia on 16 November 2006. Stem bark of *F. ceilanica* Thunb (SC0206) and *F. racemosa* JACK ex WALL (voucher sample SC0306) was collected on 16 November 2006 from Purwodadi Botanic Garden, East Java, Indonesia. A methanolic extract of the stem bark of *F. auriculata* Jack, Mall (sample collected 26 February 2001; voucher FU0101) was a generous gift from Dr Achmad Fuad, Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia. All plants were identified by staff of Purwodadi Botanic Garden, East Java, Indonesia; voucher samples are stored in the Faculty of Pharmacy, Airlangga University. Plant materials were chopped and air-dried at room temperature for 5 days and then powdered using a blender.

Powdered stem bark of *F. racemosa* (530 g) was macerated with hexanes (3 × 24 h), then with DCM, and finally by MeOH. The methanolic extract (18 g) was fractionated by vacuum liquid chromatography, with a gradient of CHCl<sub>3</sub>, MeOH, and H<sub>2</sub>O in increasing polarity, and afforded 9 fractions. Fraction 8, namely FRM8, was then partitioned by NP flash chromatography, using EtOAc, MeOH, and H<sub>2</sub>O in order of increasing polarity. The first three fractions, namely FRM801, 802, and 803 eluting in EtOAc/MeOH (9:1), EtOAc/MeOH/H<sub>2</sub>O (9:1:1), and EtOAc/MeOH/H<sub>2</sub>O (8:2:1), respectively contained secologanoside (2) (134.6 mg). The fourth fraction, FRM804, was subjected to flash chromatography on SiO<sub>2</sub> using a solvent mixture of EtOAc/MeOH/H<sub>2</sub>O (8:2:1), followed by EtOAc/MeOH/H<sub>2</sub>O (5:5:1), and yielded 9 fractions. Fractions 7 and 8, namely FRM8047 and FRM8048, were then subjected to RP-HPLC using a gradient of 0–50% MeOH in H<sub>2</sub>O for 30 min, with UV detection at 240 nm, and flowrate 1.5 mL min<sup>-1</sup>. The component eluting at 20.9 min was fagraeoside (1) (3.5 mg).

Powdered stem bark of *F. blumei* (1 kg) was macerated in hexanes, followed by DCM and MeOH using the same procedure

as *F. racemosa*. A portion of the MeOH extract (382 mg) was chromatographed on RP-silica gel using a solvent gradient from 0 to 100% MeOH in H<sub>2</sub>O. A fraction eluting in 0–20% MeOH/H<sub>2</sub>O contained di-*O*-methylcresatin (3) (3.7 mg), potalioside B (4) (1.4 mg), and a mixture of adoxosidic acid (5) (3.4 mg) and adoxoside (6), which were further purified by RP-HPLC (0–50% MeOH in H<sub>2</sub>O for 60 min, detection at 254 nm). The DCM extract (11.8 g) was chromatographed on kieselgel 60H using hexanes, DCM, and MeOH in order of increasing polarity, followed by column chromatography on Diaion HP-20, using a solvent gradient (20% H<sub>2</sub>O in MeOH to 100% MeOH) yielding 5 fractions. Fraction 1 was then subjected to RP-HPLC (50% MeOH in H<sub>2</sub>O for 50 min, detection at 254 nm) to afford (+)-pinoresinol (7) (3.8 mg) and saicifolol (8) (1.2 mg).

Extraction of powdered stem bark of *F. auriculata* was undertaken using the same procedure employed for *F. racemosa* and *F. blumei*. The MeOH extract (368 mg) was subjected to RP-flash column chromatography with a solvent gradient from 0 to 100% MeOH in H<sub>2</sub>O, and yielded 19 fractions. Fraction 4 was further purified by RP-HPLC with a gradient of 20 to 30% MeOH in water over 60 min to afford loganic acid (9) (1.16 mg). Fraction 5, eluting in 20% MeOH/H<sub>2</sub>O contained potalioside B (4) (8.79 mg), while fraction 6 (50% MeOH/H<sub>2</sub>O) afforded sweroside (10) (33.9 mg).

Stem bark of *F. ceilanica* (488 g) was extracted in the same way. A portion of the methanolic extract (3.9 g) was partitioned by RP-flash chromatography using a solvent gradient from 0 to 100% MeOH in H<sub>2</sub>O. Repeated RP-flash chromatography was undertaken on a fraction eluting in 100% H<sub>2</sub>O, followed by semi preparative RP-TLC with MeOH/H<sub>2</sub>O (3/7) to obtain five fractions. Fractions 2 and 3 were further purified by RP-HPLC (15–50% MeOH in H<sub>2</sub>O containing 0.1% TFA for 30 min, detection at 254 nm) to give taxifolin-6-*C*-glucoside (11) (22.9 mg) from fraction 2, and aromadendrin-6-*C*-glucoside (12) (14.1 mg), loganic acid (9) (5.8 mg), and secologanoside (2) (14.1 mg) from fraction 3.

**Fagraeoside (1).** Pale yellow amorphous solid (3.5 mg);  $[\alpha]_D^{25}$  -38.6 (c 0.57, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{max}$  (log  $\epsilon$ ) 241 (3.53) nm; <sup>1</sup>H and <sup>13</sup>C NMR (D<sub>2</sub>O, 750 and 100 MHz), see Table 1; <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 750 MHz)  $\delta$  7.82 (1H, br s, NH), 7.28 (1H, d, *J* = 2.4, H-3), 5.47 (1H, ddd, *J* = 17.2, 10.3, 10.0, H-8), 5.36 (1H, d, *J* = 1.7, H-1), 5.30 (1H, dd, *J* = 17.1, 2.0, H-10a), 5.22 (1H, dd, *J* = 10.3, 2.0, H-10b), 5.19 (1H, dd, *J* = 11.6, 3.0, H-7), 4.98 (1H, br d, *J* = 6.8, H-4'), 4.51 (1H, d, *J* = 8.0, H-1'), 3.68 (1H, m, H-6'a), 3.42 (1H, m, H-6'b), 3.14 (2H, m, H-3' overlapping with H-5'), 2.03 (1H, m, H-4''), 2.98 (1H, m, H-2''), 2.85 (1H, m, H-5), 2.72 (1H, d, *J* = 16.0, H-3''a), 2.62 (1H, m, H-9), 2.17 (1H, dd, *J* = 16.0, 6.8, H-3''b), 2.00 (1H, m, H-6a), 1.20 (1H, m, H-6b); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>, 750 MHz, partial data from HMBC)  $\delta$  161.8 (C-11), 146.6 (C-3), 132.9 (C-8), 119.8 (C-10), 107.0 (C-4), 98.6 (C-1'), 95.4 (C-1), 77.1 (C-5'), 76.7 (C-3'), 72.8 (C-2'), 69.8 (C-4'), 62.6 (C-7), 61.0 (C-6'), 50.6 (C-4''), 42.0 (C-9), 34.4 (C-3''), 32.9 (C-6), 23.8 (C-5); HRESIMS calcd. for C<sub>20</sub>H<sub>26</sub>N<sub>2</sub>O<sub>11</sub>, 469.1458. Found: *m/z* 469.1468 [M - H]<sup>-</sup>.

**Secologanoside (2):** Amorphous solid (134.6 mg);  $[\alpha]_D^{20}$  -98.3 (c 0.12, H<sub>2</sub>O); UV (H<sub>2</sub>O)  $\lambda_{max}$  (log  $\epsilon$ ) 241 (3.53) nm; the <sup>1</sup>H and <sup>13</sup>C NMR data were comparable with published values;<sup>101</sup> LRESIMS *m/z* 413.2 [M + Na]<sup>+</sup>.

### Determination of Glucose Absolute Configuration

The glucose moieties were characterised according to the method of Hayes et al.<sup>111</sup> A sample of either fagraeoside

(0.74 mg) or secologanoside (5.63 mg) was heated for 2 h at 100°C with 10% HCl in MeOH (0.5 mL). After cooling, the solution was concentrated under N<sub>2</sub>, then partitioned between water and CHCl<sub>3</sub> (0.5 mL), and the aqueous layer concentrated under N<sub>2</sub>. The residue was dissolved in a mixture of trifluoroacetic anhydride (TFAA)/DCM (1/1, 0.2 mL) and then analysed by enantioselective chromatography (Chirasil-L-Val capillary column (25 m × 0.32 mm × 0.2 µm), FID detection, carrier gas: helium, injector: 200°C, column flow 1.32 mL min<sup>-1</sup>, split ratio: 56/0, and pressure 40 kPa. Temperature program: initial temperature 50°C for 6 min, raised at 4°C min<sup>-1</sup> to a final temperature of 160°C for 5 min). For comparison, standards were prepared from D- and L-glucose. Under these conditions, retention times were L-Glc (26.22 and 29.88 min), D-Glc (26.31 and 30.07 min). For glycoside 1, peaks were observed at 26.30 and 30.06 min, and for glycoside 2, peaks were observed at 26.26 and 30.02 min corresponding to D-glucose. Co-injection of either glucose pentatrifluoroacetate product with the D-glucose reference standard gave in each case a single peak.

#### Evaluation of Anti-inflammatory Activity and Cytotoxicity

The prostaglandin E<sub>2</sub> assay was undertaken using Swiss albino mouse embryo fibroblast cells 3T3 (American Type Culture Collection (ATCC), Manassas, VA) and a PGE<sub>2</sub> enzyme immuno-assay kit (Prostaglandin E<sub>2</sub> ELA kit-monoclonal, Cayman Chemical Co., Ann Arbor, MI, catalogue no. 5141010) using acetyl salicylic acid as a positive control. Cytotoxicity in 3T3 murine fibroblasts, and in P388D1 murine lymphoblasts (ATCC) was assayed using an ATP-Lite kit (PerkinElmer, Waltham, MA) with chlorambucil as a reference compound. Details of both assays have recently been published.<sup>[21]</sup>

#### Evaluation of Acetylcholinesterase Inhibition

Compounds were diluted in MeOH to a concentration of 1 mg mL<sup>-1</sup>. Galanthamine HBr salt was diluted in MeOH to a concentration of 100 µM. A solution of acetylthiocholine iodide in Millipore water (15 mM, 30 µL) was placed in each well of a 96-well plate. Then in sequence, 5,5'-dithiobis-(2-nitrobenzoic acid) in Millipore water (3 mM, 150 µL) and a buffer solution of bovine serum albumin (0.1%) made up in TRIS-HCl buffer (50 mM, pH 8.0, 60 µL) were added, followed by the compound under assay (30 µL) or by galanthamine HBr salt (30 µL). A blank well contained methanol (30 µL) instead of the compound under assay. The plate was shaken carefully to homogenize the mixture and the absorbance of individual wells at 415 nm was measured 10 times at intervals of 30 s. Acetylcholinesterase solution (0.22 U mL<sup>-1</sup>, 30 µL) was then added, the individual plates shaken and the absorbance measurements repeated a further 10 times. The experiments were done in quadruplicate in three independent experiments.

#### Accessory Publication

The Accessory Publication contains copies of representative NMR spectra, including the <sup>1</sup>H and <sup>13</sup>C spectra of fagraeoside (1) in D<sub>2</sub>O, together with HMBC, NOESY, and ROESY data, and the <sup>1</sup>H and <sup>13</sup>C spectra of secologanoside (2) in D<sub>2</sub>O. Also included are characterisation details and spectroscopic data of compounds 2–12. The Accessory Publication is available on the Journal's website.

#### Acknowledgements

The authors thank AusAID for the award of an Australian Development Scholarship, The Queensland Pharmacy Research Trust (Pharmaceutical Society of Australia, Queensland branch) for financial support for the acetylcholinesterase inhibition assay, and Dr Achmad Fuad (Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia) for the *F. auriculata* sample. We thank Dr Patricia Hayes for assistance with the enantioselective GC characterisation of D-glucose. Mr Graham McFarlane and Dr Tri Le assisted with MS and NMR measurements respectively.

#### References

- [1] H. N. Ridley, *The Flora of the Malay Peninsula*, Vol. 2 1967 (L. Reeve & Co., Ltd.: London).
- [2] B. J. Conn, *Handbooks of the Flora of Papua New Guinea* 1995 (Melbourne University Press: Melbourne).
- [3] C. Wiart, *Medicinal Plants of the Asia-Pacific: Drugs for the Future* 2006 (World Scientific Publishing Ltd.: Singapore).
- [4] M. A. Qasim, S. K. Roy, M. Kamil, M. Ilyas, *Phytochemistry* 1987, 26, 2871. doi:10.1016/S0031-9422(00)83614-6
- [5] E. Okuyama, K. Suzunura, M. Yamazaki, *Chem. Pharm. Bull. (Tokyo)* 1995, 43, 2200.
- [6] M. Cuendet, K. Hostettmann, O. Potterat, W. Dyatmiko, *Helv. Chim. Acta* 1997, 80, 1144. doi:10.1002/HLCA.19970800411
- [7] A. S. C. Wan, Y. L. Chow, *J. Pharm. Pharmacol.* 1964, 16, 484.
- [8] S. Hayashi, H. Kameoka, S. Hashimoto, K. Furukawa, T. Arai, *J. Essent. Oil Res.* 1995, 7, 505.
- [9] I. Calis, O. Sticher, *Phytochemistry* 1984, 23, 2539. doi:10.1016/S0031-9422(00)84094-7
- [10] C. N. Borissow, C. L. Graham, R. T. Syvitski, T. R. Reid, J. Blay, D. L. Jakeman, *Chemosocnem* 2007, 8, 1198. doi:10.1002/CHEM.200700204
- [11] P. Y. Hayes, R. Lehmann, K. Penman, W. Kitching, J. J. De Voss, *Phytochemistry* 2009, 70, 105. doi:10.1016/J.PHYTOCHEM.2008.10.019
- [12] X. C. Li, H. N. ElSohly, L. A. Walker, A. M. Clark, *Planta Med.* 2005, 71, 977. doi:10.1055/S-2005-871233
- [13] D. R. Gardner, J. Narum, D. Zook, F. R. Sternitz, *J. Nat. Prod.* 1987, 50, 485. doi:10.1021/NP50051A024
- [14] S. R. Jensen, B. J. Nielsen, *Biochem. Syst. Ecol.* 1979, 7, 103. doi:10.1016/0305-1978(79)90016-4
- [15] N. R. Guz, F. R. Sternitz, *Phytochemistry* 2000, 54, 897. doi:10.1016/S0031-9422(00)00101-1
- [16] A. G. González, R. Estévez-Reyes, C. Mato, *J. Nat. Prod.* 1989, 52, 1139. doi:10.1021/NP50063A057
- [17] I. Calis, M. F. Lahloub, O. Sticher, *Helv. Chim. Acta* 1984, 67, 160. doi:10.1002/HLCA.19840670119
- [18] Y. Zhou, Y. T. Di, S. Gesang, S. L. Peng, L. S. Ding, *Helv. Chim. Acta* 2006, 79, 34. doi:10.1002/HLCA.200609016
- [19] J. T. Mbafor, Z. T. Fomum, R. Promsaththa, D. R. Sanson, M. S. Tempesta, *J. Nat. Prod.* 1989, 52, 417. doi:10.1021/NP50062A039
- [20] Y. Hayashi, S. Ohara, T. Takahashi, *Mokuzai Gakkaishi* 1987, 33, 511.
- [21] H. Wohlmut, M. A. Deseo, D. J. Brashett, D. R. Thompson, G. MacFarlane, L. M. Stevenson, D. N. Leach, *J. Nat. Prod.* 2010, 73, 743. doi:10.1021/NP900688R
- [22] R. Tundis, M. R. Loizzo, F. Menichini, G. A. Starti, F. Menichini, *Mini Rev. Med. Chem.* 2008, 8, 399. doi:10.2174/138955708783955926
- [23] D. Dinda, D. R. Chowdhury, B. C. Mohanta, *Chem. Pharm. Bull. (Tokyo)* 2009, 57, 765. doi:10.1248/CPB.57.765
- [24] G. L. Ellman, K. D. Courtney, V. Andres, R. M. Featherstone, *Biochem. Pharmacol.* 1961, 7, 88. doi:10.1016/0006-2952(61)90145-9
- [25] I. K. Rhee, M. van de Meent, K. Ingkaninan, R. Verpoorte, *J. Chromatogr. A* 2001, 915, 217. doi:10.1016/S0021-9673(01)00624-0
- [26] K. Machida, H. Sasaki, T. Iijima, M. Kikuchi, *Chem. Pharm. Bull. (Tokyo)* 2002, 50, 1041. doi:10.1248/CPB.50.1041
- [27] M. B. Evans, A. D. Dale, C. J. Little, *Chromatographia* 1980, 13, 5. doi:10.1007/BF02302709

# Phytochemical Study of Fagraea spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties

ORIGINALITY REPORT

**20%**

SIMILARITY INDEX

**9%**

INTERNET SOURCES

**19%**

PUBLICATIONS

**0%**

STUDENT PAPERS

PRIMARY SOURCES

<b>1</b>	<a href="http://www.slideshare.net">www.slideshare.net</a> Internet Source	<b>3%</b>
<b>2</b>	Patricia Y. Hayes, Reg Lehmann, Kerry Penman, William Kitching, James J. De Voss. "Steroidal saponins from the roots of Trillium erectum (Beth root)", Phytochemistry, 2009 Publication	<b>2%</b>
<b>3</b>	Kitajima, M.. "Camptothecin-related alkaloids from hairy roots of Ophiorrhiza pumila", Tetrahedron, 20021104 Publication	<b>1%</b>
<b>4</b>	Suciati, James A. Fraser, Lynette K. Lambert, Gregory K. Pierens, Paul V. Bernhardt, Mary J. Garson. " Secondary Metabolites of the Sponge-Derived Fungus ", Journal of Natural Products, 2013 Publication	<b>1%</b>
<b>5</b>	<a href="http://www.scribd.com">www.scribd.com</a> Internet Source	<b>1%</b>

- 6 Rudiyansyah, , and Mary J. Garson. "Secondary Metabolites from the Wood Bark of *Durio zibethinus* and *Durio kutejensis*", Journal of Natural Products, 2006. 1%
- Publication
- 
- 7 Patricia Y. Hayes. "Complete 1H and 13C assignments of the four major saponins from *Dioscorea villosa* (wild yam)", Magnetic Resonance in Chemistry, 11/2007 1%
- Publication
- 
- 8 diyhpl.us 1%
- Internet Source
- 
- 9 Ping-Chung Kuo, Hsin-Yi Hung, Chi-Wei Nian, Tsong-Long Hwang et al. " Chemical Constituents and Anti-inflammatory Principles from the Fruits of ", Journal of Natural Products, 2017 1%
- Publication
- 
- 10 Muriel Cuendet. "Iridoid Glucosides with Free Radical Scavenging Properties from *Fagraea blumei*", Helvetica Chimica Acta, 06/30/1997 1%
- Publication
- 
- 11 Henrik Franzyk. "SYNTHESIS OF NOVEL HYDROXYMETHYL SUBSTITUTED ANALOGUES RELATED TO CARBOVIR AND NEPLANOCIN A", Nucleosides Nucleotides & Nucleic Acids, 2002 1%

- 12** Adib Charafeddine. "First Synthesis of 2'-Deoxyfluoropuromycin Analogues: Experimental Insight into the Mechanism of the Staudinger Reaction", *Chemistry - A European Journal*, 06/25/2007  
Publication <1%
- 
- 13** Waffo-Teguo, P.. "Acetylated glucuronide triterpene bidesmosidic saponins from *Symplocos glomerata*", *Phytochemistry*, 200403  
Publication <1%
- 
- 14** Ch. Turghun, M. Bakri, G. A. Zou, Kh. M. Bobakulov, H. A. Aisa. "Phenolic Compounds from Leaves of *Nitraria sibirica*", *Chemistry of Natural Compounds*, 2018  
Publication <1%
- 
- 15** Essam Abdel-Sattar. "Saponin Glycosides from *Osteospermum vaillantii*", *Pharmaceutical Biology (Formerly International Journal of Pharmacognosy)*, 12/1/2001  
Publication <1%
- 
- 16** Lawrence Onyango Arot Manguro. "Phenolics of *Moringa oleifera* leaves", *Natural Product Research*, 01/2007  
Publication <1%

Hai-Feng Chen. "Novel furostanol saponins

- 17 from the bulbs of *Allium macrostemon* B. and their bioactivity on  $[Ca^{2+}]_i$  increase induced by KCl", *Journal of Asian Natural Products Research*, 1/1/2006  
Publication <1%
- 
- 18 Nicholas J. Matovic. "The Truth about False Unicorn (*Chamaelirium luteum*): Total Synthesis of 23R,24S-Chiograsterol B Defines the Structure and Stereochemistry of the Major Saponins from this Medicinal Herb", *Chemistry - A European Journal*, 06/27/2011  
Publication <1%
- 
- 19 Ken W.L. Yong, Angela A. Saïim, Mary J. Garson. "New oxygenated diterpenes from an Australian nudibranch of the genus *Chromodoris*", *Tetrahedron*, 2008  
Publication <1%
- 
- 20 AKDEMİR, Zeliha Ş., TATLI, İ. İrem, BEDİR, Erdal and KHAN, Ikhlas A.. "Iridoid and phenylethanoid glycosides from *Verbascum lasianthum*", *TUBITAK*, 2004.  
Publication <1%
- 
- 21 aem.asm.org  
Internet Source <1%
- 
- 22 Chans, Guillermo M., Antonio Nieto-Camacho, Teresa Ramirez-Apan, Simón Hernández-Ortega, Cecilio Álvarez-Toledano, and <1%

Elizabeth Gomez. "Synthetic, Spectroscopic, Crystallographic, and Biological Studies of Seven-Coordinated Diorganotin(IV) Complexes Derived from Schiff Bases and Pyridinic Carboxylic Acids", Australian Journal of Chemistry, 2015.

Publication

---

**23** Tan Pei Jean. "Bidesmosidic Oleanane Saponins from *Xerospermum noronhianum*", Helvetica Chimica Acta, 10/2009 <1%

Publication

---

**24** van Wyk, A.W.W.. "Semisynthesis of labdane diterpene metabolites from the nudibranch *Pleurobranchaea meckelii*", Tetrahedron, 20071203 <1%

Publication

---

**25** Dinda, Biswanath, Sudhan Debnath, and Yoshihiro Harigaya. "Naturally Occurring Iridoids. A Review, Part 1", CHEMICAL & PHARMACEUTICAL BULLETIN, 2007. <1%

Publication

---

**26** Dinda, Biswanath, Debashis Roy Chowdhury, and Bikas Chandra Mohanta. "Naturally Occurring Iridoids, Secoiridoids and Their Bioactivity. An Updated Review, Part 3", CHEMICAL & PHARMACEUTICAL BULLETIN, 2009. <1%

Publication



- 
- 27** Jun Uzawa. "Pregnane glycosides from the stems of *Marsdenia tenacissima*", *Journal of Asian Natural Products Research*, 6/1/2004  
Publication <1 %
- 
- 28** Mudianta, I Wayan, Tina Skinner-Adams, Katherine T. Andrews, Rohan A. Davis, Tri A. Hadi, Patricia Y. Hayes, and Mary J. Garson. "Psammaphysin Derivatives from the Balinese Marine Sponge *Aplysinella strongylata*", *Journal of Natural Products*, 2012.  
Publication <1 %
- 
- 29** Fuad O. Abdullah, Faiq H. S. Hussain, Marco Clericuzio, Alessio Porta, Giovanni Vidari. "A New Iridoid Dimer and Other Constituents from the Traditional Kurdish Plant ", *Chemistry & Biodiversity*, 2017  
Publication <1 %
- 
- 30** Xiao-Jing Wang, Gui-Jie Zhang, Peng-Yu Zhuang, Yan Zhang et al. " Lycojaponicumins A–C, Three Alkaloids with an Unprecedented Skeleton from ", *Organic Letters*, 2012  
Publication <1 %
- 
- 31** Riviere, C.. "Chemotaxonomic interest of iridoids isolated from a Malagasy species: *Perichlaena richardii*", *Biochemical Systematics and Ecology*, 201108/12  
Publication <1 %
-

- 32** Gangwar, Mayank Goel, R.K. Nath, Gopal. "Mallotus philippinensis Muell. Arg (Euphorbiaceae): ethnopharmacology and phytochemistry review.(Rep", BioMed Research International, Annual 2014 Issue  
Publication <1%
- 
- 33** scmb.uq.edu.au  
Internet Source <1%
- 
- 34** U. R. Mikkelsen. "Prostaglandin synthesis can be inhibited locally by infusion of NSAIDS through microdialysis catheters in human skeletal muscle", Journal of Applied Physiology, 11/21/2007  
Publication <1%
- 
- 35** Shao-Jiang Song, Ling-Zhi Li, Pin-Yi Gao, Ying Peng, Jing-Yu Yang, Chun-Fu Wu. "Terpenoids and hexenes from the leaves of Crataegus pinnatifida", Food Chemistry, 2011  
Publication <1%
- 
- 36** Liqin Ding, Feng Zhao, Lixia Chen, Zhihu Jiang, Yue Liu, Zhuomin Li, Feng Qiu, Xinsheng Yao. "New monoterpene glycosides from Paeonia suffruticosa Andrews and their inhibition on NO production in LPS-induced RAW 264.7 cells", Bioorganic & Medicinal Chemistry Letters, 2012  
Publication <1%
- 
- 37** Elmi N. Zainuddin, Renate Mentel, Victor Wray,

---

Rolf Jansen, Manfred Nimtz, Michael Lalk, Sabine Mundt. "Cyclic Depsipeptides, Ichthyopeptins A and B, from ", Journal of Natural Products, 2007

Publication

<1%

38

Gang Ding, Shuchun Liu, Liangdong Guo, Yuguang Zhou, Yongsheng Che. "Antifungal Metabolites from the Plant Endophytic Fungus ", Journal of Natural Products, 2008

Publication

<1%

39

De Rosa, S.. "Minor cacospongionolide derivatives from the sponge *Fasciospongia cavernosa*", Tetrahedron, 20070226

Publication

<1%

40

[www.tno.co.jp](http://www.tno.co.jp)

Internet Source

<1%

41

Jin Li, Tao Wang, Peng Yu, A. Peterson, R. Weber, D. Soerens, D. Grubisha, D. Bennett, J. M. Cook. "General Approach for the Synthesis of Ajmaline/Sarpagine Indole Alkaloids: Enantiospecific Total Synthesis of (+)-Ajmaline, Alkaloid G, and Norsuaveoline via the Asymmetric Pictet-Spengler Reaction", Journal of the American Chemical Society, 1999

Publication

<1%

42

Lee, M.H.. "Bioactive constituents of *Spatholobus suberectus* in regulating

<1%

tyrosinase-related proteins and mRNA in HEMn cells", *Phytochemistry*, 200606  
Publication

- 
- 43 Ciavatta, M.L.. "Studies on puupehenone-metabolites of a *Dysidea* sp.: structure and biological activity", *Tetrahedron*, 20070205  
Publication <1%
- 
- 44 Naima Boutaghane, Laurence Voutquenne-Nazabadioko, Dominique Harakat, Alain Simon, Zahia Kabouche. "Triterpene saponins of *Genista ulicina* Spach", *Phytochemistry*, 2013  
Publication <1%
- 
- 45 Shu-hua Qi. "The chemical constituents of *Munronia Henryi*", *Journal of Asian Natural Products Research*, 7/1/2003  
Publication <1%
- 
- 46 Hernandez-Guerrero, C.J.. "Sesterterpene metabolites from the sponge *Hyatella intestinalis*", *Tetrahedron*, 20060605  
Publication <1%
- 
- 47 [royalgrancanaria.nl](http://royalgrancanaria.nl)  
Internet Source <1%
- 
- 48 [eprints.soton.ac.uk](http://eprints.soton.ac.uk)  
Internet Source <1%
- 
- 49 Min Xu. "Iridoidal glucosides from *Gentiana*

rhodantha", Journal of Asian Natural Products  
Research, 6/2008

Publication

<1%

50

Yoko Shimokawa, Yusuke Akao, Yusuke  
Hirasawa, Khalijah Awang et al. " Gneyulins A  
and B, Stilbene Trimers, and Noidesols A and  
B, Dihydroflavonol- -Glucosides, from the Bark  
of ", Journal of Natural Products, 2010

Publication

<1%

51

Marcos, I.S.. "Synthetic studies towards  
picrasane quassinoids", Tetrahedron, 20070305

Publication

<1%

52

"Phytochemicals in Human Health Protection,  
Nutrition, and Plant Defense", Springer Nature,  
1999

Publication

<1%

53

Liang, Jinru, Yoichiro Ito, Xinxin Zhang, Jiao  
He, and Wenji Sun. "Rapid preparative  
separation of six bioactive compounds from  
*Gentiana crassicaulis* Duthie ex Burk. using  
microwave-assisted extraction coupled with  
high-speed counter-current chromatography",  
Journal of Separation Science, 2013.

Publication

<1%

54

Ken W.L. Yong, Aroon Jankam, John N.A.  
Hooper, Apichart Suksamrarn, Mary J. Garson.  
"Stereochemical evaluation of sesquiterpene

<1%

quinones from two sponges of the genus Dactylospongia and the implication for enantioselective processes in marine terpene biosynthesis", Tetrahedron, 2008

Publication

55

Feng Zhang, Zhi-Jun Wu, Lian-na Sun, Jun Wang, Xia Tao, Wan-Sheng Chen. "Iridoid glucosides and a C13-norisoprenoid from Lamiophlomis rotata and their effects on NF- $\kappa$ B activation", Bioorganic & Medicinal Chemistry Letters, 2012

<1%

Publication

56

Li, S.H.. "Novel taxoids from the Chinese yew Taxus yunnanensis", Tetrahedron, 20030101

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Phytochemical Study of Fagraea spp. Uncovers a New Terpene Alkaloid with Anti-Inflammatory Properties

GRADEMARK REPORT

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6