

DE GRUYTER

2018 · VOLUME 29 · ISSUE 6

ISSN 0792-4855 · e-ISSN 2791-0296

JOURNAL OF BASIC AND CLINICAL PHYSIOLOGY AND PHARMACOLOGY

EDITOR-IN-CHIEF
Michal Horowitz

DE
G

www.degruyter.com/jbcpp



Journal of Basic and Clinical Physiology and Pharmacology

Editor In Chief: Michal Horowitz

Editorial Board: Karen Avraham, Kusal K. Das, Yoram Epstein, Elliot S. Gershon MD, Einat Kodesh, Ron Kohen, David Lichtstein, Alina Maloyan, Raphael Mechoulam, Joachim Roth, Suzanne Schneider, Esther Shohami, Haim Sohmer, Toshikazu Yoshikawa and Joseph Tam



OVERVIEW LATEST ISSUE ISSUES SUBMIT **EDITORIAL**

Editor-in-Chief

Michal Horowitz

m.horowitz@mail.huji.ac.il

Editorial Board

Karen Avraham, Tel Aviv University, Tel Aviv, Israel

Kusal Das, BLDE University, Bijapur, India

Yoram Epstein, Tel Aviv, Tel Aviv University, Israel

Elliot S. Gershon, The University of Chicago Medicine, Chicago, IL

Einat Kodesh, University of Haifa, Haifa, Israel

Ron Kohen, Jerusalem, Hebrew University of Jerusalem, Israel

David Lichtstein, Hebrew University Jerusalem, Jerusalem, Israel

Alina Maloyan, University of Texas Health Science Center, San Antonio, TX

Raphael Mechoulam, The Hebrew University Jerusalem, Jerusalem, Israel

Joachim Roth, Justus-Liebig-University Giessen, Giessen, Germany

Suzanne Schneider, University of New Mexico, Albuquerque, NM

Esther Shohami, Hebrew University of Jerusalem, Jerusalem, Israel

Haim Sohmer, Hebrew University of Jerusalem, Jerusalem, Israel

Joseph Tam, Hebrew University of Jerusalem, Jerusalem, Israel

Toshikazu Yoshikawa, Kyoto Prefectural University of Medicine, Kyoto, Japan

Editorial Office

Katharina Appelt

DE
GUYTER

DE GRUYTER

SUBJECT AREAS

PUBLICATIONS

OPEN ACCESS

SERVICES

ABC

E-mail: jocpp.editorial@degruyter.com

IMPRINTS

Birkhäuser

De Gruyter Mouton

De Gruyter Oldenbourg

De Gruyter Saur

Deutscher Kunstverlag

Edition Klaus Schwarz

Jovis

CONTACT & SERVICE

Help Center

Contact

Career

Our Locations

Imprint

STAY IN TOUCH



Blogs

Newsletter



Journal of Basic and Clinical Physiology and Pharmacology

Editor-in-Chief: Horowitz, Michal

Editorial Board: Avraham , Karen / Das, Kusal K. / Epstein, Yoram / S. Gershon MD, Elliot / Kodesh , Einat / Kohen, Ron / Lichtstein, David / Maloyan, Alina / Mechoulam, Raphael / Roth, Joachim / Schneider, Suzanne / Shohami, Esther / Sohmer, Haim / Yoshikawa, Toshikazu / Tam, Joseph

CiteScore 2018: 1.66

SCImago Journal Rank (SJR) 2018: 0.434

Source Normalized Impact per Paper (SNIP) 2018: 0.699



Online

ISSN

2191-0286

[See all formats and pricing](#)

[Select Volume and Issue](#)

Volume 30, Issue 6 (Nov 2019)

Special Conference Issue: 2nd International Graduate Student Conference on Pharmaceutical Sciences (IGSCPS)

Reviews

[Pharmacist contributions in the treatment of diabetes mellitus in Southeast Asia: a narrative review](#)

Dwiputri, Ayu Wulan / Pristianty, Liza / Hermansyah, Andi

Published Online: 01/23/2020

- [Get Access to Full Text](#)

[The implementation of a chronic disease management program \(Prolanis\) in Indonesia: a literature review](#)

Rachmawati, Sesty / Prihhastuti-Puspitasari, Hanni / Zairina, Elida

Published Online: 12/20/2019

- [Get Access to Full Text](#)

Original Articles

[Quercetin attenuates acute predator stress exposure-evoked innate fear and behavioral perturbation](#)

Anggreini, Putri / Ardianto, Chrismawan / Rahmadi, Mahardian / Khotib, Junaidi

Published Online: 11/28/2019

- [Get Access to Full Text](#)

[Evaluating current practices and policies in the use of injectable medicines for treating myalgia in a primary care center in Pamekasan, Indonesia](#)

Prasetio, Eko / Utami, Wahyu / Othman, Zulhabri / Wardani, Ari / Rahem, Abdul / Hermansyah, Andi

Published Online: 02/07/2020

- [Get Access to Full Text](#)

[The contemporary role and potential of pharmacist contribution for community health using social media](#)

Hermansyah, Andi / Sukorini, Anila Impian / Asmani, Fadli / Suwito, Kandi Aryani / Rahayu, Titik Puji

Published Online: 12/04/2019

- [Get Access to Full Text](#)

[Evaluation of rational drug use based on World Health Organization prescribing indicators in a primary care center in Pamekasan East Java, Indonesia](#)

Prasetio, Eko / Utami, Wahyu / Othman, Zulhabri / Wardani, Ari / Rahem, Abdul / Hermansyah, Andi

Published Online: 02/07/2020

- [Get Access to Full Text](#)

[The prescription patterns of second-generation antipsychotics in schizophrenia outpatient setting](#)

Julaeha, Julaeha / Athiyah, Umi / Hermansyah, Andi

Published Online: 12/14/2019

- [Get Access to Full Text](#)

Factors affecting community pharmacist's service for women with chronic diseases during pregnancy and breastfeeding: application of the Health Belief Model

Anggraini, Septi / Utami, Wahyu / Zairina, Elida

Published Online: 12/19/2019

- [Get Access to Full Text](#)

Overview of therapeutic changes in antiepileptic drugs in adult patients

Ratri, Dinda M.N. / Rahmadi, Mahardian / Islamiyah, Wardah R. / Harum, Nur Faridah

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Exploration of barriers affecting job satisfaction among community pharmacists

Rijaluddin, Muhammad Khalid / Utami, Wahyu / Othman, Zulhabri / Puspitasari, Hanni Prihhastuti / Rahem, Abdul / Sukorini, Anila Impian / Hermansyah, Andi

Published Online: 01/13/2020

- [Get Access to Full Text](#)

Patients' characteristics and their adherence to insulin therapy

Fitriani, Yunti / Pristianty, Liza / Hermansyah, Andi

Published Online: 01/15/2020

- [Get Access to Full Text](#)

Correlation of chemotherapy costs with quality of life in nasopharyngeal cancer patients

Romdhoni, Achmad Chusnu / Aulia, Riskha / Utaminingtyas, Ririn Prasetyo / Suharjono / Alderman, Christopher Paul

Published Online: 01/23/2020

- [Get Access to Full Text](#)

Impact of educational preeclampsia prevention booklet on knowledge and adherence to low dose aspirin among pregnant women with high risk for preeclampsia

Karunia, Rella Indah / Purnamayanti, Anita / Prasetyadi, Fransiscus O.H.

Published Online: 01/13/2020

- [Get Access to Full Text](#)

Translation, cultural adaptation, and validation of the quality of well being self-administered questionnaire in general population in Indonesia

Andayani, Tri Murti / Kristina, Susi Ari / Endarti, Dwi

Published Online: 12/20/2019

- [Get Access to Full Text](#)

Knowledge, attitude, and practice of pharmacists towards management of hypertension in primary care centers

Wijaya, I Nyoman / Athiyah, Umi / Fasich / Hermansyah, Andi

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Medication adherence in diabetes mellitus patients at Tanjung Karang Primary Health Care Center, Mataram

Andanalusia, Mahacita / Athiyah, Umi / Nita, Yunita

Published Online: 12/18/2019

- [Get Access to Full Text](#)

Decreasing angiogenesis vasa vasorum through Lp-PLA2 and H2O2 inhibition by PSP from Ganoderma lucidum in atherosclerosis: in vivo diabetes mellitus type 2

Wihastuti, Titin Andri / Amiruddin, Reyhan / Cesa, Fibe Yulinda / Alkaf, Amalia Istiqamah / Setiawan, Meddy / Heriansyah, Teuku

Published Online: 02/07/2020

- [Get Access to Full Text](#)

Antiaggregation effect of clopidogrel in coronary heart disease patients using omeprazole

Munthe, Dian Hasiannami Boru / Sargo, Siti Sjamsiah / Yogiarto, Mohammad

Published Online: 01/25/2020

- [Get Access to Full Text](#)

The effect of curcuma (*Curcuma xanthorrhiza roxb.*) extract as an adjuvant of captopril therapy on cardiac histopathology of male mice (*Mus musculus*) with hypertension

Hijriani, Nursela / Yusetyani, Lilik / Hasmono, Didik

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Coenzyme Q10 nanostructured lipid carriers as an inducer of the skin fibroblast cell and its irritability test in a mice model

Shoviantari, Fenita / Erawati, Tristiana / Soeratri, Widji

Published Online: 12/20/2019

- [Get Access to Full Text](#)

Medical problems in patients with chronic kidney disease undergoing hemodialysis and their therapy

Suprapti, Budi / Nilamsari, Wenny Putri / Rachmania / Widodo / Alderman, Chris

Published Online: 11/20/2019

- [Get Access to Full Text](#)

ADMET properties of novel 5-O-benzoylpinostrobin derivatives

Pratama, Mohammad Rizki Fadhil / Poerwono, Hadi / Siswodiharjo, Siswandono

Published Online: 12/18/2019

- [Get Access to Full Text](#)

Development of nonalcoholic fatty liver disease model by high-fat diet in rats

Wardani, Hijrawati Ayu / Rahmadi, Mahardian / Ardianto, Chrismawan / Balan, Santhra Segaran / Kamaruddin, Norshafarina Shari / Khotib, Junaidi

Published Online: 11/25/2019

- [Get Access to Full Text](#)

Molecular docking of novel 5-O-benzoylpinostrobin derivatives as wild type and L858R/T790M/V948R mutant EGFR inhibitor

Pratama, Mohammad Rizki Fadhil / Poerwono, Hadi / Siswodihardjo, Siswandono

Published Online: 12/19/2019

- [Get Access to Full Text](#)

The relationship between the level of education and accuracy of insulin injection techniques in DM patients with measurement of HbA1c values

Achmad, Anisyah / Sholihah, Fatchur Rohmi Latifatus / Oktavianti, Wanda Fenny / Sasiarini, Laksmi

Published Online: 01/25/2020

- [Get Access to Full Text](#)

The effect of premixed insulin to blood glucose concentration in patients with type 2 diabetes mellitus

Puspitasari, Arina D. / Kusuma, Hayu / Ratri, Dinda M.N. / Wibisono, Cahyo / Suprapti, Budi

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Intravenous insulin therapy in diabetes mellitus with hyperglycemic crisis and intercurrent illness

Suprapti, Budi / Syarfina, Fairuza / Ardianto, Chrismawan / Wibisono, Cahyo

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Adherence behavior assessment of oral antidiabetic medication use: a study of patient decisions in long-term disease management in primary health care centers in Surabaya

Aditama, Lisa / Athiyah, Umi / Utami, Wahyu / Rahem, Abdul

Published Online: 01/18/2020

- [Get Access to Full Text](#)

Evaluation to the chemotherapy use in patients with diffuse large B-cell lymphoma

Dirani, Dirani / Suharjono / Sedana, Made / Wahyuni, Siti / Ardianto, Chrismawan / Alderman, Chris

Published Online: 01/13/2020

- [Get Access to Full Text](#)

Analysis of the use and cost of stress ulcer prophylaxis for surgical inpatients

Wijaya, Dhani / Padolo, Elfri / Ardianto, Chrismawan / Sumarno / Matulatan, Fendy / Alderman, Chris / Suharjono

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Antineuroinflammation activity of n-butanol fraction of Marsilea crenata Presl. in microglia HMC3 cell line

Ma'arif, Burhan / Mirza, Denis Mery / Hasanah, Mu'akibatul / Laswati, Hening / Agil, Mangestuti

Published Online: 01/22/2020

- [Get Access to Full Text](#)

The enhancement of Arg1 and activated ER β expression in microglia HMC3 by induction of 96% ethanol extract of Marsilea crenata Presl. leaves

Ma'arif, Burhan / Agil, Mangestuti / Laswati, Hening

Published Online: 01/25/2020

- [Get Access to Full Text](#)

Ternary solid dispersion to improve solubility and dissolution of meloxicam

Setyawan, Dwi / Dewi, Meivita Yusmala / Isadiartuti, Dewi

Published Online: 12/14/2019

- [Get Access to Full Text](#)

Improving solubility and dissolution of meloxicam by solid dispersion using hydroxypropyl methylcellulose 2910 3 cps and nicotinamide

Fathanah, Ana / Setyawan, Dwi / Sari, Retno

Published Online: 12/14/2019

- [Get Access to Full Text](#)

o-Hydroxycinnamic derivatives as prospective anti-platelet candidates: in silico pharmacokinetic screening and evaluation of their binding sites on COX-1 and P2Y12 receptors

Nofianti, Kholis Amalia / Ekowati, Juni

Published Online: 12/19/2019

- [Get Access to Full Text](#)

The change of proinflammatory cytokine tumor necrosis factor α level in the use of meloxicam in rat model of osteoarthritis

Khotib, Junaidi / Utami, Naning Windi / Gani, Maria Apriliani / Ardianto, Chrismawan

Published Online: 12/14/2019

- [Get Access to Full Text](#)

Attenuation of IL-1 β on the use of glucosamine as an adjuvant in meloxicam treatment in rat models with osteoarthritis

Khotib, Junaidi / Pratiwi, Asri Putri / Ardianto, Chrismawan / Rahmadi, Mahardian

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Analysis of effectiveness and drug related problems of pain reliever for knee osteoarthritis: weighing clinical risk and benefit

Khotib, Junaidi / Setiawan, Henny Utami / Nurhan, Ahmad Dzulfikri / Rahadiansyah, Erreza / Ardianto, Chrismawan / Rahmadi, Mahardian

Published Online: 02/11/2020

- [Get Access to Full Text](#)

Comparison of antibiotic prescriptions in adults and children with upper respiratory tract infections in Bangka Tengah primary health care centers

Novan Y. I., Pratama / Primadi, Avianto / Mahfudz / Suharjono

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Profile of sociodemographics, sources of infection, antiretrovirals and CD4 counts on HIV/AIDS outpatients in Turen Primary Health Centre, Indonesia

Dhini, Ellyvina S. / Setiadi, Antonius Adji P. / Wibowo, Yosi I.

Published Online: 12/19/2019

- [Get Access to Full Text](#)

Synergistic anti-hepatitis C virus activity of Ruta angustifolia extract with NS3 protein inhibitor

Wahyuni, Tutik Sri / Mahfud, Humairoh / Permatasari, Adita Ayu / Widyawaruyanti, Aty / Fuad, Achmad

Published Online: 12/14/2019

- [Get Access to Full Text](#)

In vitro equivalence of generic and branded amoxicillin tablet by microbiological assay method

Avianto, Primadi / Mahfudz / Suharjono / Isnaeni / Alderman, Christopher Paul

Published Online: 01/11/2020

- [Get Access to Full Text](#)

Knowledge and attitude: two fundamental factors that determine patient compliance in antibiotic therapy

Pristianty, Liza / Kurniati, Vivi Laily / Hidayati, Ika Ratna

Published Online: 02/12/2020

- [Get Access to Full Text](#)

Molecular docking study of sappan wood extract to inhibit PBP2A enzyme on methicillin-resistant Staphylococcus aureus (MRSA)

Gondokesumo, Marisca Evalina / Kurniawan, Ihsan Mulyadi

Published Online: 01/18/2020

- [Get Access to Full Text](#)

Effect of curcumin analogue synthetic product from cullilawan oil for the liver damage treatment in male mice (*Mus musculus* L.)

Kapelle, Imanuel Berly Delvis / Manalu, Wasmen / Souhoka, Fensia Analda

Published Online: 01/23/2020

- [Get Access to Full Text](#)

Case Report

A case report of generalized tetanus in a 42-year-old man with dental infection

Ulfa, Maria / Husna, Nuril Auliya

Published Online: 12/07/2019

- [Get Access to Full Text](#)

Access brought to you by: Airlangga University Library (UNAIR)

Copyright © 2011–2020 by Walter de Gruyter GmbH

[Powered by PubFactory](#)

also developed by scimago:  SCIMAGO INSTITUTIONS RANKINGS



Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

- Home
- Journal Rankings
- Country Rankings
- Viz Tools
- Help
- About Us

Journal of Basic and Clinical Physiology and Pharmacology

Country [Germany - IIII](#) [SJR Ranking of Germany](#)

Subject Area and Category [Biochemistry, Genetics and Molecular Biology](#)
[Physiology](#)

[Medicine](#)
[Medicine \(miscellaneous\)](#)

[Pharmacology, Toxicology and Pharmaceutics](#)
[Drug Discovery](#)
[Pharmacology](#)

Publisher [Walter de Gruyter GmbH](#)

Publication type Journals

ISSN 21910286, 07926855

Coverage 1985-1988, 1990-ongoing

30

H Index

Scope The Journal of Basic and Clinical Physiology and Pharmacology (JBCPP) is a peer-reviewed bi-monthly published journal in experimental medicine. JBCPP publishes novel research in the physiological and pharmacological sciences, including brain research; cardiovascular-pulmonary interactions; exercise; thermal control; haematology; immune response; inflammation; metabolism; oxidative stress; and phytotherapy. As the borders between physiology, pharmacology and biochemistry become increasingly blurred, we also welcome papers using cutting-edge techniques in cellular and/or molecular biology to link descriptive or behavioral studies with cellular and molecular mechanisms underlying the integrative processes. Topics: Behavior and Neuroprotection, Reproduction, Genotoxicity and Cytotoxicity, Vascular Conditions, Cardiovascular Function, Cardiovascular-Pulmonary Interactions, Oxidative Stress, Metabolism, Immune Response, Hematological Profile, Inflammation, Infection, Phytotherapy.

- [Homepage](#)
- [How to publish in this journal](#)
- [Contact](#)
- [Join the conversation about this journal](#)


Online Conference

Call for participation to 23rd International Conference on Multidisciplinary Studies

euser.org

OPEN

Quartiles



SJR



Citations per document



0.75



1.8




Online Conference

Call for participation to 23rd International Conference on Multidisciplinary Studies

OPEN



Unfortunately, we cannot help you with your request, we suggest you to visit the journal's homepage or contact the journal's editorial staff, so they could inform you more deeply.



I couldn't find how to publish the article at this journal. Could you possibly send the requirements of the journal, please?

Journal of Basic and Clinical Physiology and...

Q3 Drug Discovery
best quartile

SJR 2018
0.43

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com/journalsearch.php?q=22980&tip=sid&tip=sid">Journal of Basic and Clinical Physiology and Pharmacology</a>
```



Melanie Ortiz 5 months ago

SCImago Team

Dear Nilufar,

You can see the updated information just above. Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®



Fenita Shoviantari¹ / Tristiana Erawati^{2,3} / Widji Soeratri³

Coenzyme Q10 nanostructured lipid carriers as an inducer of the skin fibroblast cell and its irritability test in a mice model

¹ Department of Industrial Pharmacy, Institut Ilmu Kesehatan Bhakti Wiyata Kediri, Kota Kediri, Jawa Timur 64114, Indonesia. <https://orcid.org/0000-0002-2020-7098>.

² Universitas Airlangga, Department of Pharmaceutic, Faculty of Pharmacy, Gedung Nanizar Zaman Joenoes, UNAIR Kampus C, Jl Mulyorejo 60115 Surabaya, E-mail: era_ffua@yahoo.co.id

³ Department of Pharmaceutics, Airlangga University, Jl. Mulyorejo Surabaya, Jawa Timur, Indonesia, E-mail: era_ffua@yahoo.co.id

Abstract:

Background: Coenzyme Q10 is a fat-soluble antioxidant that can help to prevent collagen and elastin damage and avoid wrinkles. Coenzyme Q10 has several disadvantages to be formulated in topical dosage forms, such as low water solubility and large molecular weight. These make coenzyme Q10 retained in the stratum corneum and cause low skin penetration, so proper formulation is required to get products that can penetrate the skin layer. A nanostructured lipid carrier (NLC) consists of a matrix of solid lipids and liquid lipids in a certain amount with nanoparticle size; it may help increase the penetration of active ingredients.

Methods: For the antiaging activity test, mice were grouped into four treatment groups and killed on the 14th day; then the back of the skin was stained with Masson trichrome staining. For the irritation test, the mice were grouped into three groups and killed after 24 h; then the back of the mice was stained with hematoxylin-eosin staining.

Results: The number of fibroblasts in mice with NLC coenzyme Q10 is highest from all test groups. The irritation test results after 24 h of application preparation showed that NLC coenzyme Q10 did not irritate the skin of the back of male mice.

Conclusions: One percent coenzyme Q10 loaded in NLC induced the number of fibroblast cells in the mice model and showed no irritability effect in histopathology preparations.

Keywords: antiaging, antioxidant, collagen, Masson, staining, trichrome

DOI: 10.1515/jbcpp-2019-0320

Received: October 29, 2019; **Accepted:** November 15, 2019

Introduction

UV exposure activated the reactive oxygen species (ROS) that can increase collagen breakdown, decreasing procollagen and hyaluronic acid production in the skin, which caused wrinkles of skin aging [1]. The widely used therapy to overcome this process is antiaging cosmetics. The commonly used antiaging components are retinoids (vitamin A) and their derivatives, vitamin C, vitamin E, coenzyme Q10, α lipoic acid, and vitamin B3 [2], [3], and most of them are antioxidants. Coenzyme Q10 or ubiquinone is one of antioxidants found in every cell of the body. It can inhibit oxidative stress generated by UV light and it is the first antioxidant that is depleted from the skin. Coenzyme Q10 is an antioxidant that can protect the dermal matrix and fibroblast, and increases the collagen and hyaluronic acid in the skin. Unfortunately, the amount of coenzyme Q10 decreases with age [3], [4]. Coenzyme Q10 is reportedly capable of reducing the production of reactive oxygen species and DNA damage induced by UVA radiation in human keratinocytes *in vitro*. It has been shown to reduce UVA-induced matrix metalloproteinase (MMP) in human dermal fibroblasts. It may inhibit the production of interleukin-6, which stimulates fibroblasts in the dermis by paracrine to regulate MMP production, contribute to protecting the skin fibers from degradation, and rejuvenate the wrinkled skin [5]. Coenzyme Q10 has low water solubility (0.193 $\mu\text{g}/\text{mL}$ in water), large molecular weight (863.36 g/mol), and high lipophilicity ($\log P > 10$) [6], which make coenzyme Q10 poorly penetrate the skin. So coenzyme Q10 requires proper formulation to obtain products with good bioavailability that can penetrate the skin layer [6], [7].

Tristiana Erawati is the corresponding author.
© 2019 Walter de Gruyter GmbH, Berlin/Boston.

Nanolipid carrier systems such as solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) have been developed from nanotechnology-based colloidal systems. The discovery of SLN and NLC has attracted great attention from the pharmaceutical and cosmetics industries. SLN is a spherical particle composed of solid lipids at room temperature with the addition of emulsifier, dissolved in the aqueous phase, while NLC is a refined form of SLN [8]. NLC consists of a matrix of solid lipids and liquid lipids in a certain amount. NLC remains in its solid form by controlling the levels of liquid lipids added to the liquid formulation, so that controlled drug release properties for NLC can be achieved [9]. NLC has a particle size distribution of up to 500 nm. It can reduce the formation of crystals in the lipids and also can increase drug entrapment [10]. The small droplet size of NLC gives a large surface area. It increases the number of active ingredients that can be trapped in it. So the NLC can carry bioactive molecules in a more controlled manner which is expected to release the drug to be greater, and its solubilization power increases [11] and the rate of release (flux) is directly proportional to the concentration so that the greater the active ingredient is trapped in the system, the release flux is also higher [12].

Materials and methods

Preparation of coenzyme Q10 in NLC

This formulation contains 1% coenzyme Q10 as an active ingredient, Tween 80 and Span 80 as surfactants, olive oil as liquid lipid, cetyl palmitate as solid lipid, ethanol 96% as cosurfactant, and acetate buffer pH 4.2 ± 0.2 as the water phase. Solid lipid, liquid lipid, and coenzyme Q10 were melted at 60 °C. While, surfactant and the mixture of cosurfactant and acetate buffer were heated at 60 °C in a different beaker glass. After the solid lipid was completely melted and the temperature reached 60 °C, the mixture of lipids was stirred with Ultra-Turrax High Shear Homogenizer at 5000 rpm; then the surfactant was added and mixed well. Finally, the mixture of cosurfactant and acetate buffer was added drop by drop until the end, rising the speed at 16,000 rpm for 3 min.

Physical quality test of coenzyme Q10 in NLC

Physical quality tested from NLC coenzyme Q10 are particle size and polydispersity index using a Delsa Nano Particle Size Analyzer, pH meter, viscosity using a cone and plate viscometer, entrapment efficiency with a centrifugation method, and droplet shape using a transmission electron microscope (TEM).

Animal preparation

The number of fibroblasts and collagen density were evaluated *in vivo* using an animal model, *Mus musculus*. Male mice weighing 30–40 g were used. Under anesthetization with ketamine intraperitoneal, the hair on the back was cleaned with a mechanical hair clipper approximately 2×2 cm. Each mouse was given a special cage so that it cannot touch the part to be smeared on the sample to be tested. All procedures of this research have been approved by The Animal Care and Use Committee of Universitas Airlangga, Indonesia, with Ethical Clearance number 645-KE.

In vivo skin fibroblast and collagen density testing

The mice were divided into four groups. The first group was mice treated with coenzyme Q10 in NLC to see the effectiveness of the active ingredient in the preparation, and the second group was mice treated with coenzyme Q10 dissolved in olive oil in order to see the activity of coenzyme Q10 dissolved in the oil phase only. The third group was a control group of mice's skin that was not given any treatment to see the condition of normal mice's skin without any treatment. The fourth and last group was mice that were only exposed to UVB light without any treatment to determine the condition of the skin that is exposed to UV light without any treatment. About 50 mg of the sample was applied to the back skin of mice twice a day (except the third group), 20 min before being exposed to UVB for 20 s (to allow time for absorption of topical material into the skin) and 4 h after irradiation (the formation of ROS starting 4 h after exposure). UV irradiation was done every 2 days, namely on days 1, 3, 5, 7, 9, 11, and 13, with the aim to get rid of the effects of acute irradiation. The application of the preparation was still carried out on days without UV irradiation. After 14 days, the mice were killed by

cervical dislocation. The skin was cut using a microtome and then stained with Masson trichrome staining. The preparations were observed under a light microscope.

***In vivo* skin irritability testing**

The mice were divided into three groups. The first group was mice treated with coenzyme Q10 in NLC, the second group was mice treated with coenzyme Q10 dissolved in olive oil, and the last group was a control group of mice's skin that was not given any treatment to see the condition of normal mice's skin without any treatment. Approximately 50 mg of the test sample was applied to the shaved back skin of mice. The mice were killed by dislocation after 24 h. The cut skin part was soaked with formalin solution for further hematoxylin-eosin staining. The preparations were observed under a light microscope. The observation of skin irritation was performed by histopathology scoring.

Results

Physical quality test of coenzyme Q10 in NLC

NLC coenzyme Q10 has particle size $69.7 \pm 0.93 \mu\text{m}$, with polydispersity index 0.35 ± 0.05 , pH value 4.45 ± 0.006 , and viscosity 387 ± 3.55 . The result of the coenzyme Q10 NLC entrapment efficiency test is $74\% \pm 0.87\%$. It showed high entrapment efficiency ($>70\%$). The morphology of NLC coenzyme Q10 particles showed that the system has spherical particles as shown in Figure 1.




Figure 1: NLC coenzyme Q10 particles with a transmission electron microscope.

***In vivo* skin fibroblast and collagen density testing**

Fibroblast and collagen density from mice skin preparation was observed under a light microscope and is shown in Table 1 and Figure 2.




Figure 2: Picture of collagen in histopathology preparation at the back of male mice after 14 days using NLC CoQ10 (A) and the solution of CoQ10 in olive oil (B), also a control group without sample application (C) and a UV control group without sample application (UV). The picture was taken with a light microscope Nikon H600L (scale bar = 100 nm).

Table 1: Result of the number of fibroblasts and density collagen in the back skin of mice after 14 days using NLC CoQ10, and CoQ10 solution in olive oil, the control group of mice without sample application, and the UV control group of mice without sample application.

Group	Number of fibroblast cells	Collagen density
CoQ10 in NLC	94.00 ± 5.13	1.44 ± 0.19
CoQ10 in olive oil	56.87 ± 9.43	1.33 ± 0.33
Control	57.44 ± 23.26	1.44 ± 0.23
UV	30.22 ± 0.51	0.56 ± 0.19

In vivo skin irritability testing

The histopathology scoring of mice is shown in Table 2 and Figure 3.

Table 2: Result of histopathology scoring of mice after 24 h of sample application.

Test parameter	NLC CoQ10	CoQ10 in olive oil	Control
Epidermis			
Liquefaction	0.44 ± 0.49	0.67 ± 0.47	0 ± 0
Subepidermis			
Edema	0.11 ± 0.31	0.44 ± 0.49	0 ± 0
Subepidermis			
Collagen fiber swelling	0 ± 0	0.44 ± 0.49	0 ± 0
Infiltration of inflammatory cell	0.22 ± 0.41	0.11 ± 0.31	0 ± 0
Hypodermis			
Collagen fiber swelling	0 ± 0	0 ± 0	0 ± 0
Infiltration of inflammatory cell	0 ± 0	0 ± 0	0 ± 0
Skin appendages			
Degeneration	0 ± 0	0 ± 0	0 ± 0
Irritation score	0.73 ± 0.51	1.17 ± 0.51	0 ± 0




Figure 3: Microscopic images of mice skin backs after 24 h of coenzyme Q10 application in (A) NLC, (B) olive oil, and (C) controls; pictures were taken using the Nikon H600L light microscope at 100× magnification.

Discussion

NLC consists of solid fat, liquid fat, surfactant, cosurfactant, and water phase. Liquid fat used in this study was olive oil and the solid fat used was cetyl palmitate. There are three largest fatty acid contents in olive oil, namely oleic acid, palmitic acid, and linoleic acid, which in topical preparations can be used as emulsifying agents and enhancers to increase the penetration of transdermal preparations by increasing the bioavailability of water-insoluble drug ingredients and modifying the structure of stratum corneum [13]. To create a stable system, surfactants and cosurfactants are needed in right composition. Surfactants are used to reduce surface tension and interface tension. Based on the head group, surfactants are further divided into four surfactants, namely anionic, cationic, nonionic, and zwitterion surfactants [14]. The NLC formulation was made using a combination of nonionic surfactants, namely Tween 80 and Span 80. Nonionic surfactants were chosen because they are safer than other surfactants in terms of skin irritation [15], [16]. Meanwhile, cosurfactants chosen tend to have shorter C atoms, for example, ethanol, which also has the effect of increasing penetration in transdermal preparations [13]. This short chain alcohol was chosen to obtain NLC with a small particle size.

The size of the coenzyme Q10 NLC particle has been following the requirements of a diameter of 40–1000 nm with the polydispersity index value close to 0, which means homogeneous size. The value of pH NLC coenzyme Q10 was 4.4, which is similar to the pH of the buffer used in this study. The buffer pH 4.4 was used to maintain the stability of the active ingredient. Moreover, the pH value below the pKa of the active ingredient aimed to minimize the ionization of the active ingredient so that the active ingredient would remain in its molecular form and easily penetrate the skin. Besides, pH measurements are also needed to prevent irritation and dryness of the skin. NLC coenzyme Q10 has a pH value of around 4.45 ± 0.006 . The result of the coenzyme Q10 NLC entrapment efficiency test is $74\% \pm 0.87\%$. It showed quite high entrapment efficiency (>70%). The morphology of NLC coenzyme Q10 particles is analyzed using the TEM. Photographs taken with the TEM show that the system has spherical particles, which produce smaller contact area, thus preventing aggregation and friction between each other. From the results of the physical quality test of the preparations, it is known that the preparations have met the requirements for pre-clinical activity testing.

The dermis is the layer that is just below the epidermis layer. It is formed from many amorphous intercellular substances that act as “cement” in all components of the dermis. The main cell in the dermis is fibroblast. This cell produces an intercellular component known as collagen fibers. Collagen and elastin are proteins in the form of fibers. The fibers are intertwined with one another and make the dermis strong and elastic. Collagen fibers provide strength to the skin, whereas elastin fibers thinner than collagen are responsible for skin elasticity

and have the ability to return to their original shape after stretching. If both of these fibers are damaged, for example, due to the influence of age or excessive exposure to sunlight, the skin may become saggy, may not return to its original shape after stretching, and looks thin and wrinkled [17]. The collagen density test with the number of fibroblasts was performed *in vivo* on the skin of the back mice for 14 days. Fibroblasts are the most common cells found in connective tissue and synthesize some components of the extracellular matrix, such as collagen. The more amount of fibroblasts it has, the more collagen it will form. In the calculation of the number of fibroblasts, it was found that the group of mice given NLC coenzyme Q10 had a higher amount of fibroblasts (94.00 ± 5.13) than the group given coenzyme Q10 in olive oil (56.87 ± 9.43), control group (57.44 ± 23.26), and the UV group (30.22 ± 0.51). This suggests that the coenzyme Q10 prepared in the NLC delivery systems has the same effectiveness in the number of fibroblasts and can also increase the number of fibroblasts in the dermis of mice skin. Meanwhile, the dissolved coenzyme Q10 in olive oil has the same amount of fibroblasts as the control group, indicating that coenzyme Q10 is incapable of increasing the amount of fibroblasts if it is not helped by a particular delivery system.

The antiaging activity test was followed by scoring in the collagen density test in the same histopathology preparation with the number of fibroblasts. The collagen density test was performed by histopathology scoring, with detailed scores: score 0 means that the mice's skin has a collagen density of 25%–50%, score 1 means collagen density 50%–75%, and score 2 means collagen density >75%. The results of the histopathology test of the back skin of mice with Masson trichrome dye showed that the group of mice given NLC coenzyme Q10 had a higher collagen density score than the group of mice given coenzyme Q10 solution in olive oil, the group of mice that were not treated (control), and the group of mice which were only exposed to UV rays. From the results of scores on collagen density and microscopic images, it can be seen that mice groups exposed only to UV rays had the lowest scores (0.56 ± 0.20), suggesting that UV rays can damage collagen in the dermis of the skin. This is in contrast to the groups treated with coenzyme Q10 prepared in NLC and dissolved in olive oil; the collagen density of these three groups tends to be the same as the control group. From these results it can be concluded that coenzyme Q10 can protect against the adverse effects of UV rays.

Safety of cosmetics use is one of the purposes for such preparations. One of the safety features of topical preparations can be demonstrated by the absence of skin irritation on use. Examination of skin irritation can be done with histopathological observation. Examination of skin irritation by histopathological observation was carried out by the microscopic tissue scoring study using a light microscope. The results of histopathological scores of the back skin of mice after 24 h of dosage application showed that NLC coenzyme Q10 had an average score of 0.73 ± 0.51 so as not to cause structural changes in the back skin of mice, while coenzyme Q10 dissolved in olive oil had a mean score averaging 1.17 ± 0.51 and the control group had a score of 0 ± 0 . This indicates that the NLC system has less irritation risk than the active ingredient solution in olive oil. This is probably due to the addition of cetyl palmitate, which is a solid fat that has the ability of a moisturizer to minimize the occurrence of irritation.

The results show that the NLC of coenzyme Q10 enhances the quality of the particulate appearance of the dosage form. The *in vivo* test shows that NLC improves the skin fibroblast count, the collagen density, and decreases the irritability of the dosage form as compared to those that are olive oil based. Future plans include carrying out further tests regarding the stability of the preparations and the acceptability assays. The results of this study can be one of the potential antiaging cosmetics that can fight fine lines and wrinkles in a relatively short period of time.

Conclusions

One percent coenzyme Q10 loaded in NLC induced the number of fibroblast cells in the mice model and showed no irritability effect in histopathology preparations.

Acknowledgments

Thanks to Yayasan Bhakti Wiyata who allowed finishing and publishing this research. This paper has been presented at the 8th APPEN Conference and 2nd HPC Conference on 8–9 October 2019, Universitas Airlangga, Surabaya, Indonesia.

Research funding: None declared.

Author contributions: All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

Competing interests: Authors state no conflict of interest.

Ethical approval: All procedure of this research has been approved by The Animal Care and Use Committee of Universitas Airlangga, Indonesia, with Ethical Clearance number 645-KE.

References

- [1] Ganceviciene R, Liakou AI, Theodoridis A, Makrantonaki E, Zouboulis CC. Skin anti-aging strategies. *Dermato-Endocrinol* 2012;4:308–19.
- [2] Burke KE. Implementation of cosmetic dermatology into therapeutic: antiaging regimens. In: Draelos DZ, editor(s). *Cosmetic dermatology products and procedures*. Durham, NC: Wiley Blackwell Publication, 2010:480–7.
- [3] Ramos-e-Silva M, Celem LR, Ramos-e-Silva S, Fucci-da-Costa AP. Anti-aging cosmetics: facts and controversies. *Clin Dermatol* 2013;31:750–8.
- [4] Baumann LS. The Baumann skin typing system. In: Farage M, Miller K, Maibach H. editors. *Textbook of aging skin*. 2nd ed. New York: Springer, Berlin, Heidelberg, 2009:34–41.
- [5] Inui M, Ooe M, Fujii K, Matsunaka H, Yoshida M, Ichihashi M. Mechanisms of inhibitory effects of CoQ10 on UVB-induced wrinkle formation in vitro and in vivo. *BioFactors* 2008;32:237–43.
- [6] Lucangioli S, Tripodi V. The importance of the formulation in the effectiveness of coenzyme Q10 supplementation in mitochondrial disease therapy. *Der Pharmacia Sinica* 2012;3:406–7.
- [7] Korkmaz E, Gokce E, Ozer O. Development and evaluation of coenzyme Q10 loaded solid lipid nanoparticle hydrogel for enhanced dermal delivery. *Acta Pharm* 2013;63:517–29.
- [8] Müller RH, Radtke M, Wissing SA. Solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) in cosmetic and dermatological preparations. *Adv Drug Deliv Rev Adv*. 2002;54:S131–55.
- [9] Puglia C, Blasi P, Rizza L, Schoubben A, Bonina F, Rossi C, et al. Lipid nanoparticles for prolonged topical delivery: an in vitro and in vivo investigation. *Int J Pharm* 2008;357:295–304.
- [10] Phatak AA, Chaudhari PD. Development and evaluation of nanostructured lipid carrier (NLC) based topical delivery of an anti-inflammatory drug. *J Pharm Res* 2013;7:677–85.
- [11] Kogan A, Garti N. Nanoemulsions as transdermal drug delivery vehicles. *Adv Colloid Interface Sci* 2006;123–6:369–85.
- [12] Derle DV. *Essentials of physical pharmacy*. Hyderabad [India]: PharmaMed Press. 2008;73–9.
- [13] Rowe RC, Sheskey PJ, Quinn ME. *Handbook of pharmaceutical excipients*, 6th ed. London: The Pharmaceutical Press, 2009:5–6; 17–9; 75–6; 389–90; 466–8; 470–2; 473–4; 549–53; 675–8.
- [14] Anwar E. *Eksipien dalam Sediaan Farmasi Karakterisasi dan Aplikasi*. Penerbit Dian Rakyat, Jakarta, 2012.
- [15] Kumar B, Jain SK, Prajapati SK, Mahor A, Kumar A. Development and characterization of transdermal microemulsion gel for an antiviral drug. *Int J Pharm Sci Res* 2010;1:57–73.
- [16] Santos P, Watkinson AC, Hadgraft J, Lane ME. Application of microemulsions in dermal and transdermal drug delivery. *Skin Pharmacol Physiol* 2008;21:246–59.
- [17] Shai A, Maibach H, Baran R. Skin structure. *Handbook of cosmetic skin care*. London: Informa Healthcare, 2009:4–13.