

DAFTAR PUSTAKA

1. Butterworth JF, Mackey DC, Wasnick JD. Postanesthesia care. In : Morgan, Mikhail's, eds. Clinical anesthesiology, 5th ed. New York : McGraw-Hill, 2013 :1263.
2. Gan TJ, Sinha AC, Kovac AL, et al. A randomized, double-blind, multicenter trial comparing transdermal scopolamine plus ondansetron to ondansetron alone for the prevention of postoperative nausea and vomiting in the outpatient setting. *Anesth Analg.* 2009;108:1498-504.
3. Ho KY, Gan TJ. Postoperative nausea and vomiting. In : Lobato LB, Gravenstein N, Kirby RR. *Complication in anesthesiology.* Philadelphia : Lippincot Williams & Wilkins; 2008 : 571-80.
4. Wang TF, Liu YH, Chu CC, et al. Low dose haloperidol prevents postoperative nausea and vomiting after ambulatory laparoscopic surgery. *Acta anaesthesiol scand.* 2008;52:280-84.
5. Ismail S. Practice of use of antiemetic in patients for laparoscopic gynaecological surgery and its impact on the early (1st two hrs) postoperative period. *J Pak Med Assoc, Vol.58, No.4, April 2008.*
6. Shafique A, Saleem J, Taqi A, et al. Comparison of dexamethasone and ondansetron for the prophylaxis of post operative nausea and vomiting in laparoscopic gynaecological procedures. *A.P.C.M vol: 6 No. 2 July-December 2012.*
7. Farhat K, Pasha AK, Kazi WA. Comparison of ondansetron and metocloperamide for PONV prophylaxis in laparoscopic cholecystectomy. *J Anesthe Clinic Res.* 2013;4: 297.
8. Gan TJ. Risk factors for postoperative nausea and vomiting. *Anesth Analg.* 2006; 102:1884-98.
9. Le TP, Gan TJ. Update on the management of postoperative nausea and vomiting and postdischarge nausea and vomiting in ambulatory surgery. *Anesthesiology clin* 28. 2010 ;225-249.
10. Hill RP, Lubarsky DA, Phillips BB. Cost-effectiveness of prophylactic antiemetic therapy with ondansetron, droperidol, or placebo. *Anesthesiology.* 2000; 92(4):958-67.
11. Murphy GS, Szokol JW, Greenberg SB, et al. Preoperative Dexamethasone enhances quality of recovery after laparoscopic cholecystectomy: effect on in-hospital and postdischarge recovery outcomes. *Anesthesiology.* 2011 Apr;114(4):882-90.
12. Gan TJ, Meyer T, et al. Consensus guidelines for managing postoperative nausea and vomiting. *Anest analg.* 2003;97: 62-71.
13. Kjelberg FK, Henzi I, Tramer MR. Treatment of established postoperative nausea and vomiting: a quantitative systematic review. *BMC Anesthesiology.* 2001;oct;1:2.
14. Ku CM, Ong BC. Postoperative nausea and vomiting : a review of current literature. *Singapore med J.* 2003; 44 (7): 366-374.
15. Chandrakantan A, Glass PSA. Multimodal therapies for postoperative nausea and vomiting, and pain. *British journal of anaesthesia.* 2011;107 (S1): i27-i40 .

16. Schell RM, Bowe EA. Anesthetics and anesthetic adjuvants. In : Cole DJ, Schlunt M, Hines RL, eds . Adult perioperative anesthesia : the requisites in anesthesiology. Philadelphia: Elsevier Mosby; 2004,144-145.
17. Kusuma SMP. Perbandingan dexametason dengan ondansetron sebagai profilaksis mual dan muntah pasca operasi ginekologi. 2011:48-51.
18. Islam S, Jain PN. Post-operative nausea and vomiting (PONV) : a review article. *Indian J Anaesth.* 2004; 48(4):253-8.
19. Gan TJ. Management of postoperative nausea and vomiting. *The American Society of Anesthesiologists.* 2009; 37(6): 69-80.
20. Azim AASA. Combination of midazolam & dexamethasone in the prophylaxis of ponv is alternative of haloperidol in female patients undergoing gynecological laparoscopic operation. *Ain Shams Journal of Anesthesiology.* 2009; Jan; Vol 2:1-9.
21. Hernandez JG, Alatorre ALO, Contreras MD, et al. Preoperative dexamethasone reduces postoperative pain, nausea and vomiting following mastectomy for breast cancer. *BMC Cancer.* 2010; 10:692.
22. Chaundary S, Sethi AK, Motiani P, et al. Pre-operative intravenous fluid therapy with crystalloids or colloids on postoperative nausea & vomiting. *Indian J Med Res.* 2008;127:577-81.
23. Cheng CR, Sessler DI, Apfel CC. Does neostigmin administration produce a clinically important increase in postoperative nausea and vomiting?. *Anesth Analg.* 2005;101(5):1349-55.
24. Apfel CC, Korttila K, Abdalla M, et al. A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *N Engl J Med.* 2004; 350:2441-51.
25. Thomas R, Jones N. Prospective randomized, double-blind comparative study of dexamethasone, ondansetron, and ondansetron plus dexamethasone as prophylactic antiemetic therapy in patients undergoing day-case gynaecological surgery. *Br J Anaesth.* 2001;87: 588-92.
26. Johnston KD. 5-HT₃-receptor-associated mechanisms of PONV a risk factor targeted approach to anti-emetic therapy. *J Anaesth Clin Pharmacol.* 2010; 26(1):3-10.
27. Stoelting RK. Pharmacology & physiology in anesthetic practice .4 th ed. New York: Lippincott Williams & Wilkins; 2006.
28. Gautam B, Shresta BR, Lama P, et al. Antiemetic prophylaxis against postoperative nausea and vomiting with ondansetron-dexamethasone combination compared to ondansetron alone for patients undergoing laparoscopic cholecystectomy. *Kathmandu University Medical Journal.* 2008; 6(3):319-28.
29. Scuderi PE, James RL, Harris L, et al. Multimodal antiemetic management prevents early postoperative vomiting after outpatient laparoscopy. *Anesth Analg.* 2000;91(6):1408-14.
30. Ananth C. A comparative study of ondansetron and granisetron for prevention of nausea and vomiting following laparoscopic surgeries. 2009; apr: 27-28.
31. Goll V, Akca O, Greiff R, et al. Ondansetron is no more effective than supplemental intraoperative oxygen for prevention of postoperative nausea and vomiting. *Anesth Analg.* 2001;92: 112-7.

32. Sisyanto RDN. Pengaruh peningkatan tekanan intra abdomen padaa pasien operasi laparaskopi terhadap perubahan kadar CO₂ dan laktat dalam darah. 2013;7: 50.