


# CDD 2016

## The 4<sup>th</sup>

# Current Drug Development International Conference 2016

June 1-3, 2016

Phuket Graceland Resort & Spa,  
Patong Beach, Phuket, Thailand



**Proceeding  
The 4th Current Drug Development  
International Conference 2016 (CDD 2016)**

**Editors**

• Juraithip Wungsintaweekul  
Faculty of Pharmaceutical Sciences  
Prince of Songkla University  
Hat Yai, Songkhla, Thailand

• Sirinya Keawnoi

**@Copyright**

Faculty of Pharmaceutical Sciences  
Faculty of Industrial Sciences and Technology  
School of Pharmacy

Prince of Songkla University  
Universiti Malaysia Pahang  
Walailak University

**First edition** 200 copies

**Price** 500 Baht

**National Library of Thailand Cataloging in Publication Data**

Faculty of Pharmaceutical Sciences  
Proceeding the 4<sup>th</sup> Current Drug Development International Conference (CDD 2016)  
Songkhla • 2016 • 252 pages

**ISBN** 978-616-271-303-3

**Design & Printing**

Neo point  
1/59 Ratuhit Road  
Tambol hatyai Amphur hatyai  
Songkhla 90110 Thailand

**Published by Faculty of Pharmaceutical Sciences , Prince of Songkla University**



LG Life Sciences (Thailand) Co., Ltd.  
Farmaline Co., Ltd.  
ITS (Thailand) Co., Ltd.  
Applied Chemical and Instrument Co., Ltd.  
Charpa Techcenter Co., Ltd.  
The Government Pharmaceutical Organization  
Aerocare Co., Ltd.  
Bangkok Lab & Cosmetic Co., Ltd.  
Siamclassic Industry Co., Ltd.  
Thai Can Biotech Co., Ltd.  
Conversant Technology Co., Ltd.  
Abbott Laboratories Co., Ltd.  
RI Technologies Co., Ltd.  
Gold Mints Products Co., Ltd.  
Hollywood International Co., Ltd.  
Spacemed Co., Ltd.  
Haad Thip Public Co., Ltd. (Hatyai Office)  
Innovative Instrument Co., Ltd.  
Bara Scientific Co., Ltd.  
Greater Pharma Co., Ltd.

### Organization Support

- Faculty of Pharmaceutical Sciences, Prince of Songkla University
- Faculty of Industrial Sciences and Technology, Universiti Malaysia Pahang
- Drug and Cosmetic Research and Development Unit, School of Pharmacy, Walailak University
- Drug Delivery System Excellence Center, Prince of Songkla University
- Phytomedicine and Pharmaceutical Biotechnology Excellence Center, Prince of Songkla University

## Organizer Information

### Conference Chair

Pharkphoom Panichayupakaranant

Director of Phytomedicine and Pharmaceutical Biotechnology Excellence Center

### Organizing Committee

Sirirat	Pinsuwan
Pharkphoom	Panichayupakaranant
Teerapol	Srichana
Juraithip	Wungsintaweekul
Sukanya	Dej-adisai
Narubodee	Phadoongsombat
Chitchamai	Ovatlarnporn
Jiraporn	Chingunpitak
Athip	Sakunphueak

### Editor in Chief

Juraithip	Wungsintaweekul
-----------	-----------------

### Reviewers

Apichart	Atipairin	Walailak University, Thailand
Chutima	Jantarat	Walailak University, Thailand
Jiraporn	Chingunpitak	Walailak University, Thailand
Somchai	Sawatdee	Walailak University, Thailand
Brian	Hodgson	Prince of Songkla University, Thailand
Chatchai	Wattanapiromsakul	Prince of Songkla University, Thailand
Juraithip	Wungsintaweekul	Prince of Songkla University, Thailand
Prapaporn	Boonme	Prince of Songkla University, Thailand
Roongnapa	Srichana	Prince of Songkla University, Thailand
Supinya	Tewtrakul	Prince of Songkla University, Thailand
Jianbo	Xiao	University of Macau, China
Mashitah	M Yusoff	Universiti Malaysia Pahang, Malaysia
Mohd Hasbi	Ab. Rahim	Universiti Malaysia Pahang, Malaysia



**Session B: Pharmaceutical Chemistry, Pharmaceutical Technology & Drug Delivery System and Biopharmaceutics**

Paper Code	Paper name	Name	Surname
<b>Pharmaceutical Chemistry</b>			
PB001	Carbon Dioxide Supercritical Fluid Extraction of <i>t</i> -Resveratrol from <i>Arachis hypogaea</i> Using Central Composite Design	Amornrut	Chaidedgumjorn
PB002	Phytochemicals and cytotoxicity studies of <i>Smithatris supraneanae</i> W.J. Kress & K.Larsen	Khanittha	Chawanorasest
PB003	Simultaneous Quantitative HPLC Method for Determination of <i>trans</i> -Resveratrol and Its Glucoside in Thai Beverages	Malai	Satiraphan
PB004	Development of <i>in vitro</i> assay for the detection of fetal hemoglobin inducers	Patamaporn	Pruksakorn
PB005	Synthesis of Benzhydrol and Its Acetamido Analogues as Antituberculosis Agents	Pitikan	Kanjanapruk
PB006	Synthesis of Some Biologically Important Thiazolidinone Compounds	Sedat	Sevmezler
PB007	Antiproliferative compound of <i>Dipterocarpus obtusifolius</i> root	Woranan	Chanchedlha
<b>Pharmaceutical Technology &amp; Drug Delivery System</b>			
PB008	Analytical Method Validation for Determination of Quinine Sulfate in Extemporaneous Oral Suspension	Aitsara	Mitchuayrod
PB009	Development of Lidocaine Mucoadhesive Films	Araya	Raiwa
PB010	Influence of hydrophobicity or hydrophilicity of nano silica in flow property of cohesive API in a powder mixing: a comparative evaluation	Bappaditya	Chatterjee
PB011	Water Absorption and Mucoadhesive Properties of <i>Scaphium macropodum</i> Beaum. (Malva Nut)	Benjabhorn	Sethabouppha

Paper Code	Paper name	Name Surname
PB012	Effect Of Incubation Time On Protein Loading And Encapsulation Efficiency Of Ovalbumin-Alginate Microspheres	Dewi Melani Hariyadi
PB013	Activity of Cyprofloxacin from implant with cross link agent genipin in composites to <i>Staphylococcus aureus</i> ATCC25923 dan <i>Escherichia coli</i> ATCC 25922	Esti Hendradi
PB014	Preparation and Characterization of Chlortetracycline Hydrochloride Solid Dispersions	Jenjira Apiwongngam
PB015	Screening of Cytotoxic and Antioxidant Activities of <i>Jatropha curcas</i> Linn Latex Extracts	kittiya Tinpun
PB016	Formulation Development of Mucoadhesive Microspheres from Extract of <i>Piper betle</i>	Namfa Sermkaew
PB017	Antimicrobial Effect of <i>Piper betle</i> Leaf Extract for Foot Anti-malodor	Natamon Dorkbuakaew
PB018	In vitro Antioxidant Activity of Mouthwash Containing Stingless Bee's Propolis Extract	Natthan Charemsriwilaiwat
✓ PB019	Increasing of Lycopene's Antioxidant Stability In Solid Lipid Nanoparticle (SLN) and Nanostructure Lipid And Carrier (NLC) in Use As Antiaging	Noorma Rosita
PB020	Development of Pluronic Lecithin Organogel loaded <i>Kaempferia parviflora</i> Extract for Transdermal Delivery	Paisit Wattanasri
PB021	Effect of skin penetration enhancers containing liposome on ATRA loading capacity	Ponwanit Charoenputtakun
PB022	Effect of Tween on transport function of organic cation transporter 2	Sirima Soodvilai
PB023	<i>In vitro</i> release kinetic of sildenafil dry foam tablets	Somchai Sawatdee
PB024	Development of Dong quai extracts-loaded solid lipid nanoparticles cream and its preliminary characterization	Somkamon Manchun



## Increasing of Lycopene's Antioxidant Stability In Solid Lipid Nanoparticle (SLN) and Nanostructure Lipid And Carrier (NLC) in Use As Antiaging

**Noorma Rosita, Widji Soeratri**

Faculty of Pharmacy, University of Airlangga

itanr@yahoo.com noorma-r@ff.unair.ac.id

**Abstract-** Lycopene is known as lipophilic antioxidant are often used as an anti-aging cosmetic active ingredient. Antioxidants is easily degraded so that its activity is reduced. To improve the stability, lycopene was loaded in nanolipid structure, such as: Solid Lipid nanoparticles (SLN) and Nanostructure lipid carrier (NLC). Lycopene loaded in SLN and NLC are made with High Shear homogenisation (HSH) method. An antioxidant stability is determined based on IC<sub>50</sub> value changes. IC<sub>50</sub> is determined using 1,1-Diphenyl-2-picrylhydrazyl (DPPH) after being kept in controlled conditions on days 1 and 30. Their oclusivity was determined also. The conclusion is the NLC is able to improve the stability of the lycopene antioxidant and have oclusivity greater than the SLN. Oclusivity nature can indicate the ability of the formula to retain skin moisture.

**Keywords:** Lycopene, Solid Lipid Nanoparticles (SLN) and Nanostructure Lipid Carrier (NLC), setil alcohol, oleic acid.

### Background

Lycopene is a lipophilic antioxidant that easily oxidized. It is necessary to improve the stability of lycopene so that it remains effective. One of the delivery system that proper is nanolipid-based systems, such as Solid Lipid Nanoparticles (SLN) and Nanostructure Lipid Carrier (NLC). The emolien effect of lipid can increase skin softness.

This study compared the SLN and NLC ability in maintaining the lycopene antioxidant stability and also on their oclusivity. Solid lipid used was cetyl alcohol, while the liquid lipid used to form the NLC was oleic acid 1.5%. Oleic acid is a fatty acid with the highest content in olive oil and has penetrant enhacher effect. Systems that do not contain liquid lipid referred to SLN. Furthermore lycopene-NLC and lycopene-SLN stability antioxidant were determined on days 1 and 30 by determining of IC<sub>50</sub> in controlled conditions. Their oclusivity in vitro was also determined. The oclusivity effect related with the ability of retaining skin moisture.

### Materials and Methods

SLN made of lycopene 10 % , 10 % cetyl alcohol , 1.5 % tween 80 , 20% propilenglikol and acetate buffer pH 4.2 ad 100 % with High Shear homogenization method [ 1 ] with a speed of 24,000 rpm conducted for 8 minutes at 4 cycle. The NLC made by the same methode and formula , but 1.5 % cetyl alcohol was replaced with oleic acid . Furthermore, their lipid characteristic was analysed by their recrystallization index that was calculated based on the enthalpy value was determined by Differential Thermal Analysis ( DTA ) [ 2 ]

$$\% \text{ Crystallinity index} = \frac{\Delta H \text{ formula}}{\Delta H \text{ Lipid phase} \times \text{Lipid Phase Concentration}} \times 100\%$$

The crystal diffraction pattern determined by X Ray Defraktometer. Do also checks pH , morphology , particle size and particle size distribution with Dynamic Light Scattering ( DLS ) . Trapping efficiency is determined by the method of dialysis and determined by spectrophotometry . Stability antioxidants determined in accordance with the method IC50 1,1-Diphenyl - 2 - picrylhydrazyl ( DPPH ) on day 1 dan30 in controlled conditions . IC50 is the concentration of materials to be dampening the free radicals by 50 % . The larger the IC50 , the smaller the antioxidants activity. Percent reduction is calculated by the formula [ 3 ]

$$\% \text{ peredaman} = \left\{ \frac{A_{\text{control}} - A_{\text{sampel}}}{A_{\text{control}}} \right\} \times 100\%$$

Oclusivity test performed with in vitro methods . Calculated based on the percentage of water that evaporates in a vial covered with a membrane that is smeared preparations and which are not . Membrane used is a membrane filter (cellulose filter with pores 0.45 mm , Whatman no. 4) . Observations were made during a week . The oclusivity (F) is calculated by the following formula [ 4 ] :

$$F = \left( \frac{A-B}{A} \right) \times 100$$

Information :


A = The amount of water lost in the control

B = The amount of lost water in the sample

### Results and Discussion

Based on the data DTA thermogram single cetyl alcohol, base SLN and NLC base note that due to the addition of oleic acid peak calorimetry be more gentle. It shows become more amorphous. Based on calculations, percent recrystallization index, NLC is lower than the SLN, respectively: 0.32% dan 1,01%. It stretcher in accordance with the stated Hu et.al (2005) [5] that the decline in the degree of crystallinity of a material due to a lower order of the crystal lattice. The fall in the regularity of the crystal lattice has resulted in active ingredients more easily trapped and not easily terekspulsi out during storage. This makes the NLC able to trap larger lycopene ( $75.52 \pm 2.75\%$ ) than the SLN ( $57.30 \pm 5.10\%$ ). The presence of lipid liquid, in this case oleic acid, significantly decreased the size of the particles. SLN particle size of  $226.67 \pm 22.72$  nm while NLC  $134.03 \pm 19.01$  nm. Neither the particle size distribution of  $0.63 \pm 0.03\%$  of SLN and NLC  $0.48 \pm 0.06\%$ .

Stability is expressed based on changes in antioxidant power IC50. The smaller the IC50, the greater the antioxidant activity. From Figure 1 it appears that at the initial antioxidants activity determination, lycopene has antioxidant power greater than when lycopene is made either SLN or NLC. This caused by lycopene was trapped in the SLN and NLC lipid matrix. After storing for 30 days, the opposite happens that pure lycopene antioxidant activity declined sharply. While the antioxidant activity of lycopene in SLN lower than in NLC. It shows that the NLC system was better able to maintain the stability of lycopene antioxidant compared with SLN. It was related to lycopene trapping efficiency of the system. Lycopene efficiency entrapment of NLC was  $75.52 \pm 2.75\%$ , while in the SLN:  $57.30 \pm 5.10\%$ . That made the lycopene had better protected from environmental influences that can accelerate degradation.



**Figure 1** Diagram IC50 of lycopene , lycopene in the lycopene in the SLN and NLC on the 1st and 30<sup>th</sup>

The result of NLC occlusivity NLC was  $40.33 \pm 2.08\%$ , higher than the SLN ( $34.67 \pm 2.51\%$ ). This can be due to the smaller size of lycopene NLC particle there for no gaps between the particles . Occlusivity effect of nano-sized particles larger than micro-sized particles on the skin. It leads to reduced transcellular evaporation water lost . That's why the nano-sized cosmetic preparation can maintain skin moistur .

**Conclusion.** The conclusion of this study were: 1) SLN and NLC system were able to improve the stability of the lycopene antioxidant. 2) . The NLC delivery system ( with oleic acid as a liquid lipid ) better to improve the stability of the lycopene antioxidant than the SLN .

### Reference(s)

- [1] Mukhrejee, S., Ray, S., Thakur, R.S (2009) Solid Lipid Nanoparticles: A Modern Formulation Approach in Drug Delivery System. **Indian J Pharm Sci**, Vol 71, p. 349–358
- [2] Venkateswarlu, V., Manjunath, K. (2004) Preparation, characterization and in vitro release kinetics of clozapine solid lipid nanoparticles, **J. Control. Release**, 95, pp: 627-638
- [3] Rajesh, Patel., Natvar, Patel. (2011) In Vitro Antioxidant Activity Of Coumarin Compounds By DPPH, Super Oxide And Nitric Oxide Free Radical Scavenging Methods. **Journal of Advanced Pharmacy Education & Research**, Vol 1, p. 52-68



- [4] Loo, C., H., Basri M., Ismail R., Lau H.L.N., Tejo B.A., Kanthimathi M. S. & Choo Y.M. (2013) Effect of Composition in Nanostructured Lipid Carrier (NLC) on Skin Hydration and Occlusion, *International Journal of Nanomedicine*, 8, 13
- [5] Hu, Fu-Qiang., Jiang , Sai-Ping., Du, Yong-Zhong., Yuan, Hong., Ye, Yi-Qing., Zeng, Su., (2005). *Preparation and Characterization of Stearic Acid Nanostructured Lipid Carriers by Solvent Diffusion Method in an Aqueous System*. *Colloids and Surfaces B: Biointerfaces*, Vol. 45, p.167-173.