

KEPUSTAKAAN

PERPUSTAKAAN
UNIVERSITAS ANILANGKA
JAKARTA

1. Edgcombe H, Carter K, Yarrow S. Anesthesia in the prone position. Br. J. Anaesth. 2008; 100(2): 165-183.
2. Manson HM. Prone position of patients on operating room. Support equipment: intra-operative complication. Allen Medical System. 2009: 1-17.
3. Joo Jin, Kim YH, Lee J, Choi JH. Difference in the value of arterial and end-tidal carbondioxide tension according to different surgical positions: Does it reliably reflect ventilation-perfusion mismatch?. Korean J. Anesthesiol. 2012; 63(3): 216-220.
4. Hedenstierna G. Respiratory Physiology in Miller DR. Anesthesia 7th edition. 2005; 15: 361-389.
5. Raw DA, Beattie JK, Hunter JM. Anaesthesia for spinal surgery in adults. Br. J. Anaesth. 2003; 91(6): 886-904.
6. Peterson J, Ax M, Frey J, Sanches-Crespo A, Lindahl SGE, Mure M. Positive end-expiratory pressure redistributes regional blood flow and ventilation differently in supine and prone humans. Prone position; Beyond Physiology in Lippincott Williams & Wilkins. Anesthesiology. 2010; 113: 1262-1264.
7. Mure M, Domino KB, Lindahl SGE, Hlastala MP, Altemeier WA et al. Regional ventilation-perfusion distribution is more uniform in the prone position. J. Appl. Physiol. 2000; 88: 1076-1083.
8. Pelosi P, Croci M, Calappi E, Cerisara M, Mulazzi D et al. The Prone Position During General Anesthesia Minimally Affect Respiratory Mechanics While Improving Functional Residual Capacity and Increasing Oxygen Tension. Anest Analg. 1995; 80: 955-960.
9. Sud S, Sud M, Friedrich JO, Adhikari NKJ. Effect of mechanical ventilation in the prone position on clinical outcomes in patients with acute hypoxic respiratory failure: a systematic review and meta-analysis. CMAJ. 2008; 178(9): 1153-1161.

10. Gattinoni L, Protti A. Ventilation in the prone position: For some but not for all?. CMAJ. 2008; 178(9): 1174-1176.
11. Abdeyazdan Z, Nematollahi M, Ghazavi Z, Mohhamadizadeh M. The effect of supine and prone position on oxygenation in premature infants undergoing mechanical ventilation. IJNMR. 2010; 15(4): 229-233.
12. Mounier R, Adrie C, Francais A, Garrouste-Orgeas M, Cheval C et al. Study of prone position to reduce ventilator-associated pneumonia in hypoxaemic patients. Eur. Resph. J. 2010; 35: 795-804.
13. Gisolf J, Wilders R, Immink RV, van Lieshout JJ, Karemker JM. Tidal volume, cardiac output and functional residual capacity determine end-tidal CO₂ transient during standing up in humans. J. Physiol. 2003; 554(2): 579-590.
14. Sawhney A, Kumar N, Sreenivas V, Gupta S, Tyagi V, Puliye JM. Prone versus supine position in mechanically ventilated children: a pilot study. Med. Sci. Monit, 2005; 11(5): CR235-240.
15. Guerin C. Ventilation in the prone position in patient with acute lung injury/acute respiratory distress syndrome. In Lippincott Williams & Wilkins Curr. Opin. Crit. Care. 2006; 12: 50-54.
16. Pelosi P, Tubiolo D, Mascheroni D, Vicardi P, Stefani C et al. Effect of the Prone Position on Respiratory Mechanics and Gas Exchange during Acute Lung Injury. Am J Respir Crit Care Med. 1998; 157: 387-393.
17. Romero CM, Cornejo RA, Galvez LR, Llanos OP, Tobar EA et al. Extended prone position ventilation in severe acute respiratory distress syndrome: A pilot feasibility study. JCrC. 2008; 02: 1-8.
18. St. John RE. End-Tidal Carbon Dioxide Monitoring. Crit Care Nurse. 2003; 23: 83-88.
19. Morgan EG, Mikhail MS, Murray MJ. Capnography. Patient Monitor. Clinical Anesthesiology 4th edition. 2006; 6: 141-144.