



ICPPS 2014

Proceeding

The 1st International Conference on Pharmaceutics & Pharmaceutical Sciences

Drug Delivery Systems:

From Drug-Discovery, Pre-formulation, Formulation and Technological Approaches for
Poorly Soluble Drugs and Protein



The 1st International Conference on Pharmaceutics & Pharmaceutical Sciences Proceedings

ISBN : 978-602-72333-0-0

(Letter of National ISBN Agency No. 4127/E.8/p/03.2015 Date 18 March 2015)

1st edition Proceeding

Published by:

Faculty of Pharmacy Universitas Airlangga
Surabaya, Indonesia

Address:

Kampus B Jl. Dharmawangsa Dalam
Surabaya 60286
Phone +62 31 5033710
Fax +62 31 5020514
Website: www.icpps2014.com or www.ff.unair.ac.id
Email: icppinfo@gmail.com

ISBN 978-602-72333-0-0



TABLE of CONTENT

Preface from Chairman	ii
Committee	ii
Table of Contents	iii
Author Index	iii

AUTHOR INDEX

COMPARISON OF SODIUM STARCH GLYCOLATE AND CROSSCARMELLOSE SODIUM AS SUPERDISINTEGRANT IN MEFENAMIC ACID FAST DISINTEGRATING TABLET Adeltrudis Adelsa D, Oktavia Eka Puspita, Amalia Ayuningtyas, Marulita Isadora	1
STUDY EXPRESSION OF HUMAN ERYTHROPOIETIN EXPRESSION IN MAMMALIAN CELL Adi Santoso, Popi Hadiwisnuwardhani, Yana Rubiana, Yulaika Romadhani, Endah Puji Septisetyani, Dyaningtyas D.P. Putri	4
ANTIOXIDANT STABILITY ASSAY OF ALPHA TOCOPHERYL ACETATE IN SOLID LIPID NANOPARTICLE SYSTEM (LIPID BASE BEESWAX AND MONOSTEARIC GLISERYL 50:50) Anggie Widhi, Noorma Rosita, Widji Soeratri	8
A BIOACTIVE BOVINE HYDROXYAPATITE–GELATIN IMPLANT FOR IN VITRO GENTAMICIN RELEASE Aniek Setiya Budiatin, M. Zainuddin, Junaidi Khotib, Diah Himawati	13
EFFECT OF COMPARISON SURFACTANT AND COSURFACTANT IN WATER/OIL MICROEMULSION IN RELEASE OF OVALBUMIN Microemulsion Water/Oil with Surfactant (Span 80-Tween 80) : Cosurfactant (Ethanol) =5:1, 6:1, and 7:1 Anisa Rizki Amalia, Riesta Primaharinastiti, Esti Hendradi	18
ANALYSIS OF MYCOLIC ACIDS CLEAVAGE PRODUCT OF <i>Mycobacterium tuberculosis</i> BY GAS CHROMATOGRAPHY-FLAME IONIZATION DETECTOR Asri Darmawati, Deby Kusumaningrum, Isnaeni, Muhamad Zainuddin	21
PERIPLASMIC EXPRESSION OF GENE ENCODING ANTI-EGFRVIII SINGLE-CHAIN VARIABLE FRAGMENT ANTIBODY USING PeB LEADER SEQUENCE IN <i>ESCHERICHIA COLI</i> Kartika Sari Dewi, Debbie Sofie Retnoningrum, Catur Riani, Asrul Muhamad Fuad	24
IN VIVO ANTIMALARIAL ACTIVITY OF ETHANOL EXTRACT AND ETHYL ACETATE FRACTION OF <i>Alectryon serratus</i> LEAVES ON <i>Plasmodium berghei</i> INFECTED MICE Aty Widyawaruyanti, Uswatun Khasanah, Lidya Tumewu, Hilkatul Ilmi, Achmad Fuad Hafid, Indah S Tantular	30
PROFILE OF COMMUNITY PHARMACISTS KNOWLEDGE IN PATIENT ASSESSMENT WITH INFLUENZA SYMPTOMS AND ITS PRODUCTS Azza Faturrohmah, Arie Sulistyarini, Ana Yuda	33

SOLUBILITY AND DISSOLUTION STUDY OF KETOPROFEN – HIDROXYPROPYL- β -CYCLODEXTRIN INCLUSION COMPLEX (Prepared by Kneading Method) Bambang Widjaja, Achmad Radjaram, Arafah Zulhana	37
FORMULATION AND STABILITY TESTING OF MELOXICAM SOLID DISPERSION GEL Budipratiwi Wisudyaningsih, Inka Dewi Nur Anggaraini, Fersiya Wardani	40
EFFECT OF MENTHOL AS PENETRATION ENHANCER TO DICLOFENAC SODIUM MEMBRANE-TYPED TRANSDERMAL PATCH CHARACTERIZATION Destria Indah Sari, Esti Hendradi, Junaidi Khotib	43
PHYSICAL CHARACTERISTICS AND RELEASE STUDY OF OVALBUMIN FROM ALGINATE MICROSPHERES PREPARED BY DIFFERENT CONCENTRATION OF ALGINATE AND BaCl ₂ USING AEROSOLIZATION TECHNIQUE Dewi Melani Hariyadi, Tristiana Erawati, Sisilia Ermawahyuningtyas	46
MUCOADHESIVE TABLET OF ETHANOLIC EXTRACT OF SAMBILOTO (<i>Andrographis paniculata</i>) AS ANTIDIABETIC USING CHITOSAN Dhadhang Wahyu Kurniawan, Hening Pratiwi, and Lingga Ikaditya	50
PHYSICAL INTERACTION STUDY OF IBUPROFEN-STEARIC ACID BINARY MIXTURE Diajeng Putri Paramita, Dwi Setyawan, Dewi Isadiartuti	59
MOLECULAR MODELING AND SYNTHESIS OF 1-(3,4-Dichlorobenzoyl)-1,3-dimethylurea Dian Agung Pangaribowo, Siswandono, Bambang Tri Purwanto	63
EXPRESSION OF RECOMBINANT HUMAN GRANULOCYTE-COLONY STIMULATING FACTOR WITHIN PERIPLASMIC COMPARTMENT OF <i>Escherichia coli</i> USING PeIB LEADER PEPTIDE Dian Fitria Agustiyanti, Asrul Muhamad Fuad	66
EVALUATION OF ANTIHYPERURICEMIC ACTIVITY FROM BULBS OF BAWANG TIWAI (<i>Eleutherine palmifolia</i> (L.) Merr.) BY IN VITRO AND IN VIVO STUDIES Dian Ratih Laksmiwati, Rininta Firdaus, Yulinda, Mediana Astika	72
ANTIOXIDANT ACTIVITY OF 96% ETHANOL EXTRACT OF COMBINATION OF STRAWBERRY FRUIT (<i>Fragaria x ananassa</i> Duch.) AND STARFRUIT (<i>Averrhoa carambola</i> L.) USING ABTS FREE RADICAL SCAVENGING METHOD Diana Serlahwaty, Indira Natalia Timang	76
ENHANCEMENT OF SOLUBILITY AND DISSOLUTION ATORVASTATIN BY MICROCRYSTALLIZATION METHOD Dolih Gozali, Yoga Windu Wardhana, Ronny Tandela	79
IN VITRO ANTIMALARIAL ACTIVITY OF DICHLOROMETHANE SUB-FRACTION OF <i>Eucalyptus globulus</i> L. Stem AGAINST <i>Plasmodium falciparum</i> Elis Suwarni, Achmad Fuad Hafid, Aty Widyawaruyanti	86
<i>Arcangelisia flava</i> INCREASES RATS' LEUKOCYTES BUT HAS BIPHASIC EFFECT ON RATS' LYMPHOCYTE Endah Puspitasari, Evi Umayah Ulfa, Vita Ariati, Mohammad SulthonHabibi	89

IN VITRO ANTIMALARIAL ACTIVITY OF CHLOROFORM SUBFRACTION OF SALAM BADAK LEAVES (<i>Acmena acuminatissima</i>)	
Erna Cahyaningsih, Achmad Fuad Hafid, Aty Widyawaruyanti	92
CHARACTERIZATION OF DOSAGE FORM AND PENETRATION DICLOFENAC SODIUM WITH MICROEMULSION SYSTEM IN HPMC 4000 GEL BASE (Microemulsion W/O with ratio use of surfactant Span 80 – Tween 80 : Cosurfactant Ethanol 96% = 6:1)	
Esti Hendradi, Tutiek Purwanti, Karina Wahyu Irawati	95
CONSTRUCTION AND VALIDATION OF THE STRUCTURE-BASED VIRTUAL SCREENING PROTOCOLS WITH PDB CODE OF 3LN1 TO DISCOVER CYCLOOXYGENASE-2 INHIBITORS	
Mumpuni E, Nurrochmad A, Pranowo HD, Jenie UA, Istiyastono EP	99
VALIDATED UV SPECTROPHOTOMETRIC METHOD FOR THE DETERMINATION OF ASPIRIN IN RABBIT PLASMA : APPLICATION TO BIOAVAILABILITY STUDY OF ASPIRIN MICROCAPSULE IN RABBIT	
Faizatun, Novi yantih, Teguh Iman Saputra	102
EFFECT OF COMPARISON OF SURFACTANT AND COSURFACTANT W/O MICROEMULSION OVALBUMIN WITH SOYBEAN OIL TO PHYSICOCHEMICAL CHARACTERIZATION (w/o Microemulsion with Surfactant Span 80- Tween 80 : Cosurfactant Ethanol 96% = 5:1; 6:1 and 7:1)	
Farida Mutiara Sari, Riesta Primaharinastiti, Esti Hendradi	105
pH INFLUENCE IN DESALTING PROCESS OF CRUDE PERTUSSIS TOXIN (PT) AND FILAMENTOUS HEMAGGLUTININ (FHA) PURIFICATION FROM <i>Bordetella pertussis</i> BY SEPHADEX G-25 COLUMN CHROMATOGRAPHY	
Faris Adrianto, Esti Hendradi, Neni Nurainy, Isnaeni	108
SEPARATION OF COSMETIC PRESERVATIVES USING SILICA-BASED MONOLITHIC COLUMN	
Febri Annuryanti, Riesta Primaharinastiti, Moch. Yuwono	111
PREPARATION AND CHARACTERIZATION OF TELMISARTAN-CITRIC ACID CO-CRYSTAL	
Fikri Alatas, Hestairy Ratih, Sundani Nurono Soewandhie	114
PATIENTS' AND CAREGIVERS' LIQUID MEDICATION ADMINISTRATION ERRORS	
Gusti Noorrizka Veronika Achmad, Gesnita Nugraheni	117
THE POTENCY OF CANARIUM OIL (<i>Canarium indicum</i>) AS A MATERIAL FOR STRUCTURED LIPID PRODUCTION	
Hamidah Rahman, Johnner P Sitompul, Kusnandar Anggadiredja, Tutus Gusdinar	121
EFFECT OF TREHALOSE ON THERMAL PROPERTIES OF PHOSPHOLIPID-DDA AND TPGS MIXTURES	
Helmy Yusuf	124
PREPARATION AND CHARACTERIZATION OF FLUKONAZOLE- β -CYCLODEXTRIN INCLUSION COMPLEXES	
Hestairy Ratih, Fikri Alatas, Erin Karlina	127
ISOLATION AND IDENTIFICATION OF ANTIOXIDANT COMPOUND BY BIOPRODUCTION OF ENDOPHYTIC FUNGI OF TURMERIC (<i>Curcuma longa</i> L.) ISOLATE CL.SMI.RF11	
Hindra Rahmawati, Bustanussalam, Partomuan Simanjuntak	130

MODIFICATION PROCESS OF NATURAL CASSAVA STARCH : THE STUDY OF CHARACTERISTICS AND PHYSICAL PROPERTIES Prasetya, Jemmy A, C.I.S. Arisanti, N.P.P.A. Dewi, G.A.R. Astuti, N.W.N Yulianingsih, I M.A.G. Wirasuta	133
DRUG USE PROFILE OF DIABETIC PATIENTS IN EAST SURABAYA PRIMARY HEALTH CARE I Nyoman Wijaya, Azza Faturrohmah, Ana Yuda, Mufarriha, Tesa Geovani Santoso, Dina Kartika, Hikmah Prasasti N, Whanni Wido Agustin	136
GLYCINE MAX DETAM II VARIETY AS PREVENTIVE AND CURATIVE ORGAN DAMAGE DUE TO EXPOSURE TO ,LEAD (Pb) Rika Yulia, Sylvan Septian Ressandy, Gusti Ayu Putu Puspikaryani, I Putu Agus Yulyastrawan, Dewa Ayu Kusuma Dewi	142
AN ACTIVITY TEST OF MATOA LEAVES EXTRACT AS HEART RATE FREQUENCY REDUCTION WITH ADRENALINE INDUCTION Ika Purwidyaningrum, Elin Yulinah Sukandar, Irdha Fidrianny	144
EFFORT TO REDUCE COMPRESSIBILITY OF RAMIPRIL THROUGH CRYSTAL ENGINEERING Indra, Sundani N Soewandhi	147
IN VITRO ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY OF ETHANOLIC LEAF EXTRACT AND FRACTIONS OF <i>Rauvolfia serpentina</i> (L.) Benth. ex Kurz Julie Anne D. Bolaños, Ivan L. Lawag	150
PERIPLASMIC EXPRESSION OF GENE ENCODING ANTI-EGFRvIII SINGLE-CHAIN VARIABLE FRAGMENT ANTIBODY USING PelB LEADER SEQUENCE IN <i>Escherichia coli</i> Kartika Sari Dewi, Debbie Sofie Retnoningrum, Catur Riani, Asrul Muhamad Fuad	153
CHARACTERIZATION AND LD ₅₀ VALUE DETERMINATION OF 1,5-bis(3'-ethoxy-4'-hydroxyphenyl)-1,4-pentadiene-3-one (EHP) Lestari Rahayu, Septian, Esti Mumpuni	159
DEVELOPMENT OF MELOXICAM TRANSDERMAL MATRIX TYPE PATCH USING POLYVINYL PYRROLIDONE, HYDROXYPROPYL METHYLCELLULOSE, AND ETHYL CELLULOSE COMBINATION Lidya Ameliana, Monica Iwud, Selly Rio	162
ANTIHEPATITIS C VIRUS ACTIVITY SCREENING ON <i>Harpullia arborea</i> EXTRACTS AND ISOLATED COMPOUND Lidya Tumewu, Evhy Apryani, Mei Ria Santi, Tutik Sri Wahyuni, Adita Ayu Permanasari, Myrna Adianti, Chie Aoki, Aty Widyawaruyanti, Achmad Fuad Hafid, Maria Inge Lusida, Soetjipto, Hak Hotta.....	165
HPLC METHOD PRECISION TO ASSAY OF A-MANGOSTIN IN Mangosteen (<i>Garcinia mangostana</i> L.) FRUIT RIND EXTRACT FORMULATED IN ORAL SOLUTION Liliek Nurhidayati, Siti Sofiah, Ros Sumarny, Kevin Caesar	168

PREPARATION AND CHARACTERIZATION OF NARINGENIN-LOADED CHITOSAN NANOPARTICLES FOR CHEMOPREVENTION	
Lina Winarti, Lusia Oktora Ruma Kumala Sari	170
RELATIONSHIP OF KNOWLEDGE AND PATIENT BEHAVIOR ON SELF MEDICATION PIROXICAM (Studies of Pharmacy in Sukun District , Malang City)	
Liza Pristanty, Reshtia Eriana Putri, Hidayah Rachmawati	173
EFFECT OF CHRONIC USE OF ENERGY DRINK ON KIDNEY	
Mahardian Rahmadi, Zamrotul Izzah, Mareta Rindang A, Aniek Setya B, Suharjono	176
SCREENING OF SURFACE MODIFIERS TO PRODUCE STABLE NANOSUSPENSION : A GENERAL GUIDANCE	
Maria Lucia Ardhani Dwi Lestari	179
DEVELOPMENT OF SIMPLE POLYPHENOL SENSOR BASED ON SODIUM META PERIODATE AND 3-METHYL-2-BENZOTHIAZOLINONE HYDRAZONE FOR COFFEE SAMPLES	
Moch. Amrun Hidayat, Nindya Puspitaningtyas, Agus Abdul Gani, Bambang Kuswandi	181
VALIDATION OF AN HPLC ANALYTICAL METHOD FOR DETERMINATION OF LEVOFLOXACIN IN OPHTHALMIC PREPARATIONS	
Mochamad Yuwono, Rieta Primaharinastiti, Ageng Teguh Wardoyo	184
VALIDATION OF SPECTROPHOTOMETRIC METHOD FOR ESTIMATION OF EPERISONE HYDROCHLORIDE IN TABLET DOSAGE FORM	
Nia Kristiningrum, Diah Yuli Pangesti	187
ANTIOSTEOPOROTIC ACTIVITY OF 96% ETHANOLIC EXTRACTS OF ABELMOSCHUS MANIHOT L.MEDIK LEAVES AND EXERCISE ON INCREASING BONE DENSITY OF FEMALE MICE'S FEMORAL TRABECULAR	
Niliestria Ayu Faramitha Sholikhah	190
EFFECT OF -CYCLODEXTRIN ON SPF VALUE AND INHIBITION OF KOJIC ACIDSTYROSINASE ACTIVITY IN VANISHING CREAM BASE FORMULATION (ON SUNSCREEN PRODUCT CONTAINED OXYBENZONE)	
Noorma Rosita, Diana, Diana Winarita, Tristiana Erawati, Widji Soeratri	193
ANTIMICROBIAL ACTIVITY OF LACTOBACILLI PROBIOTIC MILK AND GUAVA LEAF ETHANOLIC EXTRACT (Psidium guajava) COMBINATION AGAINST BACTERIAL CAUSE OF DIARRHEA	
Nur Putri Ranti, Isnaeni, Juniar Moechtar, Febri Annuryanti	197
THE INFLUENCES OF PARTICLE SIZE AND SHAPE ON ZETA POTENTIAL OF COENZYME Q10 NANOSUSPENSION	
Nuttakorn Baisaeng	200
SYNTHESIS, MOLECULAR DOCKING, AND ANTITUMOR ACTIVITY OF N,N'-Dibenzoyl-N,N'-Dimethylurea AGAINST HUMAN BREAST CANCER CELL LINE (MCF-7)	
Nuzul Wahyuning Diyah	203
EXPRESSION OF ANTI-EGFRVIII SINGLE CHAIN FRAGMENT ANTIBODY (SCFV) ON THE SURFACE OF PICHIA PASTORIS	
Pratika Viogenta, Asrul Muhamad Fuad, Suharsono	206

SUBCLONING OF <i>csf3syn</i> (COLONY STIMULATING FACTOR-3) GENE INTO pGAPZ? AND TRANSFORMATION OF RECOMBINANT VECTOR INTO PICHIA PASTORIS Prety Ihda Arfia, Asrul Muhamad Fuad	212
THE USE OF PERICARP MANGOSTEEN (<i>Garcinia mangostana L.</i>) EXTRACT IN FORMULATION OF CREAM-TYPE O / W Rahmah Elfiyani, Naniek Setiadi Radjab, Mia Sagita Sofyan	218
CHITOSAN BASED PARTICULATE CARRIER OF DITERPENE LACTON OF SAMBILOTO PREPARED BY IONIC GELATION-SPRAY DRYING :EFFECT OF STIRRING RATE AND NOZZLE DIAMETER Retno Sari, Titin Suhartanti, Dwi Setyawan, Esti Hendradi, Widji Soeratri	222
GAS CHROMATOGRAPHY-MASS SPECTROMETRY METHOD VALIDATION FOR PESTICIDES RESIDUES ANALYSIS IN FOOD USING QuEChERS KIT Riesta Primaharinastiti, Setyo Prihatiningtyas, Mohammad Yuwono	225
CHARACTERIZATION OF PARACETAMOL ORALLY DISINTEGRATING TABLET USING GELATIN 1% AND 2% AS BINDER AND POLYPLASDONE XL-10 10% AS DISINTEGRANT (Prepared by Freeze Drying Method) Roisah Nawatila, Dwi Setyawan. Bambang Widjaja	230
ANTIOXIDANT STUDY OF COSOLVENT SOLUTION OF MANGOSTEEN (<i>Garcinia mangostana L.</i>) RIND EXTRACT IN RATS BY USING MDA PARAMETER Ros Sumarny, Liliek Nurhidayati, Yati Sumiyati, Fransiska Diana Santi	233
SCREENING OF SELECTED PHILIPPINE ROOT CROPS FOR ?-Glucosidase INHIBITION Sarah Jane S. Almazora, Ivan L. Lawag	236
NIOSOME EMULGEL FORMULATION AND STABILITY TEST OF CINNAMON (<i>Cinnamomum burmanii Nees & Th. Nees</i>) Bark ETHANOLIC EXTRACT AS ANTIOXIDANT Sasanti Darijanto, Fidriani Irla, Widhita P.A.S	239
THEOPHYLLINE RELEASE FROM SUSTAINED RELEASE TABLET USING LACTOSA AND PVP K30 AS A CHANELLING AGENT Sugiyartono, Retno Sari, Agus Syamsur Rijal, TriMulyani, Agustina Maharani	242
EFFORTS TO PRODUCE 1-(BENZOYLOXY)UREA AS ANTICANCER DRUG CANDIDATE Suko Hardjono	245
DEVELOPMENT OF PEGylated RAPAMYCIN LITHOCHOLIC ACID MICELLE FOR CANCER THERAPY Aran Tapsiri, Kanokwan Jaiprasert, Rungtip Nooma, Awadsada Sukgasem, Supang Khondee	248
FEASIBILITY OF ORAL IMMUNIZATION AGAINST JAPANESE ENCEPHALITIS VIRUS USING CHITOSAN PARTICLES Supavadee Boontha, Worawan Boonyo, Hans E. Junginger, Tasana Pitaksuteepong, Neti Waranuch, Assadang Polnok, Narong Nitatpattana, Suthee Yoksan	251

THE INFLUENCE OF HYDROXY GROUP AT ORTHO (o) AND PARA (p) POSITIONS ON METILBENZOAT AGAINST SYNTHESIS OF HIDROKSIBENZOHIDRAZIDA DERIVATIVES	
Suzana, Adita R, Melanny Ika .S, Juni Ekowati, Marcellino Rudyanto, Hadi Poerwono, Tutuk Budiaty	254
ANTIOXIDANT ACTIVITY TEST OF BEE PROPOLIS EXTRACT (Apis mellifera L.) USING DPPH (1,1-diphenyl-2-picrilhidrazyl) FREE RADICALS SCAVENGING ACTIVITY	
Titiek Martati, Shahyawidya	258
CAPSULE FORMULATION and EVALUATION COMBINATION OF AQUEOUS EXTRACT OF Phaleria's (Phaleria macrocarpa (Scheff Boerf)) FRUITS and LEAVES as ANTIHYPERTENSIVE AGENT	
Titta H Sutarna, Sri Wahyuningsih, Julia Ratnawati, Fahrouk P, Suci Nar Vikasari, Ita Nur Anisa	260
MANUFACTURE AND CHARACTERIZATION OF SOLID DISPERSION GLIKLAZID- PVP K90	
Titta H Sutarna, Fikri Alatas, Cicih Ayu Ningsih	264
ANTI-INFLAMMATORY ACTIVITY OF PARA METHOXY CINNAMIC ACID (PMCA) IN NANOEMULSION USING SOYBEAN OIL	
Tristiana Erawati M, Anneke Indraswari P, Nanda Ghernasih N.C, Noorma Rosita, Suwaldi Martodihardjo, Widji Soeratri	266
PHYSICAL CHARACTERISTICS AND PENETRATION OF DICLOFENAC SODIUM NIOSOMAL SYSTEM USING SPAN 20 AND SPAN 60	
Tutiek Purwanti, Esti Hendradi, Noverika A. Putri, Nurtya J. Devi	271
FORMULATION AND CHARACTERISATION OF JUICE OF LIME GEL USING CMC-Na BASE	
Uswatun Chasanah, Esti Hendradi, Inayah	275
HEPATOPROTECTIVE ACTIVITY OF Bidens pilosa L. IN CARBON TETRACHLORIDE INDUCED HEPATOTOXICITY IN RATS	
Vina Alvionita Soesilo, C.J. Soegihardjo	278
CYTOTOXIC ACTIVITY ASSAY AGAINTS HELA CELL LINES OF NOVEL ANTICANCER DRUG : N-(PHENYLCARBAMOYL)ISOBUTIRAMIDE	
Wimzy Rizqy Prabhata, Tri Widiandani, Siswandono	284
SIMPLE STEPS PURIFICATION OF RECOMBINANT HUMAN ERYTHROPOIETIN PRODUCED IN CHINESE HAMSTER OVARY CELL CULTURE	
Yana Rubiyana, Endah Puji Septisetyani, Adi Santoso	287
KINETICS STUDY COCRYSTALS KETOCONAZOLE-SUCCINIC ACID PREPARED WITH SLURRY METHOD BASED ON POWDER X-RAY DIFFRACTION (PXRD)	
Yuli Ainun Najih, Dwi Setyawan, Achmad Radjaram	290
CONSTRUCTION OF RECOMBINANT IMMUNOTOXIN Anti-EGFRvIII scFv::HPR FUSION PROTEIN AND INDUCIBLE EXPRESSION IN Pichia pastoris AS A TARGETED DRUG CANDIDATE	
Yuliawati, Asrul Muhamad Fuad	297
ALTERED PHARMACOKINETIC OF LEVOFLOXACIN BY COADMINISTRATION OF ATTAPULGITE	
Zamrotul Izzah, Toetik Aryani, Amalia Illiyyin, Budi Suprapti	303

ANTI-INFLAMMATORY ACTIVITY OF PARA METHOXY CINNAMIC ACID (PMCA) IN NANOEMULSION USING SOYBEAN OIL

Tristiana Erawati M., Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya, era_ffua@yahoo.co.id, **Anneke Indraswari P.**, Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya, **Nanda Ghernasih N.C.**, Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya, **Noorma Rosita**, Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya, **Suwaldi Martodihardjo**, Pharmaceutics Department of Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, **Widji Soeratri**, Pharmaceutics Department of Faculty of Pharmacy, Universitas Airlangga, Surabaya.

INTRODUCTION

Inflammation is a protective mechanism of the local microcirculation to tissue injuries caused by physical trauma, stimulation by hazardous chemicals, heat, antigen-antibody reaction and the effect of microbial.[2] It is known to be involved in the inflammatory reactions such as release of histamine, bradykinin, prostaglandins, fluid cell migration, extravasations, tissue break down and repair which are aimed at host defense and usually activated in most disease conditions. Para methoxy cinnamic acid (PMCA) known has topical anti-inflammatory effect but only 0.64 compare with Na-diclophenac.[4] Its cause PMCA is a poorly soluble drug substance (BCS II), the solubility in acetate buffer pH 4.2 ± 0.2 was 70.04 ± 0.66 mg/liter.[3] So in this study to increase the solubility PMCA loaded in nanoemulsion using soybean oil. And then the anti-inflammatory activity of PMCA in nanoemulsion was measured by the release rate, penetration rate through rat skin using Franz diffusion cell and histological test on mice's ear skin.

MATERIAL AND METHODS

Research Material:

Para methoxy cinnamic Acid (Sigma Aldrich), soybean oil (PT Kurniajaya), Tween 80 (Sigma Aldrich), Span 80 (Sigma Aldrich), ethanol 96 % (Merck), acetic acid (Merck), sodium acetate (Merck), NaCl (Merck), NaH₂PO₄ (Merck), Na₂HPO₄ (Merck), croton oil (Sigma) and aquademineralisata (PT Brataco)

Animals

Male Wistar rats (150 - 230 gm) and mince (20 - 30 gm) were taken from PUSVETMA Suraba-

ya. The animals were housed under standard conditions of temperature (25 ± 2)°C, 12/12 hours light/dark cycles and fed with standard pellets. All animal experiments were conducted with the permission from Animal Care and Use Committee (ACUC) of Veterinary Faculty, Airlangga University, Indonesia. (Reference number; 378-KE).

Research Method

Nanoemulsion formula containing PMCA PMCA in nanoemulsion produced base on the formula presented in Table 1.

Materials	Concentration (%)
PMCA	0.2
Soybean oil	2.66
Span 80	1.92
Tween 80	18.66
Ethanol 96%	3.42
Acetate buffer solution pH 4.2 ± 0.2	ad 100

Table 1. PMCA Nanoemulsion Formula

Nanoemulsion formula used in this study is the result of a previous study by Erawati et.al, using a combination of surfactant Tween 80 and Span 80 with a ratio of 9:1 (having HLB 14), the ratio of surfactant and co surfactant 6:1, the ratio of oil phase (soybean oil) and water phase (acetate buffer solution pH 4.2 ± 0.2) is 27.5:1. [3]

Solubility test

PMCA solubility test; on the nanoemulsion

Proceeding
The 1st International Conference
on Pharmaceutics & Pharmaceutical Sciences



ICPPS 2014

using soybean oil (as oil phase) added with 500 mg PMCA, then it was sonicated for 10 minutes and shaking with 150 rpm for 60 minutes. And then, the amount of PMCA dissolved in nanoemulsion was determined using spectrophotometer. After that the nanoemulsion PMCA characterized include; density, viscosity, pH, droplet morphology, droplet size and polydispersity index.

Release test

Membrane Preparation; the membrane used in the test release of PMCA in nanoemulsion system is a cellophane membrane. Membrane cut to size, then immersed in aqua-demineralization for \pm 12 hours. A moment before use, the membrane is drained until no water is dripping, and then mounted on the surface of the receptor compartment of Franz diffusion cell.

Measurement the amount of the PMCA release from the nanoemulsion; receptor compartment of Franz diffusion cell filled with phosphate buffer pH 6.0 ± 0.2 up to full. Then, 2 ml of nanoemulsion PMCA inserted into the donor compartment. Experimental temperature is set and maintained at a temperature of $32 \pm 2^\circ\text{C}$. Magnetic stirrer rotated at a speed of 100 rpm. Samples of 1 ml were taken within a certain time interval, i.e. at 0, 5, 10, 15, 30, 45 minutes, and then 1, 1.5, 2, 3, 4, 6, 8, 10, 12, 24 hours. Immediately after sampling medium was replaced with phosphate buffer pH 6.0 ± 0.2 with a volume of samples taken. Subsequently, samples were taken observed with spectrophotometer. PMCA concentration in the sample was calculated using the regression equation of standard curve. Determination of PMCA cumulative amount released per unit membrane area (mg/cm²) was calculated from the concentration obtained each time ($\mu\text{g/ml}$) which had been corrected with the Wurster's equation. Furthermore, multiplied by the number of medium and divided by the membrane surface area. The results obtained by the cumulative number of PMCA released per unit time. The release profile of PMCA, is

done by making a relations curve between the cumulative number of PMCA released (mg/cm²) versus time (minutes). The release rate of (Flux release) PMCA was obtained from the slope of the regression equation in the steady state.

Penetration test

Membrane Preparation; skin male Wistar rats that had been shaved used as the membrane in the penetration test of PMCA in nanoemulsion system.

Measurement the amount of PMCA penetrate through rat skin; receptor compartment of Franz diffusion cell filled with phosphate buffer pH 7.0 ± 0.2 up to full. Then, 2 ml of nanoemulsion PMCA inserted into the donor compartment. Samples of 1 ml were taken within a certain time interval as in release test.

RESULT AND DISCUSSION

The PMCA solubility in nanoemulsion 3.07 ± 0.19 g/liter (presented in Table 2) increased than its solubility in acetate buffer pH 4.2 ± 0.2 is 70.04 ± 0.66 mg/liter. The characteristics of nanoemulsion containing PMCA presented in Table 3; the density value was 1.0263 ± 0.0002 is almost equal to the density of water; the viscosity was 5.58 ± 0.05 cP; the pH value was 4.47 ± 0.006 it was in the range of human skin pH, so it will not cause irritation; the droplet size was 57.3 ± 7.6 nm and the polydispersity index was 0.548 ± 0.044 it was indicate a moderate particle size distribution.[1] The droplet morphology of empty nanoemulsion and nanoemulsion containing PMCA in saturated solubility by TEM.type JEM-1400 was presented in Figure 1.

Medium	The amount of PMCA dissolved
Nanoemulsion	3.07 ± 0.19 g/liter
Acetate buffer pH 4.2 ± 0.2	70.04 ± 0.66 mg/liter

Table 2. The solubility of PMCA in nanoemulsion and in acetate buffer pH 4.2 ± 0.2

**The characteristics of nanoemulsion containing PMCA**

density	1.0263 ± 0.0002
viscosity	5.58 ± 0.05 cP
pH	4.47 ± 0.006
droplet size	57.3 ± 7.6 nm
polydispersity index	0.548 ± 0.044

Table 3. The characteristics of nanoemulsion containing PMCA

The release profile and the release rate of PMCA in acetate buffer pH 4.2 ± 0.2 and in nanoemulsion were presented in figure 2 and figure 3. After investigate for 24 hours (1440 minutes) the result shows that the release rate of PMCA in nanoemulsion ($0.4024 \pm 0.0339 \mu\text{g}/\text{cm}^2/\text{minute}$) increased compare with it in acetate buffer pH 4.2 ± 0.2 is $0.0239 \pm 0.0003 \mu\text{g}/\text{cm}^2/\text{minute}$.




Figure 2. Release profile of PMCA in acetate buffer pH 4.2 ± 0.2 (control) and in nanoemulsion (F I*)




Figure 2. Penetration profile of PMCA in acetate buffer pH 4.2 ± 0.2 (control) and in nanoemulsion (F I*)




Figure 3. Histogram of release rate of PMCA in acetate buffer pH 4.2 ± 0.2 (control) and in nanoemulsion (F I*)




Figure 5. Histogram of penetration rate of PMCA in acetate buffer pH 4.2 ± 0.2 (control) and in nanoemulsion (F I*)

Proceeding
The 1st International Conference
on Pharmaceutics & Pharmaceutical Sciences



ICPPS 2014

Also the penetration rate of PMCA in nanoemulsion was increased too ($0.1513 \pm 0.0314 \mu\text{g}/\text{cm}^2/\text{minute}$) compare with the penetration rate of PMCA in acetate buffer pH 4.2 ± 0.2 ($0.0341 \pm 0.0003 \mu\text{g}/\text{cm}^2/\text{minute}$) as presented in figure 4 and 5. The increased of penetration rate of PMCA in nanoemulsion caused by 1) the solubility of PMCA in nanoemulsion increased (more than 43x compare with the solubility of PMCA in acetate buffer pH 4.2 ± 0.2) and 2) the surfactant and co surfactant in nanoemulsion formula can function as enhancers.

From figure 6 and table 4, known that the anti-inflammatory effect of PMCA in nanoemulsion (F I*) higher than the anti-inflammatory effect of PMCA in acetate buffer pH 4.2 ± 0.2 (control). The result of statistical test using ANOVA-one way ($P = 95\%$), known there are no significant different between the ears mice skin thickness after treated with PMCA in nanoemulsion (F I*) with health skin (K-) but thinner than inflammation skin (K+) also with the ears mice skin after treated with PMCA in acetate buffer pH 4.2 ± 0.2 (control).




Figure 6. The histology profile of ears mice skin, health skin (K-), inflammation skin (K+), after treated with PMCA in acetic buffer pH 4.2 ± 0.2 (control), and in nanoemulsion (F I*)

Sample	The Average Of Skin Thickness (μm)
K-	282.96 ± 36.80
K+	592.53 ± 59.81
Kontrol	541.69 ± 56.56
F I*	310.56 ± 30.76

Table 4. The average of ears mice skin after treated with PMCA in acetate buffer pH 4.2 ± 0.2 (control), and in nanoemulsion (F I*) compare with health skin (K-) and inflammation skin (K+).

CONCLUSION

The nanoemulsion with soybean oil as drug delivery system can increase the anti-inflammatory activity of PMCA.

ACKNOWLEDGMENT

Thank to Indonesia Government (DIKTI) who give Research Fund through Rector of Universitas Airlangga (through LPPM-UA).

REFERENCES

1. Duan C.Y., Xia Z.Y., Gui B.S., Xue J. F., Ouyang J.M, 2013, Changes in urinary nanocrystallites in calcium oxalate stone formers before and after potassium citrate intake. International Journal of Nanomedicine, Vol. 8, p. 909-918.
2. Mahesh G, Ramkanth S, Saleem MTS. Anti-inflammatory drugs from medicinal plants A comprehensive review. International Journal of Review in Life Sciences 2011; 1(1): 1-10.



**Proceeding
The 1st International Conference
on Pharmaceutics & Pharmaceutical Sciences**

3. Tristiana Erawati M, Esti Hendradi, Widji Soeratri, 2014, Praformulation Study Of p-Methoxycinnamic Acid (PMCA) Nanoemulsion Using Vegetable Oils (Soybean Oil, Corn Oil, VCO), Int J Pharm Pharm Sci, ISSN-0975-1491, Vol. 6, Issue 2, 2014, 99-101
4. Widji Soeratri, Tristiana Erawati M., Diny Rahmatika, Noorma Rosita, 2014, PMCA Dose as Topical Antiinflammation and Its Penetration Study Through Full-thick Rat Skin and Without Stratum Corneum, Indonesian Journal of Pharmacy and Pharmaceutical Science, ISSN- 2406-9388. Vol. 1, No. 1, June 2014 p. 28-30