

Correlation between vertical components and skeletal Class II malocclusion in ethnic Javanese

by I Gusti Aju Wahyu Ardani

Submission date: 26-Dec-2018 03:30PM (UTC+0800)

Submission ID: 1060562298

File name: tion-between-vertical-components-and-skeletal-class-i-121818.pdf (1.39M)

Word count: 4653

Character count: 23681

Correlation between vertical components and skeletal Class II malocclusion in ethnic Javanese

I Gusti Aju Wahyu Ardani¹
Ira Willyanti²
Ida B Narmada¹

¹Orthodontic Department, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia;
²Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia

Introduction: The dentoskeletal morphology of Class II malocclusion has been analyzed in several cephalometric investigations. It is crucially important to understand the vertical components in orthodontic treatment. Defining the facial type of an individual is an essential key to obtain an accurate orthodontic diagnosis.

Aim: The aim of this study was to understand the correlation of vertical components in Class II skeletal malocclusion.

Materials and methods: This is an observational descriptive study. This study used lateral cephalograms of Class II skeletal malocclusion from Universitas Airlangga Dental Hospital, between April 2015 and 2016. Cephalometric analysis was performed using digital software by a single examiner. This analysis measured gonial (G_0), upper gonial (G_1), lower gonial (G_2), Yaxis-SN angle, the length of anterior face height, and posterior face height. The correlation between each variable was analyzed using the Pearson correlation test ($P < 0.01$).

Results: There was a correlation between vertical components and Class II skeletal malocclusion.

Conclusion: Vertical components were correlated with Class II skeletal malocclusion. The greater ANB angle will be followed by greater G_2 , which worsens the condition of Class II skeletal malocclusion.

Keywords: AFH, ANB, Class II skeletal malocclusion, G_1 , G_2 , PFH, Yaxis, Yaxis-SN

Introduction

The dentoskeletal morphology of Class II malocclusion has been analyzed in several cephalometric investigations. Defining the facial type of an individual is an essential key to obtain an accurate orthodontic diagnosis, and the lateral cephalogram is presented in literatures as a relevant resource for this purpose.^{1,2} Class II division 1 (Class II/1) malocclusion represents the most common skeletal discrepancy, which orthodontists see in daily practice.³ The understanding of this morphology is a key element in planning dentofacial orthopedic treatment for this type of malocclusion. Clinically widely accepted term of "skeletal Class II" does not specify whether the mandible is retruded in relation to the maxilla, or the maxilla is protruded in relation to the mandible.^{4,5}

The Class II/1 incisal relationship was found in association with a range of vertical skeletal patterns. Some studies reported an increase in lower facial height, whereas other studies reported that the lower facial height was significantly reduced. Class II division 2 (Class II/2) malocclusion is usually associated with an increase in posterior face height (PFH), a decrease in mandibular plane angle, a decrease in anterior face height (AFH), and a more horizontal growth vector.¹

Correspondence: I Gusti Aju Wahyu Ardani
Orthodontic Department, Faculty of Dental Medicine, Campus A Universitas Airlangga Jl Prof Dr Moestopo, No 47, Surabaya 60132, East Java, Indonesia
Tel +62 31 503 0255
Fax +62 31 502 0256
Email wahju_ardani@fkg.unair.ac.id

Skeletal class II malocclusion may be caused by the maxillary prognathism of maxillary depth or the angle formed between the Frankfurt plane and the N-A line, by mandibular retrograde, ie, reduced facial depth or angle formed between the Frankfurt plane and the facial plane (N-Pog), or by the combination of both conditions. As the causes of skeletal Class II malocclusion are varied, the treatment will also differ depending on the underlying cause.⁶

The difficulty in controlling vertical dimension during orthodontic treatment is a long-recognized fact. Therefore, it is crucial to thoroughly understand the vertical face height (VFH) in orthodontic treatment. Vertical dimension, generally speaking, includes AFH and PFH.⁷

Materials and methods

Sample

The preliminary research consisted of 465 Javanese (Deutro-Melayu) adult population who were seeking orthodontic treatment at the Universitas Airlangga Dental Hospital from April 2015 to 2016. From the total sample, there were 202 patients with skeletal Class I malocclusion (ANB 1–3), 171 patients with skeletal Class II malocclusion (ANB >4), and 92 patients with skeletal Class III malocclusion (ANB <0). There were only 106 skeletal Class II malocclusion patients who meet our eligibility criteria, with the age range of 15–35 years, of whom 21 patients were males and 85 patients were females. The study was performed in accordance with the Declaration of Helsinki, and human ethics approval was attained from the Faculty of Dental Medicine, Universitas Airlangga, Health Research Ethical Clearance Commission No 271/HRECC.FODM/X/2018. All patients provided a written informed consent to review their medical records as per the policy of Faculty of Dental Medicine, Universitas Airlangga. The inclusion criteria of sample selection were ANB >4.5, male or female patient with skeletal Class II malocclusion who had never done any orthodontic treatment before full permanent dentition.

Lateral cephalogram was taken in natural head position with subject closing their teeth in centric occlusion and lips in relax position. Cephalometric analysis was performed using the digital software OrthoVision (Ewoosoft Co., Ltd, Gyeonggi-do, South Korea) by a single examiner. The analysis started with determining the landmark anatomies shown in Figure 1, such as

1. SNA: angle formed from the points sella (S), nasion (N), and A. The range of normal values is 80°–84°.
2. SNB: angle formed from the points S, N, and B. The range of normal values is 78°–82°.




Figure 1 Anatomical landmarks used in this research.

Abbreviations: Go₁, upper gonial; Go₂, lower gonial; AFH, anterior face height; UAFH, upper anterior face height; LAFH, lower anterior face height; TAFH, total anterior face height; PFH, posterior face height.

3. ANB: angle between SBA and SNB. The range of normal values is 0°–4°.
4. Go₁: angle formed from the points articulare (Ar), Go₁, and N. The range of normal values is 52°–55°.
5. Go₂: angle formed from the points N, Go₂, and menton (Me). The range of normal values is 70°–75°.
6. Upper AFH (UAFH): a line measured from points N to anterior nasal spine (ANS).
7. Lower AFH (LAFH): a line measured from points ANS to Me.
8. Total AFH (TAFH): a line measured from points N to Me. This distance indicates a long or short AFH.
9. PFH: distance from the S–Go points. This distance indicates a long or short PFH.
10. Yaxis: angle formed from the Frankfurt horizontal (FH) plane with the S–gnathion (S–Gn) line is used to determine the direction of mandibular growth. The average value is 59.4°.
11. Yaxis-SN: angle formed from SN line with S–Gn line is used to determine the direction of mandibular growth. The average value is 66°.

Statistical analysis

The mean and SD of the measurement counted were determined using SPSS 17.0 (SPSS Inc., Chicago, IL, USA) for

Windows OS. The data distribution was calculated by means of Kolmogorov–Smirnov ($P>0.05$), while the correlation between each variable was analyzed using the Pearson correlation test.

Results

The mean and SD of each parameters are shown in Table 1. The correlation test in this study used Pearson correlation with statistical significant if $P<0.01$. Table 2 shows there was no correlation between SNA and Go_1 and LAFH, whereas a negative correlation was found between SNA and Go_2 , UAFH, AFH, Yaxis, and Yaxis-SN. There was a positive correlation between SNA and PFH. While there was no correlation between SNB and Go_1 , a positive correlation was found in SNB to PFH and a negative correlation was found in SNB to Go_2 , UAFH, LAFH, AFH, Yaxis, and Yaxis-SN. There was no correlation in the relationship between ANB and Go_1 , UAFH, LAFH, and AFH. A positive correlation was found between ANB and Yaxis and Yaxis-SN, whereas negative correlation was found between Go_2 and PFH.

There was a positive correlation between Go_1 and Go_2 , whereas a negative correlation was found between Go_1 and UAFH, LAFH, AFH, and PFH. Go_1 did not correlate with Yaxis and Yaxis-SN. Go_2 was positively correlated with LAFH, AFH, Yaxis, and Yaxis-SN and was negatively correlated with PFH. There was no correlation between Go_2 and UAFH. The UAFH was positively correlated with LAFH, AFH, PFH, and Yaxis-SN, and there was no correlation between UAFH and Yaxis. LAFH was positively correlated with AFH, PFH, Yaxis, and Yaxis-SN. AFH was positively correlated with PFH, Yaxis, and Yaxis-SN. PFH did not correlate with Yaxis but negatively correlated with Yaxis-SN. Yaxis was positively correlated with Yaxis-SN.

Discussion

This study discusses the relationship between SNA, SNB, ANB, Go_1 , Go_2 , UAFH, LAFH, AFH, PFH, Yaxis, and Yaxis-SN in the patients with skeletal Class II malocclusions at Universitas Airlangga Dental Hospital. First, we discuss the relationship between SNA, SNB, and ANB with Go_1 , Go_2 , AFH, PFH, and Yaxis-SN. Ricketts determined the mandibular growth arch as the guide to predict the mandibular growth. Gonial angle position moves posteriorly on this growth arch, a distance that equaled almost one half of the total increase in mandibular growth. This study used Yaxis and Yaxis-SN, because the FH plane can be used to analyze the lower dentofacial abnormalities and using only one reference plane can provide less valid data due to the variations in the FH.⁸ According to the results of this study, Yaxis-SN has more correlation because the growth of the lower face is not separated from the growth of the upper face, and according to a previous study, SN field is more stable.⁹ Yaxis-SN can be used to measure vertical growth associated with anterior

Table 1 Descriptive data of measurement variable

Variable	Minimum	Maximum	Mean±SD
SNA (°)	73.44	97.65	82.43±3.94
SNB (°)	60.51	87.50	75.56±4.18
ANB (°)	4.54	13.58	6.86±1.73
Go_1 (°)	37.74	59.42	46.97±4.75
Go_2 (°)	55.78	95.00	79.29±6.55
UAFH (mm)	36.46	62.41	51.22±4.47
LAFH (mm)	54.25	81.14	66.51±5.55
AFH (mm)	94.16	138.73	117.72±8.18
PFH (mm)	55.10	95.16	73.32±7.29
Yaxis (°)	57.93	84.48	70.99±5.05
Yaxis-SN (°)	62.82	91.85	73.29±4.59

Table 2 Pearson correlation table

Pearson (r/p)	Go_1	Go_2	UAFH	LAFH	AFH	PFH	Yaxis	Yaxis-SN
SNA	0.006/0.948	-0.318*/0.001	-0.299*/0.002	-0.132/0.177	-0.253*/0.009	0.239*/0.014	-0.303*/0.002	-0.745*/0
SNB	0.010/0.917	-0.453*/0	-0.240*/0.013	-0.196*/0.044	-0.264*/0.006	0.319*/0.001	-0.410*/0	-0.856*/0
ANB	-0.008/0.932	0.377*/0	-0.113/0.251	0.176/0.071	0.058/0.554	-0.237*/0.015	0.311*/0.001	0.376*/0
Go_1	-	0.206*/0.034	-0.267*/0.006	-0.255*/0.008	-0.319*/0.001	-0.303*/0.002	-0.069/0.483	-0.133/0.173
Go_2	0.206*/0.034	-	0.017/0.861	0.477*/0	0.333*/0	-0.398*/0	0.510*/0	0.682*/0
UAFH	-0.267*/0.0006	0.017/0.861	-	0.325*/0.001	0.767*/0	0.468*/0	-0.049/0.616	0.248*/0.010
LAFH	-0.255*/0.008	0.477*/0	0.325*/0.001	-	0.856*/0	0.390*/0	0.370*/0	0.369*/0
AFH	-0.319*/0.001	0.333*/0	0.767*/0	0.856*/0	-	0.521*/0	0.224*/0.021	0.386*/0
PFH	-0.303*/0.002	-0.398*/0	0.468*/0	0.390*/0	0.521*/0	-	-0.130/0.186	-0.309*/0.001
Yaxis	-0.069/0.483	0.510*/0	-0.049/0.616	0.370*/0	0.224*/0.021	-0.130/0.186	-	0.580*/0
Yaxis-SN	-0.133/0.173	0.682*/0	0.248*/0.010	0.369*/0	0.386*/0	-0.309*/0.001	0.580*/0	-

Note: *Correlation is significant at the 0.01 level (two tailed).

Abbreviations: Go_1 , upper gonial; Go_2 , lower gonial; AFH, anterior face height; UAFH, upper anterior face height; LAFH, lower anterior face height; TAFH, total anterior face height; PFH, posterior face height.

and posterior changes and may determine vertical inferior anterior growth of the mandible.¹⁰

In this study, SNA and SNB are negatively correlated with Go_2 . The mean value of SNA (82.43) is normal, whereas the mean of SNB (75.56) is smaller than normal; it shows that most of the cases are retrognathic mandible. There are many variations in the anteroposterior skeletal relationships in Class II division 2 malocclusion but mostly are retrognathic mandible.² The SNB (-0.453) correlation to Go_2 is stronger than that of SNA (-0.318). This indicates that patients with small SNB will have greater Go_2 angle, which mean that the patients tend to have a downward-backward mandibular rotation. Gonial angle size tends to be inversely related to and significantly correlated with mandibular body length and the size of angles SNA and ANB.⁸ Increase in gonial angle is accompanied by an increase in mandibular length. Björk described an increasing curvature of the mandibular base and noted that this increase was accompanied by a reduction in the size of the gonial angle. This reduction, in turn, was accompanied by vertical condylar growth, whereas both the gonial angle and sagittal growth increased.¹¹ Mandibular length was not all related to gonial angle size. The mandibular body length was significantly related and inversely proportional to the gonial angle size. That is, individuals with a small gonial angle tended to have a larger mandibular body length and the SNA angle and ANB angle were also larger in cases with a smaller gonial angle.⁸

SNA and SNB angles are negatively correlated with UAFH, LAFH, AFH, Yaxis, and Yaxis-SN. The smaller the angle SNB means the length of UAFH increases and the length of LAFH increases too; this will result in the increase in the length of AFH. The SNA angle significantly correlated with Yaxis-Sn (-0.745) meaning that the larger the SNA, the lower the Yaxis-SN value. There is a very significant correlation between ANB and Yaxis-SN (-0.856); the smaller the SNB, then the mandible tends to rotate downward-backward. The mean value of ANB (6.86) greater than normal is positively correlated with Go_2 , Yaxis, and Yaxis-SN meaning that the greater the ANB angle the mandible is likely to rotate downward-backward. There is no correlation between ANB and UAFH, LAFH, and AFH. The ANB angle is negatively correlated with PFH, indicating that the greater the angle of ANB the PFH reduced. With respect to the posterior height, it weakly associated with the actual mandibular length and maxilla length and moderately and positively correlated with the anteroposterior position of the mandible and maxilla. Schudy depicted the vertical and anteroposterior growth as opposing forces, which sounds reasonable with AFH being

regarded as vertical height in the extreme case of long or short face type. In relative normal overbite case, some studies showed that the vertical growth of PFH is consistent with anteroposterior growth.⁷

The mean value of Go_1 angle (46.97) is smaller than normal with a positive correlation with Go_2 ; the smaller the Go_1 angle, the smaller Go_2 angle too. In this study, the average value of Go_2 (79.29) is greater than normal probably due to downward-backward rotation of the mandible. The angle of Go_1 is negatively correlated with UAFH, LAFH, AFH, PFH, and Yaxis-SN but not correlated with Yaxis. This shows that the smaller the value of Go_1 , the larger the values of UAFH, LAFH, and AFH, causing downward-backward rotation of the mandible.

Figure 2 shows a patient with skeletal Class II malocclusion with ANB angle (12.93), Go_2 (92.9), and Yaxis-SN (91.85). This patient showed a great ANB angle and along with the rotation of the mandible. The mean angle value of Go_2 (79.29) is greater than normal negatively correlated with PFH. The Go_2 angle is positively correlated with LAFH, AFH, Yaxis, and Yaxis-SN. The correlation between Go_2 and Yaxis-SN is very significant (0.682), proving that Go_2 strongly reflects on the mandibular rotation. This means that the larger the angle of Go_2 then the greater the length of LAFH and AFH and the lower the length of PFH, causing downward-backward mandibular rotation. The steeper the




Figure 2 Class II skeletal malocclusion with great value of ANB, Go_2 , and Yaxis-SN.

mandibular plane anteriorly inclines, the greater the AFH becomes or vice versa.¹² There is no correlation between Go_2 angle and UAFH. The high angle ramus is shorter, whereas the low angle ramus tends to be longer.¹³

Vertical dimensional control is a difficult task on orthodontic treatment. It is important to understand VFH in orthodontic treatment.¹¹ Vertical dimensions include AFH and PFH. The ratio of UAFH/LAFH (<0.8) was obtained from the mean of UAFH (51.22) and LAFH (66.51), meaning that LAFH is longer. According to statistical results data in this study, the UAFH values were positively correlated with LAFH, AFH, PFH, and Yaxis-SN. This shows that the smaller the UAFH value then LAFH, AFH, PFH, and Yaxis-SN will decrease. UAFH is significantly correlated with AFH (0.767). UAFH is not correlated with Yaxis, whereas LAFH is positively correlated with AFH, PFH, Yaxis, and Yaxis-SN. The greater the LAFH value then the greater the AFH, PFH, Yaxis, and Yaxis-SN. LAFH is significantly correlated with AFH (0.856).

From the mean of AFH (117.72) and PFH (73.32), the ratio of AFH/PFH (62.2%) was obtained showing that, in this study, patients with skeletal Class II malocclusion tend to have a balanced PFH and AFH. AFH values are positively correlated with PFH, Yaxis, and Yaxis-SN. This indicates that the longer the anterior face, the greater the length of posterior face and the greater the angles of Yaxis and Yaxis-SN, resulting in a downward-backward rotation of the mandible. According to Björk, forward mandibular rotation occurs when PFH overdevelops relative to AFH; however, in many literatures, more attention was focused on the AFH and lower AFH has been confirmed as having a strong influence on the formation of vertical facial disproportions.¹⁴ In contrast, some accepted terminologies used in describing vertical morphology, such as long face and short face and dolichofacial and brachyfacial, are mainly based on AFH.¹⁵⁻¹⁷ In normal cases, long PFH is always followed by long AFH and vice versa.⁷ Graber¹⁸ found that if the gonial angle is acute, mandibular growth is more vertical and directed more anteriorly. The value of PFH is negatively correlated with Yaxis-SN, and the greater the PFH the Yaxis-SN value will increase indicating the forward rotation of the mandible. Yaxis angle itself is positively correlated with Yaxis-SN. The Yaxis-SN angle is a measurement of mainly vertical growth very sensible to anterior and posterior changes, which collaborates in defining the vertical of anterior inferior growth of the mandible, detecting its degree of opening.¹⁹

Cases with normal overbite, forward rotation of the mandible cannot be simply regarded as the growth result of

relatively long PFH together with relatively short AFH. In other words, the reason for mandibular forward rotation is not because of increase in PFH together with a decrease in AFH but different increased dimensions of it.⁷ These study results showed that it is the underdevelopment or overdevelopment of PFH instead of AFH that plays a key role in mandibular rotation. This agrees with Björk who stated that, under ideal circumstances, the fulcrum point for anterior or forward mandibular growth rotation is located at the incisors. Therefore, as the result of forward rotation, both the AFH and PFH increase.¹⁴

Thus, a face with a low angle, normally accompanied by a long ramus and a wide frontal region (wide bizygomatic and bigonial), can easily produce an illusion of a short face, although the anterior lower face height could be normal. This also explains why many faces with a low angle are taken for granted to be assumed as a short face, whereas a wide face and a square profile are responsible for making faces appear short.

Many literatures were more focused in the AFH and LAFH values as it has been confirmed to have a strong influence on the formation of vertical facial disproportions. Reducing facial height can be done by the extrusion of molars to compensate for backward rotation by increasing the forward mandibular rotation with minimal effect on anterior vertical facial dimensions. This may be applied during treating adolescent patient with skeletal Class II caused by small size of the ramus and body of the mandible. This relationship often results in a decreased PFH, a steeper mandibular plane angle, an increased ANB angle between SNA and SNB, a normal SNA angle with a decreased SNB angle, an increased angle of convexity, and an increased overjet.

Conclusion

Patients with Class II skeletal malocclusion in Javanese population do not always have a short anterior face. From the results of this study, the greater the ANB angle then the greater the value of Go_2 and this can be caused by the downward-backward rotation of the mandible. This mandibular rotation is evidenced by the increasing value of Yaxis-SN. As the value of Go_2 increases, the AFH value will also increase. This proves that Class II in Javanese population has a high anterior face. The increased AFH will aggravate Class II skeletal malocclusion. Orthodontic treatment in Class II skeletal malocclusion with high AFH is more difficult.

Acknowledgments

This work has not been published previously and is not under consideration by another journal and is the work of the

author(s). Research grant is funded by Universitas Airlangga no 373/UN3.14/LT/2018.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Al-Khateeb EA, Al-Khateeb SN. Anteroposterior and vertical components of class II division 1 and division 2 malocclusion. *Angle Orthod.* 2009;79(5):859–866.
2. Isik F, Nalbantgil D, Sayinsu K, Arun T. A comparative study of cephalometric and arch width characteristics of class II division 1 and division 2 malocclusions. *Eur J Orthod.* 2006;28(2):179–183.
3. Pachori Y, Navlani M, Gaur T, Bhatnagar S. Treatment of skeletal class II division 1 malocclusion with mandibular deficiency using myofunctional appliances in growing individuals. *J Indian Soc Pedod Prev Dent.* 2012;30(1):56.
4. Sidlauskas A, Svalkauskiene V, Sidlauskas M. Assessment of skeletal and dental pattern of class II division 1 malocclusion with relevance to clinical practice. *Stomatologija.* 2006;8(1):3–8.
5. Al-Azzawi AM, Sahib WW, Al-Khafaji TJ. Assessment of skeletal and dental pattern of the mandible of class II division 1 malocclusion [A comparative cephalometric study]. *kufa J Nurs Sci.* 2011;1:17–48.
6. Hoffelder LB, de Lima EM, Martinelli FL, Bolognese AM. Soft-tissue changes during facial growth in skeletal class II individuals. *Am J Orthod Dentofacial Orthop.* 2007;131(4):490–495.
7. Wang MF, Otsuka T, Akimoto S, Sato S. Vertical facial height and its correlation with facial width and depth. *Int J Stomatol Occlusion Med.* 2013;6(4):120–129.
8. de Rossi M, Stuani MBS, da SLAB. Cephalometric evaluation of vertical and anteroposterior changes associated with the use of bonded rapid maxillary expansion appliance. *Dental Press J Orthod.* 2010;15:62–70.
9. Ardani I, Sanjaya ML, Sjamsudin J. Cephalometric characteristic of skeletal class II malocclusion in Javanese population at Universitas Airlangga Dental Hospital. *Contemp Clin Dent.* 2018;9(Suppl 2):S342.
10. Steiner C. Cephalometries in clinical practice. *Angle Orthod.* 1959;29:8–29.
11. Kim YH. Overbite depth indicator with particular reference to anterior open-bite. *Am J Orthod.* 1974;65(6):586–611.
12. Obaidi HA. Variation of facial heights among the class I, II and III dentoskeletal relationships (Cephalometric study). *Al-Rafidain Dent J.* 2006:98–105.
13. Yassir YA. Ramus Height and its relationship with skeletal and dental measurements. *J Oral Dent Res.* 2013;1(1):4–8.
14. Björk A. Prediction of mandibular growth rotation. *Am J Orthod.* 1969;55(6):585–599.
15. Bishara SE, Saunders WB. *Textbook of Orthodontics.* Collingwood, ON: Saunders Book Company; 2001.
16. Bishara SE, Jakobsen JR. Longitudinal changes in three normal facial types. *Am J Orthod.* 1985;88(6):466–502.
17. Schudy FF. Vertical growth versus anteroposterior growth as related to function and treatment. *Angle Orthod.* 1964;34:75–93.
18. Graber LW. Chin cup therapy for mandibular prognathism. *Am J Orthod.* 1977;72(1):23–41.
19. Paranhos LR, Benedicto EN, Nunes MF, Kairalla SA, Siqueira DF, Torres FC. Correlation of different cephalometric measurements to define facial type. *Int J Orthod Milwaukee.* 2012;23(1):31.

Clinical, Cosmetic and Investigational Dentistry

Publish your work in this journal

Clinical, Cosmetic and Investigational Dentistry is an international, peer-reviewed, open access, online journal focusing on the latest clinical and experimental research in dentistry with specific emphasis on cosmetic interventions. Innovative developments in dental materials, techniques and devices that improve outcomes and patient satisfac-

tion and preference will be highlighted. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dentistry-journal>

Correlation between vertical components and skeletal Class II malocclusion in ethnic Javanese

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Kuitert, R.. "Dentoalveolar compensation in subjects with vertical skeletal dysplasia", American Journal of Orthodontics & Dentofacial Orthopedics, 200605
Publication 1%
- 2 O. Ronning, S. A. R. Barnes, M. H. Pearson, D. M. Pledger. "Juvenile chronic arthritis: a cephalometric analysis of the facial skeleton", The European Journal of Orthodontics, 1994
Publication 1%
- 3 www.lucalombardo.net
Internet Source 1%
- 4 dental.sumandeeepuniversity.co.in
Internet Source 1%
- 5 benthamopen.com
Internet Source 1%
- 6 Castelo, Keila Maria de Sousa, and Fausto Silva Bramante. "Determination of the structural characteristics in youngsters from

CearÃ; with Class II, division 1 malocclusion",
Dental Press Journal of Orthodontics, 2012.

Publication

7	Submitted to Universidad Estadual Paulista Student Paper	1%
8	doaj.org Internet Source	1%
9	Submitted to University of Brighton Student Paper	1%
10	Ma, Xiaoyang, Antonio J. Forte, John A. Persing, Nivaldo Alonso, Nicholas L. Berlin, and Derek M. Steinbacher. "Reduced Three-Dimensional Airway Volume Is a Function of Skeletal Dymorphology in Treacher Collins Syndrome :", Plastic & Reconstructive Surgery, 2015. Publication	1%
11	www.ilapeo.com.br Internet Source	1%
12	www.readbag.com Internet Source	<1%
13	repositorio.ufrn.br Internet Source	<1%
14	scidoc.org Internet Source	<1%

15	www.dentalpress.com.br Internet Source	<1%
16	Submitted to The University of Manchester Student Paper	<1%
17	www.ortoface.com Internet Source	<1%
18	Bajracharya, Manju. "Comparison of Maxillary and Mandibular Incisor Inclination in Class II Division 1 Malocclusion among Chinese Children", Orthodontic Journal of Nepal, 2014. Publication	<1%
19	www.jisppd.com Internet Source	<1%
20	H. L. Tippett, L. K. Dodgson, N. P. Hunt, M. P. Lewis. "Indices of extracellular matrix turnover in human masseter muscles as markers of craniofacial form--a preliminary study", The European Journal of Orthodontics, 2008 Publication	<1%
21	orthomaia.com.br Internet Source	<1%
22	czas.stomat.net Internet Source	<1%
23	S. Ishwarkumar, P. Pillay, M. R. Haffajee, K. S. Satyapal. "Morphometric analysis of the	<1%

mandible in the Durban Metropolitan population of South Africa", Folia Morphologica, 2017

Publication

24

www.pagepress.org

Internet Source

<1%

25

publicationethics.org

Internet Source

<1%

26

Zicari, Anna Maria, Marzia Duse, Francesca Occasi, Valeria Luzzi, Emanuela Ortolani, Flaminia Bardanzellu, Serena Bertin, and Antonella Polimeni. "Cephalometric Pattern and Nasal Patency in Children with Primary Snoring: The Evidence of a Direct Correlation", PLoS ONE, 2014.

Publication

<1%

27

Samir Aboul-Hosn Centenero, Federico Hernández-Alfaro. "3D planning in orthognathic surgery: CAD/CAM surgical splints and prediction of the soft and hard tissues results – Our experience in 16 cases", Journal of Cranio-Maxillofacial Surgery, 2012

Publication

<1%

28

www.medfak.ni.ac.rs

Internet Source

<1%

29

S. Murata. "Use of the centroid method of

<1%

occlusion for studying the vertical and horizontal relationship of the mandible and maxilla", The European Journal of Orthodontics, 09/28/2007

Publication

30

Jens Johannes Bock. "Evaluation of Vertical Parameters in Cephalometry", Journal of Orofacial Orthopedics / Fortschritte der Kieferorthopädie, 05/2007

Publication

<1%

31

Submitted to Tennessee State University

Student Paper

<1%

32

ir.uiowa.edu

Internet Source

<1%

33

www.ajronline.org

Internet Source

<1%

34

repositorio.pucrs.br

Internet Source

<1%

35

digitalcommons.wustl.edu

Internet Source

<1%

36

etd.uthsc.edu

Internet Source

<1%

37

Ahn, J.G.. "Cephalometric appraisal of posttreatment vertical changes in adult orthodontic patients", American Journal of

<1%

Orthodontics & Dentofacial Orthopedics,
200010

Publication

38

Al-Nimri, K.S.. "Changes in mandibular incisor position in Class II Division 1 malocclusion treated with premolar extractions", American Journal of Orthodontics & Dentofacial Orthopedics, 200312

Publication

<1%

39

Lee, Young-Joo, Jong-Tae Park, and Jung-Yul Cha. "Perioral soft tissue evaluation of skeletal Class II Division 1: A lateral cephalometric study", American Journal of Orthodontics and Dentofacial Orthopedics, 2015.

Publication

<1%

40

Arriola-Guillén, Luis Ernesto, Aron Aliaga-Del Castillo, Luis Fernando Pérez-Vargas, and Carlos Flores-Mir. "Influence of maxillary posterior discrepancy on upper molar vertical position and facial vertical dimensions in subjects with or without skeletal open bite", The European Journal of Orthodontics, 2015.

Publication

<1%

41

Noemi Mazzone, Claudio Matteini, Veronica Incisivo, Belli Evaristo. "Temporomandibular Joint Disorders and Maxillomandibular Malformations", Journal of Craniofacial

<1%

42

academic.oup.com

Internet Source

<1%

43

Kajiyama, K.. "Evaluation of the modified maxillary protractor applied to Class III malocclusion with retruded maxilla in early mixed dentition", American Journal of Orthodontics & Dentofacial Orthopedics, 200011

Publication

<1%

44

Arntsen, Torill, Inger Kjær, and Liselotte Sonnesen. "Lengths of the maxillary central incisor, the nasal bone, and the anterior cranial base in different skeletal malocclusions", Acta Odontologica Scandinavica, 2009.

Publication

<1%

45

Thiruvengkatachari, B.. "Comparison of rate of canine retraction with conventional molar anchorage and titanium implant anchorage", American Journal of Orthodontics & Dentofacial Orthopedics, 200807

Publication

<1%

Correlation between vertical components and skeletal Class II malocclusion in ethnic Javanese

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
