

The Influence of Moderate Exercise on Caspase-3 Expression in Inhibiting Transformation of Oral Squamous Epithelial Cells

by Pratiwi Soesilawati

Submission date: 04-May-2023 10:08AM (UTC+0800)

Submission ID: 2083620150

File name: The_Influence.pdf (292.43K)

Word count: 2519

Character count: 13338

The Influence of Moderate Exercise on Caspase-3 Expression in Inhibiting Transformation of Oral Squamous Epithelial Cells

Anis Irmawati^{1*}, Birgita Gina Pamita², Pratiwi Soesilawati¹

1. Department of Oral Biology, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.
2. Undergraduate student, Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia.

Abstract

The prevalence of cancer is increasing nowadays. In Indonesia, the prevalency of cancer is 1,4% out of 1000 people. Cancer is mainly caused by two factors; genetic and environmental, one of them being is a carcinogenic agent benzopyrene. Physical exercise can decrease blood glucose and fatty acid levels within the blood but its effect on cancer prevention, especially on caspase-3 expression, is still unknown.

This study examines the expression of caspase-3 produced by moderate exercise in preventing the formation of transform cell epithelial of the oral squamous cell.

18 mice were divided into three study groups; control group 1 (K1) not given any physical exercise and benzopyrene, control group 2 (K2) not given any physical exercise but induced with 0,08 mg benzopyrene, treatment group (P) were given physical exercise in moderate intensity and induced with 0,08 mg benzopyrene. Buccal mucosa tissue samples were taken and stained for immunohistochemistry and examined under a light microscope at 400x at 10 different angles.

The mean of caspase-3 expression was found highest in group P (13.33), meanwhile in group K1 and K2 were 3.00 and 8.33, respectively. There was significant difference of caspase-3 expression between the groups ($p = 0.000$) Conclusion: Moderate exercise could increase caspase-3 expression in preventing the formation of the transform cell epithelial of the oral squamous cell.

Experimental article (J Int Dent Med Res 2018; 11(1): pp. 285-288)

Keywords: Moderate exercise, transforms cell epithelial, oral squamous cell, caspase-3.

Received date: 27 September 2017

Accept date: 18 October 2017

Introduction

The prevalence of cancer today is increasing. Exposure to cancer risk factors is not only present in industrialized countries, but also in the whole world.¹ Cancer is considered the second leading cause of death globally. According to data from Basic Health Research in Indonesia (RISKESDAS) 2013, the prevalence of cancer in Indonesia is considerably high at 1.4% for every 1,000 residents.²

Cancer can be caused by a multitude of factors; from a person's genetics, environment, infection and lifestyle-including smoking.³ Smoking cigarettes causes the body to be

exposed to carcinogenic substances, the risk of developing cancer are in direct correlation with death.^{1,4} Since cigarette smokers inhale smoke into the lungs, the type of cancer that develop from smoking are lung cancers, cancer of the larynx and pharynx, and cancer of the oral cavity (oral cancer). Oral cancer is considered the cancer that occurs most often in cases of head and neck cancer worldwide with about 263,000 new cases each year. The most common case of oral cancer is squamous cell carcinoma that causes changes in the oral cavity and impacts basic human functions such as talking, swallowing, chewing, and saliva production. Toxic compounds contained in cigarette smoke consists of Reactive Oxygen Species (ROS), such as polycyclic aromatic hydrocarbons complex, one of which is benzo [a] pyrene (BaP).^{1,5,6}

¹⁸ Benzopyrene is metabolized into Benzo [a] pyrene-7,8-diol-9,10-epoxide (BPDE) and binds to one of DNA purine base, guanine.⁴ BPDE and DNA bond can cause DNA mutations

*Corresponding author:

Associate Professor, Anis Irmawati,
Department of Oral Biology, Faculty of Dental Medicine,
Universitas Airlangga, Jl. Prof. Moestopo 47,
Surabaya, Indonesia.
E-mail: anis-m@fkg.unair.ac.id

that stimulate the process of apoptosis in cells.⁷ Apoptosis is the mechanism of cell death, and can be divided into 2 types, one is programmed apoptosis which is physiological and the other is pathological apoptosis which is when there is damage to the cells due to disease or due to harmful agents. If there is interference in the apoptotic pathway function, it can cause diseases in humans such as cancer, autoimmune disorders, and other neurodegenerative diseases.⁸ Complex proapoptosis pathways such as Bax, and caspase groups may be induced by physical exercise. Regular physical exercise with appropriate intensity can prevent chronic diseases such as cancer, the risk of recurrence of cancer, diabetes, and heart attacks.^{9,10,11}

Physical exercise, is mostly effective if given moderate intensity. Physical exercise being undertaken tends to be physiological, which does not cause any interference with the balance at the cellular level to the organ level. Therefore, in this study, homeostasis at cellular and organ level will not be disturbed.^{7,12}

Effect of moderate exercise on caspase-3 expression in prevention of the formation of transformation on oral squamous epithelial cells is still not clearly, therefore, the aim of this study is to analyses the effect of moderate exercise on the caspase-3 expression in prevention of the formation of transform cells on oral squamous epithelial cells.

Materials and methods

This study had been approved with ethical clearance from Committee of Ethical Clearance of Health Research, Faculty of Dental Medicine, Universitas Airlangga, Indonesia (No: 164/KKEPK.FKG/VIII/2016).

This study was an experimental laboratory study with a sample of 18 mice (*Mus musculus*) of the Swiss Webster strain, which qualifies with the calculation using the Higgins-Kleinbaum formula. Each mice was male, aged \pm 2 months, weighs 25-35 grams, adapted for 1 week, and randomly divided into 3 groups; negative control group (K1), positive control group (K2), and treatment group (P).

The negative control group (K1) was soaked in water, with a time of 70% of the mice's maximum swimming capacity, 3 times per week for 12 weeks. In the 5th week 0.04 mL of oleum olivarum was induced on the top right buccal

mucosa, 3 times a week for 4 weeks. The positive control group (K2) is soaked in water, with a time of 70% of the mice's maximum swimming capacity, 3 times per week for 12 weeks. In the 5th week the mice were induced by 0.08 mg of benzopyrene (Merck, Sigma-Aldrich Pte. Ltd., Singapore) / 0.04 ml oleum olivarum on the top right buccal mucosa, 3 times a week for 4 weeks. Lastly, the treatment group (P) was given physical exercise with weights weighing 3% of the mice's body weight (BW), with a time of 70% of the maximum swimming capacity, 3 times per week for 12 weeks. In the 5th week induced by 0.08 mg of benzopyrene / 0.04 ml oleum olivarum on the top right buccal mucosa, 3 times a week for 4 weeks. At the beginning of the 13th week, mice were sedated with ether, waited until unconscious. The next stage, tissue on the top right buccal mucosa was removed in each group. Staining was done using the immunohistochemistry method, and observation of squamous epithelial cells expressing caspase-3 (are calculated on a 10 field of view with an magnification 400x).¹³

Statistic: The data collected was analyzed with Oneway Analysis of Variance (Anova) on SPSS 19.0 for Windows.

Results




Figure 1. The expression of caspase-3 (black arrow) on K1 (a), K2 (b) and P (c) groups with immunohistochemical staining (400x magnification). Arrow (\rightarrow) indicates cells expressing caspase-3.

The figure 1 showed the expression of caspase-3 by immunohistochemical slide. The mean of caspase-3 expression was found highest in group P (13.33), meanwhile in group K1 and K2 were 3.00 and 8.33, respectively (Figure 2). The cell that express caspase-3 showed brown color due to anti caspase-3 antibody absorption.

Oneway Anova test results of caspase-3 expression, shows that there are significant differences ($p < 0.01$). The Tukey Post Hoc Test

results also showed significant differences ($p > 0.01$) between all 3 groups, the results can be seen in Table 1.




Figure 2. The average of caspase-3 expression on K1, K2 and P groups.

Group	K1	K2	P
K1	-	$p = 0,000^*$	$p = 0,000^*$
K2			$p = 0,001^*$
P			-

Table 1. Tukey Post Hoc test of *caspase-3* expression on squamous epithelial cells.

Description: * showed that the data is significant.

Discussion

The expression of caspase-3 plays an important role in the change of normal cells into cancer cells in gastric mucosa.¹⁴ Results of the research conducted proved that moderate exercise can increase the expression of caspase-3 in prevention transformation cells on oral squamous epithelial cells. The expression of caspase-3 was higher in group P due to group P being given treatment in the form of physical exercise with weights about 3% of the mice's body weight (BW), with a time of 70% of the maximum swimming capacity, 3 times per week for 12 weeks.

Moderate exercise can increase the expression of caspase-3 because it can trigger Ca^{2+} channels opening in the membrane cell. The opening of the Ca^{2+} channels will cause an influx of Ca^{2+} into the cell cytoplasm resulting in an increase in the concentration of Ca^{2+} in the cell. The increased Ca^{2+} concentration in the

cells activates signal transduction pathways involved in various cellular responses-including proliferation, transcription, contraction, exocytosis, apoptosis, immune response, and neurotransmission- one of which enables complex signal transduction Ras and Src. Active Src and Ras as signal transduction and then activate the MAPK (mitogen-Activated Protein Kinase) in the nucleus. MAPK activation causes phosphorylation of various transcription factors and act as an intermediary in the process of signal transduction from the plasma membrane to in the nucleus.^{15,16}

Benzopyrene is a compounds contained in cigarettes can enter into the tissues of the body through the skin, inhalation, or through the mouth. Benzopyrene is metabolized in the liver by the cytochrome P4501A1 enzyme (CYP1A1) with the final result¹⁷ of the carcinogenic metabolites, that is benzo [a] pyrene-7,8-dihydrodiol 9,10-epoxide (BPDE). BPDE that enter the cell can bind to DNA, thus later will be responded by p53.¹⁷ This bonding causes DNA mutations of p53, which can hinder the process of protein synthesis (transcription of wild p53). MAPK activation causes phosphorylation of various transcription factors and act as an intermediary in the process of signal transduction from the plasma membrane to the nucleus. One of the proteins that undergo transcription is wild p53 protein. Wild p53 is a protein that acts as a tumor suppressor gene that inhibits the occurrence of cancer.⁴

Bcl-2 protein complexes play an important role in controlling apoptosis through the mitochondrial pathways. Bcl-2 complex is divided into two, namely apoptosis suppressor gene and gene promoting apoptosis. Bcl-2 acts as a suppressor gene, and the gene is acting as promoting Bax. Stimulation barriers Bax Bcl-2 by causing the release of cytochrome-c. The release of cytochrome-c from the mitochondria to the cytosol is the initial phase of the process of apoptosis.^{7,18} Cytochrome c binds to Apoptosis Protease-Activating Factor (Apaf-1). The complex formed then activates caspase-9 which is responsible for the apoptotic cascade. Activation of caspase-9 causes the activation of caspase-3. Activation of caspase-3 causes CAD (Caspase Activated DNase) endonuclease to be active. In the process of cell proliferation CAD, equipped with complex inhibitor, namely ICAD (Inhibitor Caspase Activated DNase). In the

process of apoptosis, caspase-3 ICAD serves to break down and release the CAD. CAD endonuclease then degrade chromosomal DNA in the nucleus and causes chromatin to condense and form apoptotic bodies.^{8,19} Apoptotic bodies formed does not cause the formation of squamous epithelial cells transform cells of the oral cavity.

Conclusions

Moderate intensity of physical exercise increases caspase-3 expression on the epithelial transform cell of the oral squamous cell.

Declaration of interest

The authors report no conflict of interest.

References

1. Sasco AJ. Cancer and Globalization. *Biomed Pharmacother.* 2008;62(2):110-121.
2. National Institute for Health Research & Development. Riset Kesehatan Dasar (National Health Survey). Ministry of Health Republic of Indonesia. 2013:1-303.
3. Purwaningsih NMS, Sailan AT, Jalil AA, Sinon SHM. Human papillomavirus detection in oral potentially malignant disorders and oral squamous cell carcinoma. *J Int Dent Med Res.* 2017;10(2):198-201.
4. Alexandrov K, Rojas M, Rolando C. DNA Damage by Benzo(A)Pyrene in Human Cells is Increased by Cigarette Smoke and Decreased by a Filter Containing Rosemary Extract, Which Lowers Free Radicals. *Cancer Res.* 2006;66(24):11938-45.
5. Ferlay J, Shin H, Bray F, Forman D, Mathers C, Parkin D. GLOBOCAN 2008 v2.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase. No 10 [Internet] Lyon, Fr Int Agency Res Cancer; 2010. Available from [Http://globocan.iarc.fr](http://globocan.iarc.fr). Accessed 18/Jan/2016 2010.
6. Barrios R, Tsakos G, Garcia-Medina B, Martinez-Lara I, Bravo M. Oral Health-Related Quality of Life and Malnutrition in Patients Treated for Oral Cancer. *Support Care Cancer Off J Multinatl Assoc Support Care Cancer.* 2014;22:2927-33.
7. Zhou G, Richardson M, Fazili IS, Wang J, Donnelly KC, Moorthy B. Role of Retinoic Acid in the Modulation of Benzo(A)Pyrene-DNA Adducts in Human Hepatoma Cells: Implications for Cancer Prevention. *Toxicol Appl Pharmacol.* 2010;249:224-30.
8. Rastogi RP, Richa, Sinha RP. Apoptosis: Molecular Mechanisms and Pathogenicity. *EXCLI J.* 2009;8:155-81.
9. Sharkey BJ. Health Benefits of Activity and Fitness. Sharkey, B.J. (ed.), *Fit. Heal.* 5th ed, Champaign, Ill., Hum. Kinet. 2002:13-30.
10. Kirshbaum MN. A Review of the Benefits of Whole Body Exercise During and After Treatment for Breast Cancer. *J Clin Nurs.* 2007;16:104-21.
11. Rajarajeswaran P, Vishnupriya R. Exercise in Cancer. *Indian J Med Paediatr Oncol.* 2009;30:61-70.
12. Cooper, KH. *Antioxidant revolution.* Nashville: Thomas Nelson. 1994:56-9.
13. Chaudhary M, Bohra S, Gupta R, Patil S. Comparison and prediction of the extent of lesion of oral squamous cell carcinoma. *J Int Dent Med Res.* 2012;5(2):77-84.

14. Kania J, Konturek SJ, Marlicz K, Hahn EG, Konturek PC. Expression of Survivin and Caspase-3 in Gastric Cancer. *Dig Dis Sci.* 2003;48:266-71.
15. Widmaier EP, Raff H, Strang KT. *Vander's human physiology: The mechanisms of body function.* London:McGraw Hill Higher Education. 2014:445-70.
16. Kieffer BL, Evans CJ. Opioid Tolerance-In Search of the Holy Grail. *Cell.* 2002;108:587-90.
17. Purwaningsih NMS, Sailan AT, Sinon SHM, Jalil AA. Role of p16 and p53 in oral potentially malignant disorders and oral squamous cell carcinoma: A study in Malaysia. *J Int Dent Med Res.* 2017;10(1):42-7.
18. Shaha C, Tripathi R, Mishra DP. Male Germ Cell Apoptosis: Regulation and Biology. *Philos Trans R Soc Lond B Biol Sci.* 2010;365:1501-15.
19. Elmore S. Apoptosis: A Review of Programmed Cell Death. *Toxicol Pathol.* 2007;35:495-516.

The Influence of Moderate Exercise on Caspase-3 Expression in Inhibiting Transformation of Oral Squamous Epithelial Cells

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

18 %
INTERNET SOURCES

17 %
PUBLICATIONS

0 %
STUDENT PAPERS

PRIMARY SOURCES

1 [docero.net](#) Internet Source **2** %

2 [jbuon.com](#) Internet Source **1** %

3 "42 Annual Congress of AOMSI Nagpur 16–18 Nov 2017", Journal of Maxillofacial and Oral Surgery, 2017 Publication **1** %

4 [core.ac.uk](#) Internet Source **1** %

5 [ejfs.springeropen.com](#) Internet Source **1** %

6 [academic.oup.com](#) Internet Source **1** %

7 [www.i-scholar.in](#) Internet Source **1** %

8 [www.nature.com](#) Internet Source **1** %

9	connectjournals.com Internet Source	1 %
10	www.science.gov Internet Source	1 %
11	en.wikipedia.org Internet Source	1 %
12	www.edoc.co.za Internet Source	1 %
13	www.thieme-connect.com Internet Source	1 %
14	Florian C. Kurschus. "Gold fluorescent annexin A5 as a novel apoptosis detection tool", Cytometry Part A, 2009 Publication	1 %
15	Hesty Wahyuningsih, Andina Putri Aulia. "Review: Effect of Red Cabbage Juice (<i>Brassica oleracea</i> var. <i>Capitata</i> f. <i>Rubra</i>) on SGPT Level", Borneo Journal of Medical Laboratory Technology, 2020 Publication	1 %
16	Zhenkuan Xu, Wenzhe Xu, Yan Song, Bin Zhang, Feng Li, Yuguang Liu. "Blockade of store-operated calcium entry alleviates high glucose-induced neurotoxicity via inhibiting apoptosis in rat neurons", Chemico-Biological Interactions, 2016	1 %

17	mobt3ath.com Internet Source	1 %
18	www.jaypeedigital.com Internet Source	1 %
19	qmro.qmul.ac.uk Internet Source	<1 %
20	www.febrasgo.org.br Internet Source	<1 %
21	"Posters", Journal of Gastroenterology and Hepatology, 10/2004 Publication	<1 %
22	K. TAKAHASHI, O. INANAMI, M. KUWABARA. "Effects of intracellular calcium chelator BAPTA-AM on radiation-induced apoptosis regulated by activation of SAPK/JNK and caspase-3 in MOLT-4 cells", International Journal of Radiation Biology, 2009 Publication	<1 %
23	Retno Indrawati Roestamadji, Nahdah Izdihar Nastiti, Meircurius Dwi Condro Surboyo, Anis Irmawati. "The Risk of Night Shift Workers to the Glucose Blood Levels, Saliva, and Dental Caries", European Journal of Dentistry, 2019 Publication	<1 %

24 Sun, Y.. "Differential caspase-3 expression in noncancerous, premalignant and cancer tissues of stomach and its clinical implication", *Cancer Detection and Prevention*, 2006
Publication <1 %

25 icps.conference.unair.ac.id
Internet Source <1 %

26 medrech.com
Internet Source <1 %

27 www.dovepress.com
Internet Source <1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Influence of Moderate Exercise on Caspase-3 Expression in Inhibiting Transformation of Oral Squamous Epithelial Cells

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
