

Characterization and Production of Polyclonal Antibody Anti Excretory Secretory Protein of Blastocystis sp

by Mufasirin Mufasirin

Submission date: 02-Apr-2020 02:16PM (UTC+0800)

Submission ID: 1287617829

File name: Characterization_and_Production_of....pdf (416.63K)

Word count: 1710

Character count: 8912

Characterization and Production of Polyclonal Antibody Anti Excretory Secretory Protein of *Blastocystis* sp

Briantono Willy Rendragraha, Lucia Tri Suwanti¹, Rahadju Ernawati, Mufasirin Mufasirin,
Setiawan K²ardarto, Wiwiek Tyasningsih, Soelih Estoe Pangestie, Heni Puspitasari and Septian Hakim Susantoputro
Faculty of Veterinary Medicine, Airlangga University, Kampus C Unair, Mulyorejo, Surabaya 60115, East Java, Indonesia.

³
(Received : July, 2019 250/19 Accepted : August, 2019)

Abstract

This study aims to produce and characterize polyclonal antibody anti excretory secretory (ES) protein of *Blastocystis* sp. ES protein profile was analyzed using SDS-PAGE and used to immunize 2 rabbits. Rabbit's serum were analyzed using indirect ELISA and Western Blot. The result showed that molecular weight of ES protein of *Blastocystis* sp was 40 and 50 kDa and the protein was immunogenic. Both ES protein and antibody anti ES of *Blastocystis* sp can be promoted as diagnostic kit.

Key words : *Blastocystis* sp, Excretory Secretory Protein, Polyclonal Antibody.

Blastocystis sp is a protozoan parasite that widely prevalent in many countries and it causes gastrointestinal symptoms such as diarrhea, nausea, vomiting, abdominal pain, irritable bowel syndrome, and urticaria (Ajjampur and Tan., 2016). This parasite can infect human (Roberts *et al.*, 2014) and various animals like amphibian, reptiles, bird, and mammals

(Alfellani *et al.*, 2013).

Excretory secretory (ES) protein of parasites have an important role as virulence factor, it can affect and control host immune system during infection and it can be used as a biomarker to detect the presence of parasite and status of the infection in infectious disease (Gomez *et al.*, 2015). This study aims to characterize and produce polyclonal antibody anti ES protein of *Blastocystis* sp.

Materials and Methods

Blastocystis sp was isolated from dungen of beef cattle in Bangkalan, Madura. The sample was cultured using yeast extract media (Mohammed *et al.*, 2015). The growing culture was confirmed with PCR examination using primers b11400 FORC (5'-GGA ATC CTC TTA GAG GGA CAC TAT ACA T-3') and b11710 REVC (5'-TTA CTA AAA TCC AAA GTG TTC ATC GGA C-3) from Badparva *et al* (2014).

Blastocystis sp was harvested from media and centrifuged with 10.000 rpm for 10 minutes. Pellets was washed 3 times with PBS and

¹Corresponding author : Email: lucia-t-s@fkh.unair.ac.id




Fig 1. PCR Product of *Blastocystis* sp. , M = marker (Pro-mega); 1 = Negative control; 2,3 = positive culture

incubated in 37°C overnight. The supernatant was as ES protein. The profile of ES protein was analyzed with SDS-Page and visualized with a digital camera.

Two male New Zealand rabbits were immunized with 250 µg of ES protein. Immunization was done 4 times. The serum was collected before and 7 days after immunization. Antibody production was verified by indirect ELISA based on value of optical density (OD) and Western Blot. The OD value between groups was analyzed using one-way ANOVA.




Fig 2. Profile of excretory secretory protein of *Blastocystis* sp in SDS-PAGE, M = marker; 1 = ES protein of *Blastocystis* sp

Results and Discussion

Blastocystis sp from feces of cattle was cultured with yeast extract media, after 3 days, the organism reproduced and reached a number 5×10^9 /ml. Based on PCR, the sample showed positive result with band in 310 bp (Fig 1) as described in Badparva *et al* (*loc cit*).

SDS PAGE of ES protein of *Blastocystis* sp showed that there were two protein bands with molecular weight; 40 kDa and 50 kDa (Fig 2). Fadl *et al* (2016) identified 22 proteins from whole proteins of *Blastocystis*, which ranged from 12 kDa to 200 kDa, include 40 kDa and 50 kDa. According to Nagel *et al.* (2015), the 50 kDa is Matrix metalloproteinase-9 (MMP-9), member of zinc-binding matrix metalloproteinases (MMP) which degrades extracellular matrix including gellatin (gelatinase B), collagen and elastin, which explained that MMP-9 exist as pro-enzyme around 92 kDa with an active form of around 84 kDa and turncuted around 50 kDa and it has important role in tissue invasion and has corelation with immune system, including potentiation of interleukin (IL)-8. Nourrisson *et al.* (2016) reported that rocombinant legumain had a size ~40 kDa. Legumain is a lysosomal protease that localizes on cell surface, cytoplasm, and central vacuole, this protein has




Fig 3. Immunoblotting of excretory secretory protein of *Blastocystis* sp with antibody anti ES, M = marker; 1,2 = antigen with Rabbit's Serum 1:2

several functions such as mediated pro-survival function for surface molecule and inflammation reaction in host cell (Wu *et al.*, 2009).

Antibody in rabbit's serum increased significantly after immunization with ES protein. Before immunization the OD was $0.0415^c \pm 0.0007$, 7 days after 3rd and 4th immunization, respectively the OD were, $1.6217^b \pm 0.2216$, and $2.2232^a \pm 0.0300$ respectively. It indicated that ES protein of *Blastocystis* sp was immunogenic.

In blotting product, antibody was reacted with 40 kDa and 50 kDa proteins (Fig 3). Indirectly that these proteins were immunogenic. Nagel *et al* (*loc cit*) also reported that 50 kDa protein reacted with blastocystosis patient serum and 50 kDa protein was found in *Blastocystis* sp subtype 1, 2, 3, 4, 5, 7, and 8.

Summary

Molecular weight of ES protein of *Blastocystis* sp is 40 kDa and 50 kDa and both of protein are immunogenic. ES protein of *Blastocystis* sp can be promoted as kit diagnostic to detect serum blastocystosis patient and antibody anti ES as kit diagnostic to detect coproantigen.

Acknowledgement

The authors thankful to the Dean of Faculty of Veterinary Medicine, and Institute of Tropical Disease Universitas Airlangga

References

Ajjampur S. S. R and K. S. W. Tan. (2016) Pathogenic mechanisms in *Blastocystis* spp. interpreting result from in vitro and in vivo studies. *Parasitol Int* **65**: 772-779

Alfellani, M. A., D. Taner-Mulia., A. S. Jacob., C. A. Imeede., H. Yoshikawa., C. R. Stensvold and C. G. Clark. (2013) Genetic diversity of *Blastocystis* in livestock and zoo animals. *Protist* **164**: 497-509

Badparva, E., J. Sadraee., F. Kheirandish and M Frouzandeh. (2014) Genetic diversity of human *Blastocystis* isolates in Khorramabad, Central Iran. *Iranian J Parasitol* **9(1)**: 44-49

Fadl, H. O., D. M. H. El-Akkad., D. S. Abd El-Fattah., H. A. El-Bolaky and S. O. El-Bassiouni. (2016) Study of the protein profiles of *Blastocystis* isolates from symptomatic and asymptomatic subjects. *Med. J. Cairo Univ.* **84(3)**: 349-353

Gomez, S., L. Adalid-Peralta., H. Palafox-Fonsesca., V. A. Cantu-Robles., X. Soberon., E. Sciutto., G. Fragoso., R. J. Bobes., J. P. Lacleite., Ld. P. Yauner and A. Ochoa-Leyva. (2015) Genome analysis of excretory/secretory proteins in *Taenia solium* reveals their abundance of antigenic region (AAR). *Scientific Reports* **5(9683)**: 1-10

Mohammed, T. S., N. M. Sulaiman and S. B. Kamal. (2015) Preparation simplified culture for culturing *Blastocystis hominis* parasite. *J Bio, Agric and Healthcare.* **5(20)**:112-115

Nagel, R., R. J. Traub., M. M. S. Kwan., and H. Bielefeldt-Ohmann. (2015) Blastocystis spesific serum immunoglobulin in patients with irritable bowel syndrome (IBS) versus healthy controls. *Parasit Vectors* **8(453)**: 1-13

Nourrisson, C., I.Wawrzyniak., A. Cian., V. Livrelli., E. Viscogliosi., F. Delbac and P. Poirier. (2016) On *Blastocystis* secreted cysteine proteases: a legumain activated cathepsin B increases paracellular permeability of intestinal Caco-2 cell monolayers. *Parasitology* **143**: 1713.

Roberts T, D Stark, J Harkness, and J Ellis (2014) Update on the pathogenic potential and treatment options for *Blastocystis* sp. *Gut Pathog.* **6(17)**: 1-9

Wu. B., J. yin., C. Texler., M. Roussel and K. S. W. Tan. (2009) *Blastocystis* legumain is localized on the cell surface, and specific inhibition of its activity implicates a pro-survival role for the enzyme. *J Biol Chem* **285(3)**: 1790-1798.

IVJ TARIFF FOR BOOK REVIEW

1. Two numbers of each book - one for sending to the reviewing expert and the other for IVJ Library - should be submitted.
2. A Review charge of Rs. 1000/- for the expert for each book should be paid in advance.
3. For the review of each book at least one advertisement as inner page Black & White full page for Rs.6000/-.
4. All payments should be by DD or Multicity cheque drawn in favour of the Editor, Indian Veterinary Journal - payable at Chennai.
5. Review of the book will be as per the Reviewing Expert, in not more than 2 pages of IVJ.
6. Review time - a minimum of one month from the date of receipt of the book and the payment.
7. Publication of the Review - Earliest issue immediately after the receipt of the review.

- Editor

Comparative Performance of TANUVAS Aseel, Gramapriya and Indigenous Desi Bird under Backyard Condition in Dharmapuri District	...	33
R.Thangadurai and P.S.Shanmugam		
Antigenotoxic Effect of Nano-quercetin on 7,12-dimethylbenz[a]anthracene Induced Genotoxicity in Sprague Dawley Rats	...	35
S.Shahana, R.Madheswaran, P.Balachandran, A.Arivuchelvan and G.A.Balasubramaniam		
Improving the Cell Wall Estimation in Starch Rich Cereal Grains	...	38
Sonali Prusty, S.S. Kundu, K.S. Bisitha, U.B. Sontakke and Vijay Kumar Sharma		
 CLINICAL AND FIELD ARTICLES :		
Management of Cutaneous Papilloma in a Labrador Dog – A Case Report	...	42
D.Sumathi, P.Ramesh, N.Pazhanivel, M.Sandya Bhavani, K.Amirinder Singh and M.G.Jayathangaraj		
Silicone Resin Plastination of Helminth Parasites for Preservation	...	44
S.T. Bino Sundar, S.Sivagnanam and Bhaskaran Ravi Latha		
A Rare Case of Cutaneous Angiofibroma in a Cow and its Surgical Treatment	...	46
R. Uma Rani and N. Pazhanivel		
Cutaneous Streptothricosis in a Jersey Cross Bred Cow and its Therapeutic Management	...	48
S.Saravanan and K.M. Palanivel		
Preslaughter Stress in Banyuwangi Cattle During Transport	...	50
Muhammad Thohawi Elziyad Purnama, Winda Kusuma Dewi, Shabrina Fauzia Prayoga, Nadia Marva Triana, Bondan Sigit Purnomo Aji, Faisal Fikri and Iwan Sahrial Hamid		
Trauma induced Primary Hypovolemic Shock in a Deer	...	52
K.P. Prabhakaran and R. Madheswaran		
The Protective effect of <i>Swietenia macrophylla</i> Extract Nanoparticles on Pulmonary Damage in Streptozotocin-Induced Diabetic Rats	...	54
S.A. Sudjarwo and G. Wardani		
A Case Report of <i>Babesia gibsoni</i> in Dog and its Therapeutic Management	...	57
Prabhavathy Harikrishnan, C.Jayanthi and M.Ranjith Kumar		
FLASH BACK - A Golden article from the collection of digitalised archival of IVJ	...	59
Author and Subject Index		69 & 70

Characterization and Production of Polyclonal Antibody Anti Excretory Secretory Protein of Blastocystis sp

ORIGINALITY REPORT

12%

SIMILARITY INDEX

9%

INTERNET SOURCES

8%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

espace.library.uq.edu.au

Internet Source

4%

2

jjmicrobiol.com

Internet Source

3%

3

Submitted to Universitas Airlangga

Student Paper

2%

4

www.aitvm.org

Internet Source

2%

5

ira.le.ac.uk

Internet Source

1%

6

Robyn Nagel, Rebecca J. Traub, Marcella M S Kwan, Helle Bielefeldt-Ohmann. "Blastocystis specific serum immunoglobulin in patients with irritable bowel syndrome (IBS) versus healthy controls", Parasites & Vectors, 2015

Publication

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On