

DAFTAR PUSTAKA

1. White PF. Update on ambulatory anesthesia. Can J Anesth 2005; 52(6): 1-10.
2. Bisri T. Seri Buku Literasi Anestesiologi: Ambulatory anesthesia. 2007.
3. Apfelbaum JL. Current controversies in adult outpatient anesthesia. ASA, 2005.
4. D. Janet Pavlin, MD et al. Factors Affecting Discharge Time in Adult outpatients.Ambulatory Anasthesia.1998
5. Lucio Trevisani et al. Post-Anaesthetic Discharge Scoring System to assess patient recovery and discharge after colonoscopy. World Journal of Gastrointestinal Endoscopy. 2013
6. Piergaspare Palumbo et al. Modified PADSS for monitoring outpatients discharge. Annali Italiani di Chirurgia.2012
7. White PF. Ambulatory anesthesia advances into the new millenium. Anesth Analg 2000; 90: 1234-35.
8. McCarthy DE. Outpatient anesthesia. Jax-Medicine Journal 1998.
9. Friedman Z, Chung F, Wong DT. Ambulatory surgery adult patient selection criteria-a survey of canadian anesthesiologists. Can J Anesth 2004; 51(5): 437-43.
10. Gupta A, Stierer T, Zuckerman R, Sakima N, Parker SD, Fleisher LA. Comparison of recovery profile after ambulatory anesthesia with propofol, isoflurane, sevoflurane and desflurane:a systematic review. Anesth Analg 2004; 98: 632-41.
11. White PF, Kehlet H, Neal JM, Schricker T, Carr DB, Carli F, et al. The role of the anesthesiologist in fast-track surgery: from multimodal analgesia to perioperative medical care. Anesth Analg 2007; 104: 1380-96.

12. Wennervirta J, Ranta SO, Hynynen M. Awareness and recall in outpatient anesthesia. *Anesth Analg* 2002; 95: 72-77.
13. Pavlin DJ, Rapp SE, Polissar NL, Malmgren JA, Koerschgen M, Keyes A. Factors affecting discharge time in adult outpatient. *Anesth Analg* 1998; 87: 816-26.
14. Pavlin DJ, Chen C, Penalosa DA, Polissar NL, Buckley FP. Pain as a factor complicating recovery and discharge after ambulatory surgery. *Anesth Analg* 2002; 95: 627-34.
15. Raeder JC, Steine S, Vatsgaard TT. Oral ibuprofen versus paracetamol plus codeine for analgesia after ambulatory surgery. *Anesth analg* 2001; 92: 1470-72.
16. White PF. The role of non-opioid analgesic techniques in the management of pain after ambulatory surgery. *Anesth Analg* 2002; 94: 577-85.
17. Gan TJ, Meyer TA, Apfel CC, Chung F, Davis PJ, Habib AS, et al. Society for ambulatory anesthesia guidelines for the management of postoperative nausea and vomiting. *Anesth Analg* 2007; 105: 161528.
18. Gan TJ, Franiak R, Reeves J. Ondansetron orally disintegrating tablet versus placebo for the prevention of postdischarge nausea and vomiting after ambulatory surgery. *Anesth Analg* 2002; 94: 11991200.
19. Dimich I, Lingham R, Gabrielson G, et al. Comparative hemodynamic effects of labetalol and hydralazine in the treatment of postoperative hypertension. *J Clin Anesth.* 1989;1:201–206.
20. Rebecca S Twersky,Beverly K. Philip. *Handbook of Ambulatory anesthesia*. 2008.
21. White PF, Song D. New criteria for fast-tracking after outpatient anesthesia: A comparison with the modified alderete scoring system. *Anesth Analg* 1999; 88: 1069-72.
22. Marshall SI, Chung F. Discharge criteria and complications after ambulatory surgery. *Anesth Analg* 1999; 88: 508-17.

23. Schreiner MS, Nicholson SC, Martin T, et al. Should children drink before discharge from day surgery? *Anesthesiology*. 1992;76:528–533.
24. Kearney R, Mack C, Entwistle L. Withholding oral fluids from children undergoing day surgery reduces vomiting. *Paediatr Anaesth*. 1998;8:331–336.
25. Practice Guidelines for Postanesthetic Care. A report by the American Society of Anesthesiologists Task Force on Postanesthetic Care. *Anesthesiology*. 2002;96:742–752.
26. Pavlin DJ, Pavlin EG, Gunn HC, et al. Voiding in patients managed with or without ultrasound monitoring of bladder volume after outpatient surgery. *Anesth Analg*. 1999;89:90–97.
27. Pavlin DJ, Rapp SE, Polissar NL, et al. Factors affecting discharge time in adult outpatients. *Anesth Analg*. 1998;87:816–826.
28. Rosseland LA, Stubhaug A, Breivik H. Detecting postoperative urinary retention with an ultrasound scanner. *Acta Anaesthesiol Scand*. 2002;46:279–282.
29. Lau H, Lam B. Management of postoperative urinary retention: a randomized trial of in-out versus overnight catheterization. *ANZ J Surg*. 2004;74:658–661.
30. Mulroy MF, Salinas FV, Larkin KL, et al. Ambulatory surgery patients may be discharged before voiding after short-acting spinal and epidural anesthesia. *Anesthesiology*. 2002;97:315–319.