

Kaempferia galanga L. Inhibiting Effect on Vascular Endothelial Growth Factor (VEGF) and Cyclooxygenase-2 (Cox-2) Expression on Endothelium of Chorioallantoic Membrane

by Juni Ekowati

Submission date: 21-Dec-2019 03:34PM (UTC+0800)

Submission ID: 1237723036

File name: Cox-2_Expression_on_Endothelium_of_Chorioallantoic_Membrane.pdf (112.37K)

Word count: 1704

Character count: 9206

Kaempferia galanga L. Inhibiting Effect on Vascular Endothelial Growth Factor (VEGF) and Cyclooxygenase-2 (Cox-2) Expression on Endothelium of Chorioallantoic Membrane

Iwan Sahrial Hamid¹, Juni Ekowati and Muhammad Thohawi Elziyad Purnama

Department of Veterinary Pharmacology, Faculty of Veterinary Medicine, Universitas Airlangga, Surabaya 60115, Indonesia.

(Received : March, 2019 107/19 Accepted : May, 2019)

Abstract

This study was aimed to examine the extract of *Kaempferia galanga* L. to inhibit the expression of VEGF and Cox-2 on the chorioallantoic membrane of embryonated chicken eggs (ECE). This study used 24 ECE which were divided into six groups: (C+) inoculated with solvent, (C-, T1, T2, T3 and T4) inoculated with bFGF 60 ng then (T1, T2, T3 and T4) treated with the extract doses of 30, 60, 90 and celecoxib 60 mg, respectively. Expression of VEGF and Cox-2 was observed by the immunohistochemical method. It was concluded that the extract of *Kaempferia galanga* L. can inhibit the expression of VEGF and Cox-2 on the chorioallantoic membrane.

Key words: *Kaempferia galanga* L, VEGF, Cox-2, celecoxib

Development of cancer cells was induced by activating some growth factors such as Vascular Endothelial Growth Factor (VEGF), cyclooxygenase-2 (Cox-2) and tumor necrosis factor (TNF) (Prior *et al.*, 2004). *Kaempferia galanga* L contains Ethyl p-methoxycinnamate (EPMC), which has anti-inflammatory and anti-angiogenesis action through tyrosine kinase (Ekowati *et al.*, 2015). The study was conducted to assess these inhibitory effect of EPMC, on Cox-2, VEGF on Chorioallantoic membrane of embryonated eggs.

Materials and Methods

This study was approved by the ethical committee No: 923-KE, Faculty of Veterinary Medicine, Universitas Airlangga. Nine day old specific pathogen free (SPF) eggs from Pusat Veteri-

¹Corresponding author. Email : iwan-s-h@fkh.unair.ac.id

aria Farma, Surabaya. *Kaempferia galanga* L. isolated from Malang, East Java. Angiogenesis inducer used recombinant human Basic Fibroblast Growth Factor (bFGF) (Nako Japan Pure Chemical Industries Ltd.) diluted with Tris-HCl pH 7.5.

24 nine day old embryonated eggs were divided into six groups of 4 each. The positive control C+ eggs were inoculated with Tris-HCl; the negative control eggs are inoculated with DMSO solvent. The treatment groups T1, T2, T3 and T4 were inoculated with EPMC at doses of 30, 60, 90 and 120 mg in 2% DMSO and incubated at 38°C with 60% RH for 72 hr (Ribatti, 1997). The VEGF and Cox-2 expression were observed by immunohistochemical methods (Purnama *et al.*, 2019).

Results and Discussion

Observation of VEGF and Cox-2 expression was carried out on brownish-looking endothelial cells. The expression was caused by an antigen reaction with VEGF and Cox-2 antibodies as markers. The results (Table I) showed significant ($p < 0.05$) on VEGF expression after being inoculated with (T2) *Kaempferia galanga* L dose of 60 mg. Cox-2 expression was significant ($p < 0.05$) after being inoculated with (T1) *Kaempferia galanga* L dose of 30 mg.

The VEGF expression shows a brownish color in the cell cytoplasm that binds to the complex with VEGF receptor ligands (Chen *et al.*, 2016). The VEGF receptor bonds with specific ligands occur in the transmembrane domain and cytoplasm. The VEGF receptor is an inducer of angiogenesis and regulates the

Table I. Mean and standard deviation of the VEGF and Cox-2 expression

Treatments	Mean±SD	
	VEGF	Cox-2
C-	48.60 ^a ±9.34	69.60 ^d ±9.73
C+	26.60 ^c ±4.44	35.40 ^e ±4.32
T1	23.40 ^{bc} ±4.28	25.40 ^b ±3.05
T2	17.40 ^{ab} ±5.32	16.20 ^a ±4.09
T3	13.80 ^a ±3.29	15.80 ^a ±5.13
T4	11.85 ^a ±2.77	13.60 ^a ±9.99

Different superscripts in the same column indicate significant differences among treatments ($p < 0.05$)

integrity of the endothelium. Some of VEGF compounds inhibit endothelium formation and decrease angiogenesis (Di Marco *et al.*, 2009).

One of the factors that trigger the growth of angiogenesis when tumors occur, namely VEGF which initiates the formation of blood vessels (Red-Horse and Ferrara, 2006). Ethyl p-methoxycinnamate acts to inhibit angiogenesis by decreasing VEGF expression. Quercetin and Ethyl p-methoxycinnamate contained in *Kaempferia galanga L* inhibits angiogenesis and tumor necrosis factors. Quercetin can suppress tumor growth in vitro and in vivo acting as a barrier to the activity of tyrosine kinase (Kubo *et al.*, 2004).

Other active compounds in *Kaempferia galanga L* extracts such as alpha sitosterol, silymarin, 4-hydroxy-4-methyl-2-pentanone, quercetin and resveratrol are considered to have anticancer activities through a mechanism of signal transduction, inhibiting cell cycle, apoptosis, and inhibiting metastasis (Umar *et al.*, 2012). Resveratrol and quercetin at a concentrations of 100 µg/ml were able to inhibit the proliferation of angiogenesis, migration of endothelial cells, and formation of intima vascular layers (Igura *et al.*, 2001). Silymarin was developed as a cyclooxygenase-2 (Cox-2) enzyme inhibiting agent. This compound reduces the number of Human Vascular Endothelial Cells (HUVEC) at levels of 50 µg/ml (Tosetti *et al.*, 2002).

The compounds contained in *Kaempferia galanga L* extract can inhibit Cox-2 activity induced by bFGF angiogenic agents. Flavonoid

extract from *Kaempferia galanga L* decreased VEGF receptor (VEGFR2) through barrier activity from Matrix Metallo Proteinase (MMP), tyrosine kinase, and Cox-2 (Masferrer *et al.*, 2000; Hamid *et al.*, 2018).

Summary

Kaempferia galanga L EPMC extract at 60 mg can inhibit VEGF expression in chorioallantoic membrane blood vessel endothelial cells induced by bFGF. A minimum dose of 30 mg can inhibit Cox-2 expression. There was no significant difference in the administration of extracts of at the dosage of 90 mg with celecoxib 60 mg.

Acknowledgment

The authors acknowledge The Institute of Research and Innovation, Universitas Airlangga for providing fund support to carry out this study.

References

- Chen, L., Li, Z.Y., Xu, S.Y., Zhang, X.J., Zhang, Y., Luo, K. and Li, W.P. (2016) Upregulation of miR-107 inhibits glioma angiogenesis and VEGF expression. *Cell. Mol. Neurobiol.* **36**(1): 113-120.
- Di Marco, G.S., Reuter, S., Hillebrand, U., Amler, S., König, M., Larger, E. and Brand, M. (2009) The soluble VEGF receptor sFlt1 contributes to endothelial dysfunction in CKD. *J. Am. Soc. Nephrol.* **20**(10): 2235-2245.
- Ekowati, J., Hardjono, S. and Hamid, I.S. (2016) Ethyl p-methoxycinnamate from *Kaempferia galanga* inhibits angiogenesis through tyrosine kinase. *Universa. Med.* **34**(1): 43-51.
- Hamid, I.S., Aksono, E.B., Sukmanadi, M. and Purnama, M.T.E. (2018) Antiangiogenesis activity test of tin leaf (*Ficus carica L.*) on the number of blood vessels and VEGF expression of chorioallantoic membrane of embryonated chicken eggs. *Eur. J. Oncol. Pharm.* **1**(4): e00007.
- Igura, K., Ohta, T., Kuroda, Y. and Kaji, K. (2001) Resveratrol and quercetin inhibit angiogenesis in vitro. *Cancer letters.* **171**(1): 11-16.
- Kubo, I., Nihei, K.I. and Shimizu, K. (2004) Oxidation products of quercetin catalyzed by mushroom tyrosinase. *Bioorg. Med. Chem.* **12**(20): 5343-5347.
- Masferrer, J.L., Leahy, K.M., Koki, A.T., Zweifel, B.S., Settle, S.L., Woerner, B.M. and Seibert, K. (2000) Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res.* **60**(5): 1306-1311.
- Prior, B.M., Yang, H.T. and Terjung, R.L. (2004) What makes vessels grow with exercise training?. *J. App. Physiol.* **97**(3): 1119-1128.

Purnama, M.T.E., Rahmaningtyas, I.H., Pratama, A.R., Prastika, Z., Kartikasari, A.M. and Cahyo, N.P.D. (2019) Tadpole serum activity (*Rana catesbeiana*) in caspase-3 as a marker of the role of apoptosis and total cytotoxic T lymphocytes in Albino rats epithelial cells induced by neoplasia, *Vet. World*. **12**(1): 63-67.

Red-Horse, K. and Ferrara, N. (2006) Imaging tumor angiogenesis. *J. Clin. Investig.* **116**(10): 2585-2587.

Ribatti, D., Gulantris, A., Bastaki, M., Vacca, A., Lurlaro, M., Roncali, L. and Presta, M. (1997) New Model for the study of Angiogenesis and Antiangiogenesis in the Chick Embryo

Chorioallantoic Membrane. The Gelatin Sponge/Chorioallantoic Membrane Assay, *J. of Vascular Res.* **34** : 455-463.

Tosetti, F., Ferrari, N., De Flora, S. and Albini, A. (2002) 'Angioprevention': angiogenesis is a common and key target for cancer chemopreventive agents. *FASEB J.* **16**(1): 2-14.

Umar, M.I., Asmawi, M.Z., Sadikun, A., Atangwho, I.J., Yam, M.F., Altaf, R. and Ahmed, A. (2012) Bioactivity-guided isolation of ethyl-p-methoxycinnamate, an anti-inflammatory constituent, from *Kaempferia galanga* L. extracts. *Mol.* **17**(7): 8720-8734.

Indian Vet. J., September 2019, 96 (09) : 82 - 83

Pathology of Snake Envenomation in a Cat

M.S.Sreelakshmi¹, Mammen, J. Abraham, N.D. Nair, B.Abeena and R.B.Vishnurahav

Department of Veterinary Pathology, College of Veterinary and Animal Sciences, KVASU, Mannuthy, Thrissur- 680651, Kerala.

(Received : December, 2018 **444/18** Accepted : May, 2019)

Abstract

The present paper provides an account of gross and histopathological observations of viper snake envenomation in a cat.

Key words: cat, envenomation, histopathology, snakebite

Snakes and snakebites have been extensively studied because of the diversity of these species and venom. Annually, more than one lakh animal deaths occur as a result of venomous snake bites (Banga *et al.*, 2009). Eventhough, the problems associated with the poisonous snakes are more frequent in domestic animals, it is less common in cats as they avoid snakes (Sandhu and Brar, 2009).

Materials and Methods

A two year old male non-descript cat carcass was presented for post-mortem examination in the Department of Veterinary Pathology, College of Veterinary and Animal Sciences, Mannuthy with a clinical history of restlessness, salivation and death. A detailed necropsy was conducted, gross lesions were recorded and

representative tissue samples were collected in 10% neutral buffered formalin. Tissue sections were routinely processed and paraffin embedded tissues were sectioned at 5 μ and stained with Haematoxylin and Eosin (Luna, 1972).

Results and Discussion

A detailed necropsy examination revealed a small wound with oedema and haematoma on the medial aspect of right thigh region. On removal of the skin in the area there was oedema and degeneration of subcutaneous muscle (Fig. 1). Grossly, lung showed congestion, oedema and subpleural haemorrhages, liver was enlarged and congested. Subepicardial suffusions were



Fig. 1. Haematoma formation around snake bite wound

¹Corresponding author. Email: smsvet06@gmail.com

Kaempferia galanga L. Inhibiting Effect on Vascular Endothelial Growth Factor (VEGF) and Cyclooxygenase-2 (Cox-2) Expression on endothelium of Chorioallantoic Membrane

ORIGINALITY REPORT

11%

SIMILARITY INDEX

6%

INTERNET SOURCES

8%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.elsevier.es Internet Source	2%
2	T. K. Lim. "Edible Medicinal and Non-Medicinal Plants", Springer Nature, 2016 Publication	2%
3	Sekiya, S.. "Bioengineered cardiac cell sheet grafts have intrinsic angiogenic potential", Biochemical and Biophysical Research Communications, 20060310 Publication	1%
4	pubs.sciepub.com Internet Source	1%
5	scholarbank.nus.edu.sg Internet Source	1%
6	repository.lib.cuhk.edu.hk Internet Source	1%
7	cellbiosignal.jp Internet Source	

1%

8 link.springer.com
Internet Source

1%

9 Xia, Huan, Yang Li, and Xiaohong Lv.
"MicroRNA-107 inhibits tumor growth and metastasis by targeting the BDNF-mediated PI3K/AKT pathway in human non-small lung cancer", International Journal of Oncology, 2016.
Publication

1%

10 Tumor Angiogenesis, 2008.
Publication

1%

11 Tae-Jung Kim, Youn Soo Lee, Jin-Hyung Kang, Yeon-Sil Kim, Chang Suk Kang. "Prognostic significance of expression of vegf and cox-2 in nasopharyngeal carcinoma and its association with expression of C-erbB2 and EGFR", Journal of Surgical Oncology, 2011
Publication

1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Kaempferia galanga L. Inhibiting Effect on Vascular Endothelial Growth Factor (VEGF) and Cyclooxygenase-2 (Cox-2) Expression on Endothelium of Chorioallantoic Membrane

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
