

January-April 2015 | Vol.34 No.1

# UNIVERSA MEDICINA

<http://www.univmed.org>




- Zinc supplementation improves heme biosynthesis in rats exposed to lead
- *Centella asiatica* increases B-cell lymphoma 2 expression in rat prefrontal cortex
- Soursop leaf extract increases neuroglia and hepatic degeneration in female rats
- Induction of *Plasmodium falciparum* strain 2300 dormant forms by artemisinin
- Fetal blood vessel count increases in compensation of hypoxia in premature placentas
- Ethyl *p*-methoxycinnamate from *Kaempferia galanga* inhibits angiogenesis through tyrosine kinase
- Lipiodol retention pattern predicts transarterial chemoembolization therapeutic effect in hepatocellular carcinoma
- Family support is not a risk factor of negative self-esteem in HIV/AIDS women
- High toluene exposure risk increases risk of olfactory dysfunction in furniture workers

pISSN: 1907-3062 / eISSN: 2407-2230



# UNIVERSA MEDICINA

eISSN: 2407-2230  
pISSN: 1907-3062



USER

Username:   
Password:   
 Remember me

- HOME
- ABOUT
- LOGIN
- REGISTER
- SEARCH
- ANNOUNCEMENTS
- AUTHORS
- REVIEWERS
- EDITORIAL BOARD
- CONTACT US
- PUBLICATION
- ETHICS

JOURNAL CONTENT

Search

Search Scope  
All

Browse

- By Issue
- By Author
- By Title

Current Issue

Home > Editorial Board

## Editorial Board

### Editor in Chief

Adi Hidayat (Trisakti University, Indonesia)

### Deputy Editor in Chief

Pusparini (Trisakti University, Indonesia)

### Associate Editors

Julius E. Surjawidjaja (Trisakti University, Indonesia)

Nugroho Abikusno (Trisakti University, Indonesia)

Yenny (Trisakti University, Indonesia)

INFORMATION

- For Readers
- For Authors
- For Librarians

FONT SIZE

## For Authors

Review Article



Original Article



Case Report



## Journal Statistics

00191436 View My Stats



## Tools



CURRENT ISSUE

AHEAD OF PRINT

MOST CITED ARTICLES

ARCHIVES

RETRACTION & ERRATA

USEFUL LINKS

### Editorial Board

Elly Herwana (Trisakti University, Indonesia)

Edhyana Sahiratmadja (Padjadjaran University, Indonesia)

Mulyoto Pangestu (Monash University, Australia)

Dhananjay Yadav (Yonsei University, Korea)

Murad Lesmana (Trisakti University, Indonesia)

Roslida Abd Hamid (Putra Malaysia University, Malaysia)

Gulay Yilmazel (Hitit University, Turkey)

Muchtaruddin Mansyur (University of Indonesia, Indonesia)

Hans Joachim Freisleben (University of Indonesia, Indonesia/Germany)

Ruvan A I Ekanayaka (Institute of Cardiology, Sri Lanka)

Widyasari Kumala (Trisakti University, Indonesia)

Erlangga Yusuf (Universitair Ziekenhuis Antwerpen, Belgium)

Elvina Karyadi (The Micronutrient Initiative, Canada)

Rachel Soemedi (Brown University, United States)

Fatimah Binti Ibrahim (University of Malaya, Malaysia)

Indra Prasad P. Tripathi (Faculty of Science and Environment, India)

Saddek Bouharati (Setif University, Algeria)

Toni Wandura (Sari Mutiara University, Indonesia)

Daryoush Babazadeh (Shiraz University, Iran)

Xin Sun (West China Hospital of Sichuan University, China)

Farhad Safarpour Dehkordi (University of Tehran, Tehran, Iran)

Last updated on November 4, 2019



Universa Medicina by Faculty of Medicine, Trisakti University is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. Based on a work at <https://univmed.org/ejurnal/index.php/medicina/>.


### CALL FOR PAPPERS



Universa Medicina warmly welcome the contributions from you about new basic and clinical (including modern and epidemiological) research on medicine and health

Read more

### ACCREDITATION :

Directorate General of Research and Development Reinforcement Ministry of Research, Technology and Higher Education

Number 21/KPT/2018


Volume 35 Number 1, 2016 to Volume 39 Number 3, 2020





# UNIVERSA MEDICINA

eISSN: 2407-2230  
pISSN: 1907-3062



USER

Username   
Password   
 Remember me

- HOME
- ABOUT
- LOGIN
- REGISTER
- SEARCH
- ANNOUNCEMENTS
- AUTHORS
- REVIEWERS
- EDITORIAL BOARD
- CONTACT US
- PUBLICATION
- ETHICS

JOURNAL CONTENT

Search

Search Scope  
All

Browse

- By Issue
- By Author
- By Title

Vol 34, No 1 (2015)

Home > Archives > Vol 34, No 1 (2015)

## Vol 34, No 1 (2015)

DOI: <http://dx.doi.org/10.18051/UnivMed.2015.v34.1>

### Table of Contents

#### Review Article

<b>Benefits of smoking cessation for coronary heart disease patients</b> <i>Adi Hidayat</i>	PDF 1-2
<b>Zinc supplementation improves heme biosynthesis in rats exposed to lead</b> <i>Budi Santoso, Hertanto Wahyu Subagio, Lisyani Suromo, Henna Rya Sunomo</i>	PDF 3-9
<b>Centella asiatica increases B-cell lymphoma 2 expression in rat prefrontal cortex</b> <i>Kuswati Kuswati, Djoko Prakosa, Brian Wasita, Nanang Wiyono</i>	PDF 10-16
<b>Soursop leaf extract increases neuroglia and hepatic degeneration in female rats</b> <i>Ety Sri Handayani, Zainuri Sabta Nugraha, Prilly I Raleka Pahlevawati</i>	PDF 17-24
<b>Induction of Plasmodium falciparum strain 2300 dormant forms by artemisinin</b> <i>Lilik Maslachah, Yoes Prijatna Dachlan, Chairul A. Nidom, Loeki Enggar Fitri</i>	PDF 25-34
<b>Fetal blood vessel count increases in compensation of hypoxia in premature placentas</b> <i>K Kartini, Ahmad A Jusuf, Sri Widia A Jusman, M Ekawati, Ani R Prijanti</i>	PDF 35-42
<b>Ethyl p-methoxycinnamate from Kaempferia galanga inhibits angiogenesis through tyrosine kinase</b> <i>Juni Ekowati, Suko Hardjono, Iwan Sahrial Hamid</i>	PDF 43-51
<b>Lipiodol retention pattern predicts transarterial chemoembolization therapeutic effect in hepatocellular carcinoma</b> <i>Margaretha Vianny, Gunawan Santosa, Eddy Soedjianto</i>	PDF 52-60
<b>Family support is not a risk factor of negative self-esteem in HIV/AIDS women</b> <i>Jean Valeria, Surilena Surilena, Yanto Budiman, Samsuridjal Djauzi, Haridana Indah</i>	PDF 61-67
<b>High toluene exposure risk increases risk of olfactory dysfunction in furniture workers</b> <i>Magdalena Wartono, Herkutanto Herkutanto, Niken Lestari</i>	PDF 68-76

CURRENT ISSUE

AHEAD OF PRINT

MOST CITED ARTICLES

ARCHIVES

RETRACTION & ERRATA

USEFUL LINKS

#### CALL FOR PAPERS



Universa Medicina warmly welcome the contributions from you about new basic and clinical (including modern and epidemiological) research on medicine and health

Read more

#### ACCREDITATION :

Directorate General of Research and Development Reinforcement Ministry of Research, Technology and Higher Education

Number 21/KPT/2018

Volume 35 Number 1, 2016 to  
Volume 39 Number 3, 2020



Universa Medicina by Faculty of Medicine, Trisakti University is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.  
Based on a work at <https://univmed.org/ejournal/index.php/medicina/>.


#### INFORMATION

- For Readers
- For Authors
- For Librarians

#### FONT SIZE

### For Authors

#### Review Article



#### Original Article



#### Case Report



### Journal Statistics

00191263 View My Stats



### Tools



## Journal Profile

# Universa Medicina

eISSN : 2407-2230 | pISSN :

[Health](#) [Science](#)

[Universitas Trisakti](#)



S2

Sinta Score



Indexed by GARUDA

17

H-Index

14


H5-Index

1489

Citations

1159

5 Year Citations



*Penerbit:*

Fakultas Kedokteran Universitas Trisakti

---

[Website](#) | [Editor URL](#)

---

*Address:*

Fakultas Kedokteran Universitas Trisakti (kampus B) Jl. Kyai Tapa No. 260 – Grogol Jakarta  
Jakarta

---

*Email:*

---

*Phone:*

(021) 5672731 ext.2504

---

Last Updated :

2020-02-21

---



## Ethyl *p*-methoxycinnamate from *Kaempferia galanga* inhibits angiogenesis through tyrosine kinase

Juni Ekowati\*, Suko Hardjono\*, and Iwan Sahrial Hamid\*\*

### ABSTRACT

#### BACKGROUND

Many tumors express on their receptor tyrosine kinases vascular endothelial growth factor activity associated with angiogenesis. Inhibition of angiogenesis through reduction of tyrosine kinase activity is a promising strategy for cancer therapy. The present study aimed to determine the mechanism and potency of ethyl *p*-methoxycinnamate (EPMC) isolated from *Kaempferia galanga* as angiogenesis inhibitor.

#### METHODS

A laboratory experimental study was conducted using chorio-allantoic membranes (CAMs) of nine-day old chicken eggs induced by 60ng basic fibroblast growth factor (bFGF). Ethyl *p*-methoxycinnamate (EPMC) potency was determined at dosages of 30, 60, 90 and 120 µg and compared with celecoxib 60 µg as reference drug and one negative bFGF-induced control group. Neovascularization and endothelial cell count in CAM blood vessels were evaluated. To predict the antiangiogenic mechanism of EPMC, a docking study was performed with the Molegro Virtual Docker program on tyrosine kinase as receptor (PDB 1XKK).

#### RESULTS

Angiogenesis stimulation by bFGF was prevented significantly ( $p < 0.05$ ) by EPMC at dosages of 30, 60, 90 and 120 µg and this activity was dose dependent. Molecular docking showed interaction between EPMC functional groups and tyrosine kinase amino acids at Met766, Met793, Thr854, Thr790, Gln791 and Ala743. There was an association between EPMC antiangiogenic activity and docking study results.

#### CONCLUSIONS

Ethyl *p*-methoxycinnamate is a potential new angiogenesis inhibitor through interaction with tyrosine kinase. EPMC could be a promising therapeutic agent for treatment of angiogenesis-related diseases.

**Keywords:** Ethyl *p*-methoxycinnamate, chorio-allantoic membrane, angiogenesis, tyrosine kinase

\*Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Airlangga University, Surabaya

\*\*Department of Molecular Pharmacology, Faculty of Veterinary Medicine, Airlangga University, Surabaya

#### Correspondence

Dr. Juni Ekowati, Apt., M.Si  
Department of Pharmaceutical Chemistry,  
Faculty of Pharmacy,  
Airlangga University  
Kampus B Unair  
Jl. Dharmawangsa Dalam  
Surabaya 60286  
Email: j\_ekowati@yahoo.com

Univ Med 2015;34:43-51  
DOI: 10.18051/UnivMed.2016.v35.43-51  
pISSN: 1907-3062 / eISSN: 2407-2230

This open access article is distributed under a Creative Commons Attribution-Non Commercial-Share Alike 4.0 International License

## **Etil *p*-metoksisinamat dari *Kaempferia galanga* menghambat angiogenesis melalui interaksi dengan tirosin kinase**

### **ABSTRAK**

#### **LATAR BELAKANG**

Banyak tumor menunjukkan pada reseptor tirosin kinasenya ekspresi vascular endothelial growth factor yang berhubungan dengan angiogenesis. Hambatan angiogenesis melalui pengurangan aktivitas tirosin kinase adalah strategi yang menjanjikan untuk terapi kanker. Penelitian ini bertujuan untuk menentukan mekanisme dan potensi etil *p*-metoksisinamat (EPMC) sebagai penghambat angiogenesis.

#### **METODE**

Sebuah studi eksperimental laboratorium dilakukan menggunakan chorio-allantoic membrane (CAM) telur ayam tertunas berumur sembilan hari, yang diinduksi dengan basic fibroblast growth factor (bFGF) 60ng. Potensi EPMC diteliti pada dosis 30, 60, 90 dan 120 µg; dibandingkan dengan celecoxib 60 µg sebagai obat referensi dan satu grup tanpa perlakuan. Neovaskularisasi dan sel endotel pembuluh darah baru dari CAM dihitung dan dievaluasi. Untuk prediksi mekanisme antiangiogenesis EPMC, dilakukan studi doking pada reseptor tirosin kinase (PDB 1XKK) menggunakan program Molegro Virtual Docker v.5.5

#### **HASIL**

Stimulasi angiogenesis oleh bFGF pada CAM dihambat secara signifikan ( $p < 0.05$ ) oleh EPMC pada dosis 30, 60, 90 and 120 µg dan bersifat dose dependent. Celecoxib dan EPMC tersebut menyebabkan terjadinya lisis sel endotel dari CAM. Studi doking menunjukkan adanya interaksi antara gugus fungsi pada EPMC dengan residu asam amino tirosin kinase pada Met766, Met793, Thr854, Thr790, Gln791 dan Ala743. Studi doking dan aktifitas antiangiogenesis EPMC tersebut menunjukkan hasil yang berhubungan.

#### **KESIMPULAN**

Etil *p*-metoksisinamat merupakan senyawa yang berpotensi sebagai angiogenesis inhibitor melalui hambatan pada tirosin kinase.

**Kata kunci :** Etil *p*-metoksinamat, korio allantois membran, angiogenesis, tirosin kinase

## **INTRODUCTION**

Tyrosine kinase receptors of many tumors express vascular endothelial growth factor (VEGF) activity connected with angiogenesis.<sup>(1)</sup> In the last decade, angiogenesis has been explored in depth as an interesting cancer therapeutic target, since angiogenesis is an important step in tumor growth and cancer metastasis.<sup>(2,3)</sup>

Drug resistance, increased tumor progression, and signs of drug toxicity such as bleeding, fatigue, hypertension and

gastrointestinal perforation are the main clinical problems that occur in patients treated with angiogenesis inhibitors.<sup>(1)</sup> Thus, the use of natural herbal alternative agents to constrain angiogenesis is quite crucial in cancer treatment. Ethyl *p*-methoxycinnamate (EPMC), a major constituent of *Kaempferia galanga* Linn. (local name: *kencur*, Fam. *Zingiberaceae*) has been used as sunscreen,<sup>(4)</sup> analgesic<sup>(5)</sup> and anti-inflammatory.<sup>(6,7)</sup> agent, as cyclooxygenase-2 (COX-2) inhibitor, and to treat fibrosarcoma in mice.<sup>(8)</sup> It is has been established that PGE<sub>2</sub> production in preclinical breast and colon cancer



models is strongly influenced by COX-2 expression. COX-2 inhibition augments VEGF pathway blockade to destroy angiogenesis and tumor growth, avoid metastasis, and increase overall patient survival.<sup>(9-11)</sup>

It has also been reported that EPMC exerts anti-angiogenic activity in zebrafish<sup>(12)</sup> and can block endothelial function.<sup>(6)</sup> One method for the study of anti-angiogenic activity is by means of the chorio-allantoic membrane (CAM) model. The CAM is an extra-embryonic membrane formed on day 4 of incubation by fusion of the chorion and the allantois. The CAM contains extracellular matrix proteins that mimic the physiological cell environment and vascularization of cancers. Use of CAM has the advantages of easy accessibility and low cost, and is less time consuming.<sup>(13,14)</sup> The present study was done because there have been no reports of the potency of EPMC to inhibit angiogenesis in the CAM model.

He et al.<sup>(12)</sup> reported that EPMC influences multiple molecular targets of angiogenesis, therefore we evaluated the possibility of EPMC-mediated tyrosine kinase inhibition by performing a virtual molecular docking study using the Molegro Virtual Docker program (PDB 1XKK). The objectives of our study were to determine the antiangiogenic potency of EPMC and to predict whether the mechanism of action of this compound is related to its interaction with tyrosine kinase.

## METHODS

### Research design

This laboratory experimental study was conducted from May 2014 until September 2014. The research was performed in three steps, i.e. molecular docking study, isolation of EPMC from *Kaempferia galanga* and antiangiogenic assay using the CAM model.

### Docking study

Molecular docking studies of EPMC, celecoxib and the reference ligand FMM\_91[A]

(lapatinib tosylate) into the three-dimensional tyrosine kinase structure were carried out using Molegro Virtual Docker (MVD) software version 5.5. The receptor tyrosine kinase structure was obtained from the Protein Data Bank (PDB 1XKK). Ligand preparation was done using ChemBio Ultra software version 10.0.; the geometry was optimized by the MMFF 94 method and saved in Sybyl Mol2 format. All ligands were placed into the binding site of 1XKK (cavity-1) by alignment with the reference ligand FMM\_91[A] and 10 independent runs were performed.<sup>(15,16)</sup> The binding affinity between ligand and enzyme (docking score) was predicted using rerank scores. Higher negative scores (kcal/mol) indicate stronger ligand-enzyme binding, with the ligand filling the same pocket in the enzyme as in FMM\_91[A] docking. Validation of the docking study was carried out by redocking FMM\_91[A] into cavity-1 of 1XKK. The best docking results can be observed visually by comparing the structure of the docked molecules with the crystal structure of FMM\_91[A] into the binding site.

### Isolation of ethyl-*p*-methoxycinnamate from *Kaempferia galanga*

*Kaempferia galanga* Linn. was collected and determined by the botanist at Materia Medica Botanical Garden Park at Batu (near Malang), Indonesia, in August 2013. EPMC was isolated from *Kaempferia galanga* Linn. by extraction of the dried powder of the rhizomes (1kg) with petroleum ether until the filtrate was nearly transparent.<sup>(6)</sup> The extract was filtered and concentrated in a Rotavapor Model Laborota 4000 vacuum evaporator (Heidolph Instruments, Germany) at 25°C, and then kept in the freezer. The extract was carefully washed with *n*-hexane, and the *n*-hexane supernatant was then decanted. This process was repeated until the solvent was almost transparent.<sup>(6)</sup> The residue that remained after washings consisted of pale yellow, needle-like crystals, which were recrystallized with ethanol 70%, yielding white pure EPMC crystals (10.2 g), with a sharp melting point of 49°C, as

observed using a Fisher-John melting point apparatus. The chemical structure was characterized spectroscopically using a Jasco Fourier transform infra red (FT-IR) 5300 spectrophotometer and a nuclear magnetic resonance (NMR) Hitachi R-1900 spectrometer, confirming its identity as EPMC.

The IR and NMR spectroscopy results were as follows: IR (KBr) 3007, 2980, 2935, 1705, 1630, 1602, 1253, 1209, 829  $\text{cm}^{-1}$ ; [ $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 3.82 (3H, s,  $1 \times \text{OCH}_3$ ); 6.96 (1H, d,  $J = 15.9\text{ Hz}$ ,  $1 \times \text{CH}$  alkene); 6.88 (2H, d,  $J = 8.8\text{ Hz}$ ,  $2 \times \text{CH}$  benzylic); 6.96 (1H, d,  $J = 15.9\text{ Hz}$ ,  $1 \times \text{CH}$  alkene); 7.46 (2H, d,  $J = 8.8\text{ Hz}$ ,  $2 \times \text{CH}$  benzylic); 4.24 (2H, q,  $1 \times \text{CH}_2$ ); 1.32 (3H, t,  $J = 7.2\text{ Hz}$ ,  $1 \times \text{CH}_3$ ). [ $^{13}\text{C}$  ( $^1\text{H}$ ) NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$ : 14.4 ( $\text{CH}_3$ ), 55.4 ( $\text{OCH}_3$ ), 60.4 ( $\text{CH}_2$ ), 114.4 (benzylic  $\text{CH}_2$ ), 115.8 (CH alkene), 129.8 (Ar-C), 144.3 (CH alkene), 161.2 (Ar-C), 167.4 (carbonyl C)].

#### Antiangiogenic activity of EPMC

Twenty-four nine-day old chicken eggs were obtained from PUSVETMA Surabaya, and incubated at  $37^\circ\text{C}$  and 60-70% humidity. After 1 d of incubation, a  $1\text{ cm}^2$  window was opened at the top of the egg and the air was aspirated from the air chamber with a rubber ball until the CAM came in contact with the eggshell membrane. On all CAMs of the experimental groups, induction of new blood vessels was performed with bFGF 60 ng dissolved in 60 mL of stock rh bFGF at a concentration of 1 ng/mL in a laminar air flow hood under aseptic conditions. The eggs were distributed in six groups; four groups were treated with EPMC at dosages of 30, 60, 90 and 120  $\mu\text{g}$ ; one group was treated with celecoxib 60  $\mu\text{g}$  as the positive control group, and one group was left without any treatment except bFGF blood vessel induction as the negative control group. To verify EPMC antiangiogenic activity, a CAM assay was accomplished with certain modifications as described before.<sup>(13,17)</sup> Sterile 5-mm diameter Whatman qualitative filter-paper disks containing bFGF alone, or supplemented with

EPMC (at dosages of 30, 60, 90 and 120  $\mu\text{g}$ ) or with celecoxib 60  $\mu\text{g}$ , were implanted onto the CAM of each embryo. The window was resealed with adhesive tape and the eggs were returned to the incubator until day 11 of chick embryo development. After that, the upper eggshell was removed and neovascularization as positive parameter was characterized by capillary branching of the main blood vessels onto the bFGF-containing paper disks. The capillaries attached at the edges and on the upper and lower surfaces of the paper disks. The neovascularization parameter was considered to be negative if no capillary blood vessel formation occurred on the edges and upper and lower surfaces of the disk. Those capillaries within 2,5 mm around the filter paper disk were observed and photographed. Neovascularization around the disk was counted by determining the number of capillary branching of the main CAM blood vessels onto the disk. In addition to macroscopic observation as mentioned above, we also carried out microscopic observations in the following way: the CAM was removed from the shell, 10% formalin was added as preservative, and the CAM was then prepared into histological slides and stained with Hematoxylin and Eosin (HE).<sup>(13,17,18)</sup> Endothelial cell formation in the neovascular capillaries was observed in CAM cross-sections using an inverted Nikon H600L phase contrast microscope. Endothelial cell counts were determined in five visual fields of each slide at 400x magnification and compared with positive and negative controls for subsequent analysis.

#### Statistical analysis

Significant differences between the treatment and control group were assessed by one-way analysis of variance (ANOVA), supported by LSD test. Differences were considered significant at  $p < 0.05$ .

#### Ethical clearance

This assay was approved by the Ethical Commission of Airlangga University.




Figure 1. Interaction of EPMC (A), celecoxib (B), and FMM\_91 [A] (C) with amino acids residue of tyrosine kinase (PDB 1XKK)

## RESULTS

The results of the molecular docking study (rerank scores) were -87.84 kcal/mol (EPMC); -92.3184 kcal/mol (celecoxib); -143.23 kcal/mol (FMM\_91[A]), in which the best dock of EPMC, celecoxib and FMM\_91[A] into tyrosine kinase was conducted in cavity-1 of the receptor. Their interaction with tyrosine kinase (cavity-1 PDB 1XKK) is shown in Figure 1. Lapatinip tosylate displayed hydrogen bonding interaction between the following pairs: oxygen of ether moiety vs. Leu792; nitrogen of amino moiety and oxygen of furan ring vs Thr790; nitrogen of quinazoline ring vs. Lys745; oxygen of sulphonyl moiety vs. Gln791 and Lys852. Celecoxib presented hydrogen bonding interaction of the sulphonamide moiety with Ala743, Gln791 and Leu792. However, it showed steric and hydrophobic interaction of the trifluoromethyl moiety with Leu844, of the aromatic ring with Lys745, Ala743, and Met793, and of the sulphonamide moiety with Val742, Gln791 and Leu792 amino acid residues. EPMC showed

hydrogen bonding interaction of the oxygen moiety with Met793, and steric and hydrophobic interaction of the Met766, Thr854, Thr790, Gln791 and Ala743 of tyrosine kinase. It was concluded that functional groups of EPMC can interact with some tyrosine kinase amino acids, similar to those of celecoxib and lapatinip tosylate.

Treatment of CAM bFGF-induced neovascularization with celecoxib 60  $\mu\text{g}$  caused endothelial cell lysis of the blood vessels and reduced CAM neovascularization. Celecoxib suppressed CAM neovascularization by  $83.4 \pm 5.3\%$  and growth of endothelial cells in capillary branches by  $76.8 \pm 2.2\%$ . The potency of EPMC as angiogenic inhibitor was dose-dependent. At doses of 30, 60, 90 and 120  $\mu\text{g}$ , EPMC inhibited neovascularization by  $50.3 \pm 4.5\%$ ,  $67.6 \pm 4.9\%$ ,  $77.6 \pm 6.3\%$ , and  $81.9 \pm 3.2\%$  ( $p < 0.05$ ) respectively. At the doses of 30, 60, 90 and 120  $\mu\text{g}$ , EPMC inhibited endothelial cell counts in capillary branches of CAM by  $50.9 \pm 4.1\%$ ,  $55.4 \pm 2.9\%$ ,  $66.1 \pm 4.7\%$ , and  $73.2 \pm 4.5\%$ , respectively ( $p < 0.05$ ) (Table 1).

Table 1. Antiangiogenesis activity of EPMC by treatment group using CAM model

Treatment	% inhibition of neovascularization	% inhibition of endothelial cells	P
Celecoxib 60 $\mu\text{g}$	$83.4 \pm 5.3$	$76.8 \pm 2.2$	0.008
EPMS 120 $\mu\text{g}$	$81.9 \pm 3.2$	$73.2 \pm 4.5$	0.021
EPMS 90 $\mu\text{g}$	$77.6 \pm 6.3$	$66.1 \pm 4.7$	0.009
EPMS 60 $\mu\text{g}$	$67.6 \pm 4.9$	$55.4 \pm 2.9$	0.014
EPMS 30 $\mu\text{g}$	$50.3 \pm 4.5$	$50.9 \pm 4.1$	0.037

All values are represented as the mean  $\pm$  SD (n=4) and indicate significant differences compared with the control-untreated group at  $p < 0.05$




Figure 2. (A) Capillary of neovascularization of CAM induced by bFGF 60ng (white arrow), some of capillary blood vessels moving into the disk implanted on CAM; (B) Capillary of neovascularization of CAM induced by bFGF 60ng and treated with celecoxib 60  $\mu$ g (white arrow), none of capillary blood vessels moving into the disk implanted on CAM (black arrow); (C) capillary of neovascularization of CAM induced by bFGF 60 ng and treated with EPMC 60  $\mu$ g (white arrow), none of capillary blood vessels moving into the disk implanted on CAM (black arrow)

The endothelial cell counts of CAM bFGF-induced capillaries in the negative control group were higher than in the EPMC-treated groups (Figure 2).

Treatment of CAM with EPMC caused lysis of the endothelial cells (Figure 3). The higher EPMC doses caused more endothelial cell lysis and decreased CAM neovascularization more strongly. EPMC as an antiangiogenic agent at doses of 30 and 60  $\mu$ g was less active compared to celecoxib 60  $\mu$ g. There were a statistically significant differences in antiangiogenic potency between EPMC doses of 90 and 120  $\mu$ g. There were also a significant

differences between celecoxib 60  $\mu$ g and EPMC at doses 90 and 120  $\mu$ g ( $p < 0.05$ ).

## DISCUSSION

Angiogenesis plays an important role in delivering oxygen and nutrients essential for tumor progression and invasion of surrounding tissues and vessels. The development of a new vascular network consists of several steps which are controlled by various endogenous stimulators and inhibitors.<sup>(18)</sup> Antiangiogenic therapy has great potential for developing new cancer treatment methods in the future.<sup>(19)</sup>



Figure 3. Photomicrograph images of endothelial cells of CAM using an inverted phase contrast microscope (Nikon H600L; camera DS Fi2 300 megapixel) at 400 magnification. CAM induced by b bFGF (black arrow), the morphology of endothelial cells is good. (B) CAM induced by b FGF and treated by celecoxib 60  $\mu$ g. Some of the endothelial cells are lyzed (white arrow). (C) CAM induced by b FGF and treated by EPMC 60  $\mu$ g. It also appears that some of the endothelial cells are lyzed (white arrow)

He et al.<sup>(12)</sup> found by semiquantitative reverse transcription polymerase chain reaction assay that EPMC influences multiple molecular targets connected to angiogenesis. A recent study, showed the chemopreventive effect of EPMC against fibrosarcoma in mice and EPMC's capability to interact with COX-2.<sup>(8)</sup> The results of our molecular docking study on EPMC's interaction with the tyrosine kinase receptor (PDB 1XKK) predict the probable mechanism of EPMC as a cancer chemopreventive agent.

Molecular docking is a virtual method, in which the ligand is placed on the binding site in such a manner as to obtain the lowest potential energy. The docking process consists of testing the coordinate area of the target binding mode and scoring each possible ligand pose into that site. The lowest value of rerank scores is expected as the best binding mode.<sup>(15)</sup> Our results are in line with those of previous studies and tyrosine kinase is one of the molecular targets of EPMC for inhibition of angiogenesis. Altun et al.<sup>(20)</sup> reported that tyrosine kinase inhibitors are excellent agents for treating colorectal cancer and angiogenesis in a CAM model.

It is known that COX-2 may indirectly provoke angiogenesis by increasing the production of angiogenic factors, such as vascular endothelial growth factor (VEGF).<sup>(18,19)</sup> Our study also supports the antiangiogenic activity of the COX-2 inhibitor celecoxib in the CAM assay. Several reports revealed that there is a significant association between COX-2 expression and VEGF-C expression. The combination of celecoxib with tamoxifen is a potential drug for treatment of breast cancers expressing high levels of VEGF.<sup>(18)</sup> Celecoxib as COX-2 inhibitor has been reported to suppress angiogenesis and growth of secondary bone tumors.<sup>(19)</sup> Celecoxib overturns the expression of VEGF by constraining the production of PGE<sub>2</sub> and the stimulation of some signal transduction pathways by which the expression of VEGF is upregulated.<sup>(21)</sup>

Basic FGF as a proangiogenic agent causes chronic inflammation that markedly increases the

risk for the development of cancer, and therefore angiogenesis and VEGF signaling might pose an important link between inflammation and tumor development.<sup>(19,20)</sup> Histopathological analysis showed the presence of inflammatory cells on CAM induced by bFGF. Even though not directly proinflammatory, bFGF synergistically forces inflammatory mediator-induced leukocyte recruitment, thus indicating the relationship between inflammation and angiogenesis.<sup>(22,23)</sup>

Angiogenesis is triggered by increased arachidonic acid production as a result of prostaglandin PGE<sub>2</sub> metabolism catalyzed by COX-2. Prostaglandin affects vascularization and the cancer metastatic process that encourages the expression of proangiogenic factors, such as VEGF and fibroblast growth factor (FGF).<sup>(24)</sup> A previous research study reported that EPMC blocked COX-2 expression in the T47D cell line and caused lysis of that cell line.<sup>(25)</sup> This fact strengthens our prediction of the ability of antiinflammatory substances like EPMC to block the starting step of angiogenesis. Therefore, the present study was designed to investigate the potential angiogenesis inhibitory effect of EPMC on CAM. The outcomes showed that EPMC is a potent antiangiogenic substance, therefore it can be used as a cancer chemopreventive agent, in agreement with our earlier report. Although the antiangiogenic activity of EPMC is lower than that of celecoxib, EPMC can be used as lead compound for developing agents with higher antiangiogenic activities. EPMC can also be combined with other antiangiogenic drugs, e.g. bevacizumab, or with COX-2 inhibitors, such as celecoxib, so reducing the dose and decreasing the side effects of the drugs.<sup>(1,9)</sup>


One limitation of this study is the virtual experimental study of interactions between EPMC and tyrosine kinase, which eliminated the lipophilic effect of the compound. So, in our future research the experiment will be conducted by *in vitro* studies based on the present results. We will also study the inhibition of VEGF expression on CAM using EPMC.

The clinical implication of this study is the availability of an easily obtainable natural compound as a cancer chemopreventive agent. Ethyl *p*-methoxycinnamate is easily obtained from *Kaempferia galanga*, therefore its use may lower the expected cost of cancer treatment and increase life expectancy.

## CONCLUSIONS

Ethyl *p*-methoxycinnamate has antiangiogenic potency and its activity is dose-dependent. The antiangiogenic potency of EPMS is associated with its inhibition of tyrosine kinase.

## ACKNOWLEDGMENT

We thank the Ministry of National Education of the Republic of Indonesia, and the Rector of Airlangga University, Indonesia, for financial support in 2014. Thanks are also due to Prof. Dr. Siswandono, MS., Apt for MVD program support. 

## REFERENCES

1. Gotink KJ, Verheul HMW. Anti-angiogenic tyrosine kinase inhibitors: what is their mechanism of action? *Angiogenesis* 2010;13:1-14.
2. Filho AL, Lopes JM, Schmitt FC. Angiogenesis and breast cancer. *J Oncol* 2010. doi: 10.1155/2010/576384.
3. Kyzas PA, Stefanou D, Agnantis NJ. COX-2 expression correlates with VEGF-C and lymph node metastases in patients with head and neck squamous cell carcinoma. *Mod Pathol* 2005;18:153-60.
4. Athikomkulchai S, Vaymhasuwan P, Tunvichien S, et al. The development of sunscreen product from *Kaempferia galanga*. *J Health Res* 2007;21:253-6.
5. Kusumawati I, Yusuf H. Phospholipid complex as a carrier of *Kaempferia galanga* rhizoma extract to improve its analgesic activity. *IJPPS* 2011;3 Suppl 1:S44-6.
6. Umar MI, Asnawi MZ, Sadikun A, et al. Ethyl *p*-methoxycinnamate isolated from *Kaempferia galanga* inhibits inflammation by suppressing interleukin-1, tumor necrosis factor- $\alpha$  and angiogenesis by blocking endothelial functions. *Clinics* 2014;69:134-44.
7. Aroonrerk N, Kamkaen N. Anti-inflammatory activity of *Quercus Infectoria*, *Glycyrrhiza Uralensis*, *Kaempferia galanga* and *Coptis Chinensis*: the main components of Thai herbal remedies for aphthous ulcer. *J Health Res* 2009; 23:17-22.
8. Ekowati J, Bimo AT, Sukardiman, et al. Structure modification of ethyl *p*-methoxycinnamate and their bioassay as chemopreventive agent against mice's fibrosarcoma. *IJPPS* 2012;4 Suppl 3:S 528-32.
9. Greehough A, Smart HJM, Moore AE, et al. The COX-2/PGE<sub>2</sub> pathway: key roles in the hallmarks of cancer and adaptation to the tumour microenvironment. *Carcinog* 2009;30:377-86.
10. Yao L, Liu F, Hong L, et al. The function and mechanism of COX-2 in angiogenesis of gastric cancer cells. *J Exp Clin Cancer Res* 2011;30:13. doi:10.1186/1756-9966-30-13.
11. Roskoski R. Sunitinib: a VEGF and PDGF receptor protein kinase and angiogenesis inhibitor. *Biochem Biophys Res Commun* 2007; 356:323-8.
12. He ZH, Yue GG, Lau CB, et al. Antiangiogenic effects and mechanisms of trans-ethyl *p*-methoxycinnamate from *Kaempferia galanga* L. *J Agric Food Chem* 2012;60:11309-17.
13. Ribatti D. The chick embryo chorioallantoic membrane as in vivo assay to study antiangiogenesis. *Pharm* 2010;3:482-513.
14. Lokman NA, Elder ASF, Ricciardelli C, Oehler MK. Chick chorioallantoic membrane (CAM) assay as an in vivo model to study the effect of newly identified molecules on ovarian cancer invasion and metastasis. *Intern J Molec Sci* 2012;13:9959-70. doi: 10.3390/ijms13089959.
15. Kusumaningrum S, Budiarto E, Kosela S, et al. The molecular docking of 1,4-naphthoquinone derivatives as inhibitors of polo-like kinase 1 using Molegro Virtual Docker. *J App Sci* 2014;4: 47-53.
16. Diyah NW, Ekowati J, Siswandono. Synthesis and anti tumor activity evaluation of *N, N'*-dibenzoyl-*N, N'*-diethylurea against human breast cancer cell line (MCF-7). *IJPPS* 2014;6: 315-8.
17. Salamah N, Sugiyanto, Hartati MS, et al. Isolasi dan identifikasi eurycomanone akar pasak bumi (*Eurycoma longifolia*, Jack) serta uji anti angiogenik. *Majalah Farmasi Indonesia* 2009; 20:118-26.
18. Kumar BNP, Rajput S, Dey KK, et al. Celecoxib alleviates tamoxifen-instigated angiogenic effect

- by ROS-dependent VEGF/VEGFR2 autocrine signaling. *BMC Cancer* 2013;13:273. doi: 10.1186/1471-2407-13-273.
19. Klenke FM, Gebhard M, Ewerbeck V, et al. The selective COX-2 inhibitor Celecoxib suppresses angiogenesis and growth of secondary bone tumors: an intravital microscopy study in mice. *BMC Cancer* 2006;6:9. doi:10.1186/1471-2407-6-9.
  20. Altun A, Temiz TK, Balci E, et al. Effects of tyrosine kinase inhibitor E7080 and eNOS inhibitor L-NIO on colorectal cancer alone and in combination. *Chin J Cancer Res* 2013;25:572-84.
  21. Wang L, Chen W, Xie X, et al. Celecoxib inhibits tumor growth and angiogenesis in an orthotopic implantation tumor model of human cancer. *Exp Oncol* 2008;30:42-51.
  22. Fox SB, Generally DG, Harris AL. Review breast tumor angiogenesis. *Breast Cancer Res* 2007;9: 1-11.
  23. Zittermann SI, Issekutz AC. Basic fibroblast growth factor (bFGF, FGF-2) potentiates leukocyte recruitment to inflammation by enhancing endothelial adhesion molecule expression. *Am J Patol* 2006;168:835-46.
  24. Sahin E. Cyclooxygenase-1 in cancer and angiogenesis. *Angiology* 2009;60:242-53.
  25. Ekowati J, Tejo BA, Susilo I, et al. Microwave assisted synthesis of ethyl *p*-methoxycinnamate derivative and its activity to blockade COX-2 expression on breast cancer cell line. *Proceeding of the 3<sup>rd</sup> ICPAPS; 2013 June 18-19. Yogyakarta.* p.32-8.

## revisi perbaikan makalah

---

From: editor@univmed.org

To: j\_ekowati@yahoo.com

Date: Thursday, March 26, 2015, 09:27 AM GMT+7

---

Yth Sejawat Juni,

Maaf masih sedikitr lagi perlu diperbaiki dan lengkapi.

Salam,

Editor

On 22.03.2015 12:03, juni ekowaty wrote:

> Yth. Prof. Adi Hidayat

> editor Jurnal Universa Medisina

>

> Bersama ini kami sampaikan Perbaikan naskah Anti angionetic Juni

> revised I 7-3-2015 sesuai arahan revisi yang tertera pada huruf merah

> pada naskah yang dikirimkan kembali kepada kami.

>

> Hasil revisi tsb kami beri nama Anti angionetic Juni revised I

> 21-3-2015.

> Demikian, semoga hasil revisi kami ini bisa memenuhi persyaratan

> untuk

> diterbitkan pada jurnal Universa Medisina.

>

> Atas perhatian dan bantuan yang diberikan, kami sampaikan terima

> kasih.

>

> Hormat kami,

> Juni Ekowati

> Fakultas Farmasi Universitas Airlangga

>

> On Tuesday, March 10, 2015 12:06 PM, "[editor@univmed.org](mailto:editor@univmed.org)" wrote:

>

> Yth Sejawat Juni,

>

> Makalah sejawat telah direview, harap perbaiki dan lengkapi dalam

> huruf

> merah. Perbaikan diharapkan dalam waktu yang tidak terlalu lama ( 2

> minggu). Terima kasih.

>

> Salam,

>

> Adi

>

> On 06.03.2015 13:42, juni ekowaty wrote:

>> Yth. Prof. Adi Hidayat

>>

>> Bersama ini kami mohon informasi terkait paper kami No 002/AP/2015,

>> apakah ada revisi yang perlu kami lakukan?

>> Kami juga mohon info, jika revisi sudah dilakukan...dan dinyatakan

>> layak untuk diterbitkan...Jurnal ini kapan diterbitkan?

>>



>> Demikian, atas perhatian yang diberikan...kami sampaikan  
> terimakasih.  
>>  
>> Hormat kami,  
>> Juni Ekowati  
>>  
>> On Thursday, January 8, 2015 1:22 PM, juni ekowaty wrote:  
>  
>>  
>> Yth. Prof. Adi Hidayat  
>>  
>> Terima kasih informasinya.  
>> Kami menunggu info berikutnya.  
>>  
>> Hormat kami  
>>  
>> Juni Ekowati  
>>  
>> On Wednesday, January 7, 2015 3:55 PM, "[editor@univmed.org](mailto:editor@univmed.org) [1]"  
> wrote:  
>>  
>> YTH seJAWAT jUNI,  
>>  
>> Terima kasih atas kiriman makalahnya dan kami akan kiri ke reviewer  
>> untuk direview.  
>>  
>> Salam,  
>>  
>> Adi  
>>  
>> On 05.01.2015 09:21, juni ekowaty wrote:  
>>> Yth. Prof. Adi Hidayat  
>>> Editor-in-Chief  
>>> Universa Medicina  
>>>  
>>> Bersama ini kami kirimkan paper dengan judul : MOLECULAR DOCKING OF  
>>> STRUCTURALLY ETHYL P-METHOXYCINNAMATE AND ITS ANTI ANGIOGENIC  
>>> ACTIVITY  
>>> ON THE CHORIO ALLANTOIC MEMBRANE untuk dipublikasikan di Jurnal  
>>  
>>> Universa Medicina.  
>>>  
>>> Kami mengusulkan Prof. Dr. Siswandono, MS., Apt. dari Fakultas  
>>> Farmasi  
>>> Universitas Airlangga sebagai salah satu reviewer untuk paper ini.  
>>>  
>>> Besar harapan kami, paper ini bisa diterima untuk diterbitkan. Atas  
>>> perhatian dan bantuannya kami sampaikan terima kasih.  
>>>  
>>> Hormat kami,  
>>>  
>>> Dr. Juni Ekowati, MSi.,Apt.  
>>> Dept. Kimia Farmasi  
>>> Fakultas Farmasi  
>>>  
>>> Universitas Airlangga Surabaya  
>>> 60286  
>  
>  
>  
> Links:  
> -----

> [1] <mailto:editor@univmed.org>



Anti angionetic Juni revised 1 21-3-2015 revised 2 23-3-2015.doc  
1.1MB

## Re: Pengiriman Paper

---

From: juni ekowaty (j\_ekowati@yahoo.com)

To: editor@univmed.org

Date: Friday, March 6, 2015, 01:42 PM GMT+7

---

Yth. Prof. Adi Hidayat

Bersama ini kami mohon informasi terkait paper kami No 002/AP/2015, apakah ada revisi yang perlu kami lakukan?

Kami juga mohon info, jika revisi sudah dilakukan...dan dinyatakan layak untuk diterbitkan...Jurnal ini kapan diterbitkan?

Demikian, atas perhatian yang diberikan...kami sampaikan terimakasih.

Hormat kami,  
Juni Ekowati

On Thursday, January 8, 2015 1:22 PM, juni ekowaty <j\_ekowati@yahoo.com> wrote:

Yth. Prof. Adi Hidayat

Terima kasih informasinya.  
Kami menunggu info berikutnya.

Hormat kami

Juni Ekowati

On Wednesday, January 7, 2015 3:55 PM, "editor@univmed.org" <editor@univmed.org> wrote:

YTH seJAWAT JUNI,

Terima kasih atas kiriman makalahnya dan kami akan kiri ke reviewer untuk direview.

Salam,

Adi

On 05.01.2015 09:21, juni ekowaty wrote:

> Yth. Prof. Adi Hidayat

> Editor-in-Chief

> Universa Medicina

>

> Bersama ini kami kirimkan paper dengan judul : MOLECULAR DOCKING OF

- > STRUCTURALLY ETHYL P-METHOXYCINNAMATE AND ITS ANTI ANGIOGENIC
- > ACTIVITY
- > ON THE CHORIO ALLANTOIC MEMBRANE untuk dipublikasikan di Jurnal
  
- > Universa Medicina.
- >
- > Kami mengusulkan Prof. Dr. Siswandono, MS., Apt. dari Fakultas
- > Farmasi
- > Universitas Airlangga sebagai salah satu reviewer untuk paper ini.
- >
- > Besar harapan kami, paper ini bisa diterima untuk diterbitkan. Atas
- > perhatian dan bantuannya kami sampaikan terima kasih.
- >
- > Hormat kami,
- >
- > Dr. Juni Ekowati, MSi.,Apt.
- > Dept. Kimia Farmasi
- > Fakultas Farmasi
- >
- > Universitas Airlangga Surabaya
- > 60286

## Final Draft dan Surat bebas plagiat

---

From: editor@univmed.org

To: j\_ekowati@yahoo.com

Date: Tuesday, May 12, 2015, 11:15 AM GMT+7

---

Yth. Sdri. Juni

Mohon untuk dapat melengkapi surat pernyataan bebas plagiat dan kepengarangan (terlampir), dan mengoreksi ulang draft final makalah sejawat sebelum masuk kepercetakan. Untuk Bantuan Biaya cetak sebesar Rp. 1.000.000,- + Rp. 600.000,- (2 lembar cetak gambar berwarna) = Rp. 1.600.000,- (satu juta enam ratus ribu rupiah), mohon untuk ditransfer ke rekening

drg. Eddy Kasim  
Bank BCA a/c 3731072602  
KCP Kyai Tapa

Terima kasih.

Editor



Surat Pernyataan Bebas Plagiat.doc  
22kB



Surat Pernyataan Kepengarangan.doc  
25.5kB



Juni.pdf  
101kB

## trims Final Draft dan Surat bebas plagiat

---

From: editor@univmed.org

To: j\_ekowati@yahoo.com

Date: Monday, May 18, 2015, 09:15 AM GMT+7

---

YTh Sejawat Juni,

Terima kasih atas koreksinya dan pengiriman surat bebas plagiat serta bantuan biaya penerbitan Universa Medicina. Kami tunggu makalah sejawat yang berikutnya.

Salam,

Editor

On 17.05.2015 20:09, juni ekowaty wrote:

> Yth. Editor Jurnal Universa Medicina

>

> Bersama ini kami kirimkan kelengkapan surat pernyataan dan bukti

> transfer sesuai email terakhir.

>

> Pada draft final, masih ada revisi dari kami pada file Juni-rev, kami

> beri tanda dengan huruf merah, yaitu pada halaman 39, 43 dan 46.

>

> Demikian atas perhatiannya, kami sampaikan terimakasih.

>

> Salam

> Juni Ekowati

>

> On Tuesday, May 12, 2015 11:15 AM, "[editor@univmed.org](mailto:editor@univmed.org)" wrote:

>

> Yth. Sdri. Juni

>

> Mohon untuk dapat melengkapi surat pernyataan bebas plagiat dan

> kepengarangan (terlampir), dan mengoreksi ulang draft final makalah

> sejawat sebelum masuk kepercetakan. Untuk Bantuan Biaya cetak sebesar

> Rp. 1.000.000,- + Rp. 600.000,- (2 lembar cetak gambar berwarna) =

> Rp.

>

> 1.600.000,- (satu juta enam ratus ribu rupiah), mohon untuk

> ditransfer

>

> ke rekening

>

> drg. Eddy Kasim

> Bank BCA a/c 3731072602

> KCP Kyai Tapa

>

> Terima kasih.

>

> Editor

## Re: Final Draft dan Surat bebas plagiat

---

From: juni ekowaty (j\_ekowati@yahoo.com)

To: editor@univmed.org

Date: Sunday, May 17, 2015, 08:09 PM GMT+7

---

Yth. Editor Jurnal Universa Medicina

Bersama ini kami kirimkan kelengkapan surat pernyataan dan bukti transfer sesuai email terakhir. Pada draft final, masih ada revisi dari kami pada file Juni-rev, kami beri tanda dengan huruf merah, yaitu pada halaman 39, 43 dan 46.

Demikian atas perhatiannya, kami sampaikan terimakasih.

Salam  
Juni Ekowati

---

On Tuesday, May 12, 2015 11:15 AM, "editor@univmed.org" <editor@univmed.org> wrote:

Yth. Sdri. Juni

Mohon untuk dapat melengkapi surat pernyataan bebas plagiat dan kepengarangan (terlampir), dan mengoreksi ulang draft final makalah sejawat sebelum masuk kepercetakan. Untuk Bantuan Biaya cetak sebesar Rp. 1.000.000,- + Rp. 600.000,- (2 lembar cetak gambar berwarna) = Rp. 1.600.000,- (satu juta enam ratus ribu rupiah), mohon untuk ditransfer ke rekening

drg. Eddy Kasim  
Bank BCA a/c 3731072602  
KCP Kyai Tapa

Terima kasih.

Editor



PERNYATAAN JUNI.pdf  
31.4kB



Tranfer ke Universa medisina.pdf  
11.6kB



Juni-rev.doc



653kB



## percetakan

---

From: editor@univmed.org

To: j\_ekowati@yahoo.com

Date: Tuesday, June 23, 2015, 06:43 PM GMT+7

---

Yth Sejawat Yuni Ekowati,

Hard copy masih diproses di percetakan. Setelah selesai setiap penulis akan dikirim 2 hard copy + 4 reprint. Terima kasih.

Salam,

Editor

On 23.06.2015 14:22, juni ekowaty wrote:

> Yth. Editor Universa Medisina,

>

> Kami ingin bertanya apakah kami bisa memperoleh jurnal lengkap dalam bentuk hardcopy ?

> Hal ini kami perlukan untuk keperluan administrasi di instansi kami.

>

> Jika bisa, mohon kami diinformasikan bagaimana cara memperolehnya.

> Atas perhatian yang diberikan, kami sampaikan terima kasih.

>

> Salam,

> Juni Ekowati

>

> On Monday, May 18, 2015 9:15 AM, "[editor@univmed.org](mailto:editor@univmed.org)" wrote:

>

> YTh Sejawat Juni,

>

> Terima kasih atas koreksinya dan pengiriman surat bebas plagiat serta

> bantuan biaya penerbitan Universa Medicina. Kami tunggu makalah

> sejawat

> yang berikutnya.

> Salam,

>

> Editor

>

> On 17.05.2015 20:09, juni ekowaty wrote:

>> Yth. Editor Jurnal Universa Medicina

>>

>> Bersama ini kami kirimkan kelengkapan surat pernyataan dan bukti

>> transfer sesuai email terakhir.

>>

>> Pada draft final, masih ada revisi dari kami pada file Juni-rev,

> kami

>> beri tanda dengan huruf merah, yaitu pada halaman 39, 43 dan 46.

>>

>> Demikian atas perhatiannya, kami sampaikan terimakasih.

>>

>> Salam

>> Juni Ekowati

>>

>> On Tuesday, May 12, 2015 11:15 AM, "[editor@univmed.org](mailto:editor@univmed.org) [1]" wrote:

>>  
>> Yth. Sdri. Juni  
>>  
>> Mohon untuk dapat melengkapi surat pernyataan bebas plagiat dan  
>> kepengarangan (terlampir), dan mengoreksi ulang draft final makalah  
>> sejawat sebelum masuk kepercetakan. Untuk Bantuan Biaya cetak  
> sebesar  
>> Rp. 1.000.000,- + Rp. 600.000,- (2 lembar cetak gambar berwarna) =  
>> Rp.  
>>  
>> 1.600.000,- (satu juta enam ratus ribu rupiah), mohon untuk  
>> ditransfer  
>>  
>> ke rekening  
>>  
>> drg. Eddy Kasim  
>> Bank BCA a/c 3731072602  
>> KCP Kyai Tapa  
>>  
>> Terima kasih.  
>>  
>> Editor  
>  
>  
>  
> Links:  
> -----  
> [1] <mailto:editor@univmed.org>