

**TERAKREDITASI B (SINTA 2)  
KEMENRISTEKDIKTI SK No. 30/E/KPT/2018**

**Penerbit  
UNIVERSITAS SEBELAS MARET  
SURAKARTA INDONESIA**



(<https://drive.google.com/file/d/1YZX9GAzvwT3dj8x7dmR0ow1JL05KMZ/view?usp=sharing>)

usp=sharing)

(<https://jurnal.uns.ac.id/alchemy/about/>) / Editorial Team

1412-4092 (Print)

(<https://jurnal.uns.ac.id/alchemy/about/editorialTeam>)

(<https://drive.google.com/file/d/1zwb6Pcz1MIQ062ih9NOHK687UoQtIE8M/view?usp=sharing>)

usp=sharing)

## Editorial Team

2443-4183 (Online)

(<https://drive.google.com/uc?export=download&id=910>)

## Chief Editor

Checked By

» Dr. Teguh Endah Saraswati

turnitin

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/17/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/17/))

, (SCOPUS ID: 36931094200 h-index 4) Program Studi Kimia FMIPA Universitas Sebelas Maret Surakarta, Indonesia

Accreditation

User

Username

Password

Remember me

Login

## Journal Manager

» Dr. Khoirina Dwi Nugrahaningtyas

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/14/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/14/))

, Chemistry Department, Sebelas Maret University, Indonesia



([http://sinta2.ristekdikti.go](http://sinta2.ristekdikti.go.id=910))

id=910)

Currently, Alchemy Jurnal Penelitian Kimia has Sinta-2

accreditation based on Ministry of Research, Technology and Higher Education

Decree: 50/E/KPT/2016

(<https://drive.google.com/uc?export=download&id=910>)

(<https://jurnal.uns.ac.id/alchemy/manager/files/>)

Journal Content

Search

Search Scope

Search

## Editorial Board

» Assoc. Prof. Dr Nafarizal - Nayan

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/330409/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/330409/))

, Microelectronic and Nanotechnology- Shamsuddin Research Centre Universiti Tun Husein Onn Malaysia, Malaysia

» Prof. Dr. rer.nat Gunawan Indrayanto

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/230/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/230/))

, Faculty of Pharmacy Airlangga University Surabaya, Indonesia

» Dr Fitria Rahmawati

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/3/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/3/))


, Chemistry Department Sebelas Maret University Indonesia, Indonesia

» Ass. Prof. Dr Younki Lee

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/91659/'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/91659/))

, School of Material Science and Engineering, Gyeongsang National University, Korea, Republic of

Indexed By



(<https://scholar.google.co>)

user=43jf0hUAAA&hl=e



(<http://portal.garuda.ristekdikti.go>)



Browse

• By Issue

(<https://jurnal.uns.ac.id/alchemy/issue/archive/>)

• By Author

(<https://jurnal.uns.ac.id/alchemy/search/authors>)

• By Title

(<https://jurnal.uns.ac.id/alchemy/search/titles>)

• Other Journals

• **Categories**

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/374206'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/374206))

(<https://jurnal.uns.ac.id/index/search/categories>)  
School of Engineering and Materials Science, Queen Mary University  
of London, England, United Kingdom, United Kingdom

Citefactor  
(<http://www.citefactor.org>)  
jurnal-penelitian-  
kimia#.WoG1rNx200

Flagcounter



(<https://s09.flagcounter.com/more/yMdb/>)

» Prof. Dr Is Fatimah

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/432'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/432))

, Chemistry Department Universitas Islam Indonesia, Indonesia

» Dr Soerya Dewi Marliyana

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/1190'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/1190))

, Chemistry Department, Sebelas Maret University, Indonesia



Current Issue

ATOM 1.0

(<https://jurnal.uns.ac.id/alchemy/gateway/plugin/WebFeedGatewayPlugin/atom>)

RSS 2.0

(<https://jurnal.uns.ac.id/alchemy/gateway/plugin/WebFeedGatewayPlugin/rss2>)

RSS 1.0

(<https://jurnal.uns.ac.id/alchemy/gateway/plugin/WebFeedGatewayPlugin/rss>)

» Dr. Dani Gustaman Syarif

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/20'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/20))

, PSTNT BATAN BANDUNG, Indonesia

» Dr. Adi Darmawan

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/4752'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/4752))

, Faculty of Science & Mathematic Universitas Diponegoro  
Semarang, Indonesia

ROAD ISSN

world cat

(<http://www.worldcat.org/>)

jurnal-penelitian-

kimia/oclc/984813833)

crossref

(<https://search.crossref.org>)

q=alchemy&publication=A

» Dr. Eny Kusrini

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/228'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/228))

, Department of Chemical Engineering Faculty of Engineering  
Universitas Indonesia Depok, Indonesia

Mendeley

(<https://www.mendeley.co>)

jurnal-penelitian-kimi2/)

» Dr. Triana Kusumaningsih

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/164'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/164))

, Chemistry Department, Sebelas Maret University, Indonesia

## Journal Officer

» Heriyanto Heriyanto

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/266'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/266))

, Department of Chemistry, Sebelas Maret University, Indonesia,  
Indonesia

» Zylfi Faiz

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/112872'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/112872))

, Chemistry Department Sebelas Maret University, Indonesia

» Muhammad Zulkarnain

([javascript:openRTWindow\('https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/185532'\)](https://jurnal.uns.ac.id/alchemy/about/editorialTeamBio/185532))

, Chemistry Department Sebelas Maret University



(<https://drive.google.com/file/d/1YZX9GArwwT3dj8x7hmROxw1JL05KM2/view?usp=sharing>) (<https://jurnal.uns.ac.id/alchemy/issue/archive/>) / Vol 15, No 2 (2019)

1412-4092 (Print)

(<https://drive.google.com/file/d/1zwb6Pcz1M1CQ62ih8NOHK687UcQtlE8M/view?usp=sharing>) **Vol 15, No 2 (2019)**

2443-4183 (Online)

(<https://drive.google.com/uc?export=download&>

Checked By

**Vol 15, No 2 (2019) Alchemy**

## Jurnal Penelitian Kimia

turnitin®

User

Alchemy Jurnal Penelitian Kimia, Volume 15 No 2 Tahun 2019

Username

Password

Remember me

Login

### Full Issue

View or

download (<https://jurnal.uns.ac.id/alchemy/issue/viewIssue/2568/Dewan%20Editor>)

the full

issue

(<https://jurnal.uns.ac.id/alchemy/issue/viewIssue/2568/Kata%20Pengantar>)

(<https://jurnal.uns.ac.id/alchemy/issue/viewIssue/2568/Daftar%20Isi>)

Accreditation



(<http://sinta2.ristekdikti.id>)

Dewan Editor  
id=910)

Currently, Alchemy

Kata Pengantar

Jurnal Penelitian

Kimia has Sinta-2

Daftar Isi

accreditation based

on Ministry of

Research, Technology

and Higher Education

Decree: 30/E/KPT/201

(<https://drive.google.com>

[usp=sharing](https://drive.google.com))

Journal Content

Search

### Table of Contents

Search Scope

### Articles

Search

[Komposit Polianilina/Kitosan/Perak Nanowires](#)

[Sebagai Elektrokatalis Reaksi Evolusi Hidrogen](#)

(<https://jurnal.uns.ac.id/alchemy/ar>

[dalam Medium Netral](#)

(<https://jurnal.uns.ac.id/alchemy/article/view/30460>)

Browse

• **By Issue**

(<https://jurnal.uns.ac.id>

Fadli Fakhruallah, Purwantiningsih Sugita, Mohammad Khotib,

Takeo Akiyoshi, Shinya Takahashi

• **By Author**

(<https://jurnal.uns.ac.id/alchemy/search/authors/>

[Modifikasi Zeolit Alam Ende dengan Garam](#)

(<https://scholar.google.com>

• **By Title**

[Logam serta Potensinya Sebagai Katalis](#)

[user=43jf0hUAAAAJ&hl=](https://scholar.google.com)

(<https://jurnal.uns.ac.id/alchemy/article/view/28180>)

(<https://jurnal.uns.ac.id/alchemy/ar>)

• [Other Journals](#)

### **Hidroksimetilfurfural (HMF)**

(<https://jurnal.uns.ac.id/index>)

(<https://jurnal.uns.ac.id/alchemy/article/view/28180>)

(<http://garuda.ristekdikti>)

• [Categories](#)

(<https://jurnal.uns.ac.id/index/search/categories>)

Dessy Dwi Septian, Sri Sugiarti

Flagcounter

### **In-vivo Antipyretic Effect of Eel (*Anguilla bicolor bicolor*) Oil on Yeast-induced Fever on Mice**

(<https://s09.flagcounter.co>)

(<https://jurnal.uns.ac.id/alchemy/article/view/27086>)

(<https://jurnal.uns.ac.id/alchemy/ar>)

Current Issue

Heru Sasongko, Aulia Ayu Rahmawati, Yeni Farida, Sugiyarto  
Sugiyarto

ATOM 1.0

(<https://jurnal.uns.ac.id/al>)

RSS 2.0

(<https://jurnal.uns.ac.id/alchemy/gateway/plugin/WebFeedGatewayPlugin/rss2>)

### **Sintesis 3,4,4'-Trimetoksikalkon dan Karakterisasinya**

[jurnal-penelitian-kimia#.WoGl7rNx200](#)

RSS 1.0

(<https://jurnal.uns.ac.id/alchemy/gateway/plugin/WebFeedGatewayPlugin/rss>)

(<https://jurnal.uns.ac.id/alchemy/article/view/24256>)

(<https://jurnal.uns.ac.id/alchemy/ar>)

Rehana Rehana, Muhammad Salman Fahreza, Masita  
Wulandari S



### **Isolasi Mikroselulosa dari Limbah Eceng Gondok (*Eichornia crassipes*) dengan Metode Bleaching-Alkalisasi**

(<https://jurnal.uns.ac.id/alchemy/article/view/30862>)

(<https://jurnal.uns.ac.id/alchemy/ar>)

Jeesica Hermayanti Pratama, Rizka Lailatul Rohmah, Amalia  
Amalia, Teguh Endah Saraswati

### **Pengaruh Pelapisan Kitosan-Nisin terhadap Kualitas Ikan Sidat (*Anguilla bicolor bicolor*) selama Masa Penyimpanan pada Suhu Rendah**

(<https://jurnal.uns.ac.id/alchemy/article/view/33892>)

[jurnal-penelitian-kimia/oclc/984813833](#)

(<https://jurnal.uns.ac.id/alchemy/ar>)

Triana Kusumaningsih, Tri Martini, Tika Diah Utami

(<https://search.crossref.org?q=alchemy&publication=>)

### **The Temperature Effect on Ultrasonic-assisted of Synthesis Methyl Ferulate and Its Antiplatelet Assay**

(<https://jurnal.uns.ac.id/alchemy/article/view/29914>)

(<https://jurnal.uns.ac.id/alchemy/ar>)

Juni Ekowati, Rian Putra Pratama, Kholis Amalia Nofianti,  
Nuzul Wahyuning Diyah

### **Sintesis Komposit Hidroksiapatit-Lantanum Oksida (HA-La<sub>2</sub>O<sub>3</sub>) dengan Metode**

**Hidrotermal secara In-Situ dan Ex-Situ**

(<https://jurnal.uns.ac.id/alchemy/article/view/32062>)

(<https://jurnal.uns.ac.id/alchemy/ar>

Geofanny S Hutabarat, Dzikri T Qodir, Hendri Setiawan, Nur Akbar, Atiek Rostika Noviyanti

**Perbandingan Kandungan Senyawa Kimia dan**

**Aktivitas Antibakteri terhadap MRSA**

**(Methicillin-resistant Staphylococcus aureus)**

**Beberapa Minyak Atsiri Daun Salam (Syzygium**

**polyanthum)**

(<https://jurnal.uns.ac.id/alchemy/article/view/25736>)

(<https://jurnal.uns.ac.id/alchemy/ar>

Muhammad Salman Fareza, Esti Dyah Utami, Elesenda May Gita, Vintya Roosalinga Permatasari, Tryandika Telaumbanua, Nur Amalia Choironi

**Pengaruh Konsentrasi Asam Terhadap Sifat**

**Fisik dan Muatan Permukaan Selulosa**

**Termodifikasi**

(<https://jurnal.uns.ac.id/alchemy/article/view/33756>)

(<https://jurnal.uns.ac.id/alchemy/ar>

Agus Wedi Pratama, Bambang Piluharto, Dwi Indarti, Tanti Haryati, Hardian Susilo Addy

Designed By:



**UNS**  
UNIVERSITAS  
SEBELAS MARET



This work is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).



[View My Stats](#)

**UNS Internal Link**

- [Universitas Sebelas Maret](#)
- [LPPM UNS](#)
- [Webmail UNS](#)
- [Perpustakaan UNS](#)

Journal Profile

## ALCHEMY Jurnal Penelitian Kimia

eISSN : 2443-4183 | pISSN : 2443-4183

[Science](#) [Social](#) [Engineering](#)

[Universitas Sebelas Maret](#)



S2

Sinta Score



Indexed by GARUDA

7

H-Index

6

H5-Index

187

Citations

144

5 Year Citations



*Penerbit:*

Sebelas Maret University, Surakarta

[Website](#) | [Editor URL](#)

*Address:*

Jl. Sutami 36A, Ketingan, Surakarta

Surakarta

*Email:*

[alchemy@mipa.uns.ac.id](mailto:alchemy@mipa.uns.ac.id)

Phone:

0271 663375


Last Updated :

2020-02-21

2018

 SINTA Accreditations

Search..



Page 1 of 30 | Total Records : 293

Publications	Citation
<u>Pengaruh impregnasi logam titanium pada zeolit alam malang terhadap luas permukaan zeolit</u> LB Rianto, S Amalia, SN Khalifah Alchemy	18
<u>Pengaruh Impregnasi Logam Titanium pada Zeolit Alam Malang Terhadap Luas Permukaan Zeolit</u> LB Rianto, S Amalia, SN Khalifah Alchemy	18
<u>Pembuatan dan karakterisasi kitosan vanilin sebagai membran polimer elektrolit</u> E Pramono, SAP Pandu, P Candra, W Jati Jurnal Penelitian Kimia 8 (1), 70-78	12
<u>Pembuatan dan Karakterisasi Kitosam Vanilin Sebagai Membran Polimer Elektrolit</u> E Pramono, PSA Probowo, C Purnawan, J Wulansari ALCHEMY Jurnal Penelitian Kimia 8 (1), 70-78	11
<u>Komponen Kimia Buah Pare Belut (Trichosanthes anguina L.)</u> V Suryanti, SD Marlina, D Kristinawati Alchemy Jurnal Penelitian Kimia 4 (2)	11
<u>Pembuatan Mikrokapsul Kitosan Gel Tersambung Silang Etilen Glikol Diglisidil Eter (Psf-Egde-Cts) Sebagai Adsorben Zat Warna Procion Red MX 8b</u> T Kusumaningsih, DS Handayani, Y Lestari ALCHEMY jurnal penelitian kimia 8 (1), 47-56	10
<u>Aktivitas Antibakteri Minyak Atsiri Daun Sirih Merah</u> SD Marliyana, N Handayani, S Ngaisah, EN Setyowati ALCHEMY jurnal penelitian kimia 9 (2), 33-40	10
<u>PEMBUATAN MIKROKAPSUL KITOSAN GEL TERSAMBUNG SILANG ETILEN GLIKOL DIGLISIDIL ETER (Psf-Egde-Cts) SEBAGAI ADSORPEN ZAT WARNA Procion Red Mx 8b (PREPARATION OF MICROCAPSULE OF</u>	10





## The Temperature Effect on Ultrasonic-assisted of Synthesis Methyl Ferulate and Its Antiplatelet Assay

Juni Ekowati\*, Rian Putra Pratama, Kholis A. Nofianti, Nuzul W. Diyah

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Airlangga University, Surabaya, Indonesia 60286

\*Corresponding author

E-mail: [juni-e@ff.unair.ac.id](mailto:juni-e@ff.unair.ac.id)

DOI: 10.20961/alchemy.15.2.29914.272-286

Received 8 Mei 2019, Accepted 5 September 2019, Published 30 September 2019

### ABSTRACT

Ferulic acid (FA) has been reported to have antiplatelet activity through indirectly inhibiting P2Y<sub>12</sub> receptor. One technique to improve the activity of ferulic acid is to increase its lipophilicity. In this study the carboxylic group of ferulic acid was modified to methyl ferulate through the Fisher esterification reaction. Ultrasonic waves were utilized as source of energy emitted through water as the medium at two various of temperature, i.e., 55 °C and 65 °C. The purposes of this study are to produce methyl ferulate and to determine the reaction constant rate (*k*) and its energy activation (*E<sub>a</sub>*), at temperature of 55 °C and 65 °C. Moreover, the biological activity as antiplatelet was investigated at dose 20 mg/kg BW. The antiplatelet assay was conducted by clotting time and bleeding time methods. The results were analyzed by one way ANOVA program (*P*<0.05). The yield of methyl ferulate are 50.3% and 67.1% at 55 °C and 65 °C, respectively. The *k* value at 55°C is 4x10<sup>-5</sup> cons<sup>-1</sup> min<sup>-1</sup>, while that of at 65 °C is 9x10<sup>-5</sup> cons<sup>-1</sup> min<sup>-1</sup>. The clotting time and bleeding time of methyl ferulate obtained were 265 sec and 175 sec, respectively. The antiplatelet activity of methyl ferulate is better than ferulic acid.

**Keywords:** antiplatelet, bleeding time, clotting time, ferulic acid, Fisher esterification, methyl ferulate.

### INTRODUCTION


Excessive aggregation of platelets in blood vessels causes thromboembolism. Antithrombotic drugs, which include antiplatelet and anticoagulant therapy, prevent and treat many cases of heart abnormalities. However, some antithrombotic drugs have several pharmacological limitations which produce a variety of antithrombotic effects among patients due to metabolism, and the interactions to the others e.g., other drug interactions, environment, and genetics, or targets of the drug. The rapid knowledge of pharmacological antithrombotic drugs and understanding of the mechanism of thrombosis have triggered the

development of new antithrombotic drugs which has rapid onset of action, smaller interactions, and fewer variations in antithrombotic effects (Mega and Simon, 2015).

Ferulic acid has been reported to have antiplatelet activity by indirectly inhibiting P2Y12 receptor activity (Yang *et al.*, 2013; Zhang *et al.*, 2015). However, a major inconvenience of FA is its disfavored solubility in both oil and aqueous media constraining its application in formulations proposed for pharmaceutical products (Antonopoulou *et al.*, 2017). In an effort to increase the activity of ferulic acid, structural modification is carried out through the esterification reaction of the carboxylic group to produce an ester compound, namely methyl ferulate. The purpose of this esterification is to increase the lipophilicity of the compound, making it easier to enter the cell (Fauchier *et al.*, 2015).

Yang *et al.* (2013) reported the use of methyl ferulate, in combination as prodrug with dabigatran. The ester bond of dabigatran-methyl ferulate was expected to be hydrolyzed *in vivo* by esterase to release dabigatran and ferulic acid, which has a synergy antithrombotic effect both in vein and arterial thrombosis via two routes, namely antiplatelet and antithrombosis.

Methyl ferulate, as ester compound, can be obtained from carboxylic acid through Fischer esterification reaction (Solomons and Fryhle, 2011), methylation with dimethyl carbonate (Shieh and Repič, 2002), methylation with a mixture of  $\text{BF}_3$  and methanol (Demirbas, 2008), and methylation with dimethyl sulfate (Lamoureux and Agüero, 2009). Fischer esterification is the formation of ester compounds from a carboxylic acid by reaction with alcohol and using acid as catalyst (Solomons and Fryhle, 2011). This reaction is reversible, relatively inexpensive and easy to implement due to the use of acid catalysts and the ease in obtaining reagents. In addition, compared to the esterification reaction with a base catalyst, Fischer esterification produces fewer by-products (Pacheco *et al.*, 2014). This reaction also runs very slowly, but can immediately reach equilibrium in a few hours when carboxylic acid and alcohol are refluxed together with sulfuric acid or hydrochloric acid in small amounts.



**Figure 1.** Synthesis methyl ferulate through ultrasonic irradiation

Cravotto *et al.*, (2005) showed that Fischer esterification of cholic acid and methanol assisted by *p*-toluene sulfuric acid as a catalyst was conducted successfully in three methods, namely conventional methods (heating with reflux), microwave irradiation, and ultrasonic waves (Cravotto *et al.*, 2005). The conventional method gave a high yield, which is 93%, in five hours reaction time, while the esterification reaction using microwave and ultrasonic waves provided higher reaction yield (96% with ultrasonic waves and 97% with microwave irradiation) in shorter reaction times (30 minutes for ultrasonic waves and 3 minutes for microwave irradiation). This shows that reactions assisted by microwave and ultrasonic irradiation are more efficient, can obtain higher reaction results, and require shorter reaction times (Cravotto *et al.*, 2005).

This study will evaluate the synthesis of methyl ferulate (Figure 1), and its kinetic and order reaction using ultrasonic waves. Ultrasonic irradiation has been respected as a clean and suitable method in organic synthesis throughout the last three decades, contrasted with conventional methods; the procedure is more accessible. Through the ultrasonic irradiation method, many organic reactions can be prepared in a higher yield, shorter reaction time and milder conditions (Arani and Safari, 2011).

According to Pacheco *et al.* (2014), the effects of ultrasonic waves play a role in increasing chemical processes, especially in cases where methods require drastic conditions or long reaction times. Ultrasonic waves are emitted through the wave medium through by inducing molecular vibrational movements. Therefore, the distance between molecules can vary to be closer or further due to the movement of the oscillation of the molecule. If the ultrasonic waves in the medium are getting more intense, one point will be reached where the intra-molecular force of the fluid cannot maintain its molecular structure intact. As a result, the molecular structure of the liquid will break and form a cavity. This cavity is a denomination of a bubble cavity, and this process is called cavitation (Pacheco *et al.*, 2014).

In the present research, synthesis of methyl ferulate was conducted by reacting ferulic acid, methanol, and sulfuric acid as catalyst ( $H_2SO_4$ ) using ultrasonic wave method at 55 °C and 65 °C. This is because the temperature can influence the possibility of collisions between molecules in the reaction. The higher the temperature in the reaction, the higher the kinetic energy produced by compound molecules, thus more collisions occur between molecules. This phenomenon will increase the reaction rate. At the same time, a reaction at a higher temperature will also increase the yield percentage.

To conclude that the reaction at a temperature of 65 °C is greater than the reaction at 55 °C, the determination of the Fischer esterification rate synthesis of methyl ferulate was

evaluated. Theoretically, every 10 °C temperature increase will increase the reaction rate two to three times (Singh, 2006).

## METHODS

Ferulic acid (1) was bought from Aldrich; dimethyl sulfate, KOH, acetone, chloroform, sodium hydrogen carbonate, H<sub>2</sub>SO<sub>4</sub>, ether, and methanol were purchased from Merck. The synthesis reaction was conducted by ultrasonic Hwashin Technology Powersonic 400 series and was observed with TLC using UV lamp on  $\lambda = 254$  nm to spot detection. Melting points were evaluated by Fischer-John melting point apparatus without correction. UV spectra were achieved by Shimadzu HP 8452 UV-vis spectrophotometer. The concentration of reactant was monitored through Densitometer Shimadzu. IR spectra were presented using a Jasco FT-IR 5300 spectrophotometer. The <sup>1</sup>H NMR spectra were obtained from the JEOL JNM-ECS 400 instrument (<sup>1</sup>H NMR: 400 MHz) using appropriate DMSO-*d*<sub>6</sub> as a solvent.

### Synthesis

A mixture of 400 mg (2.0 mmol) of ferulic acid, 10.0 ml (246.1 mmol) methanol, and 2 drops (1.86 mmol) of concentrated H<sub>2</sub>SO<sub>4</sub> in a round bottom flask was shaken homogeneously and reacted using ultrasonic waves at the highest power. This reaction was conducted at temperatures of 55 °C and 65 °C. The completion of the reaction was evaluated by TLC at 0, 30, 60, 90, 180, and 360 minutes reaction time, by using ferulic acid as the reference compound and chloroform: dichloromethane: methanol = 4: 3: 0.2 as the mobile phase. After that, the rest of methanol was evaporated, and the residue was then neutralized by NaHCO<sub>3</sub> 5% solution. The desired compound was extracted by chloroform and dried by anhydrous MgSO<sub>4</sub>. The structure of the obtained compound was confirmed by UV, IR and <sup>1</sup>H-NMR spectroscopy.

### Determination of Linearity of Densitometry Method

The standard solution was arranged by taking 100  $\mu$ l, 200  $\mu$ l, 300  $\mu$ l, and 500  $\mu$ l of the stock solution of 40,000 ppm and diluted by methanol (p.a.) until 1000  $\mu$ l. Each concentration was spotted on the TLC plate and eluted by chloroform: dichloromethane: methanol (4:3:0.2). The area of the resulted spot was determined by a densitometer. The concentration in ppm (as ordinate) and spot area (as abscissa) were then plotted to form a line equation and calculated the linearity by linear equation as equation (1).

$$y = Ax + B \tag{1}$$

The samples of this test were five; the  $R_{table}$  of those samples was 0.8054. The curve is linear if  $R_{calculated}$  is greater than  $R_{table}$ .

### Determination of Reaction Order, Rate Constant ( $k$ ) and Activation Energy ( $E_a$ )

Determination of the reaction rate was carried out by the graph method based on the integral equation of the reaction rate between the concentration of ferulic acid ( $C_t$ ) vs. time ( $t$ ). The rate constants ( $k$ ) from each temperature, namely  $k_{55}$  and  $k_{65}$  were calculated after the rate order was known. The reaction rate constant ( $k$ ) of the two temperatures substituted in the Arrhenius equation as shown equation (2) to determine the energy of activation as shown in equation (3).

$$\log \frac{k_{65}}{k_{55}} = \frac{E_a}{2,303 R} \left( \frac{T_{65} - T_{55}}{T_{65} T_{55}} \right) \quad (2)$$

$$E_a = \left( \frac{2,303 R T_{65} T_{55}}{T_{65} - T_{55}} \right) \log \frac{k_{65}}{k_{55}} \quad (3)$$

Notes:

$k_{65}$  = rate constant at 65 °C ( $\text{cons}^{-1} \text{min}^{-1}$ )

$k_{55}$  = rate constant at 55 °C ( $\text{cons}^{-1} \text{min}^{-1}$ )

$E_a$  = activation energy (kcal/mol)

$R$  = gas constant (1.987 calories/mol.K)

$T_{65}$  = absolute temperature 338 K (65 °C+273)

$T_{55}$  = absolute temperature 328 K (55 °C+273)

### *In vivo* Antiplatelet Activity Evaluation

Bleeding time assay: Twelve healthy adult male white mice aged 8–12 weeks weighing 20–22 g, adapted and fed for one week. The mice were divided randomly into four groups, while each group consisting of three mice. Each mouse was given food in the form of pellets and drinks in the form of water every day in *ad libitum*. Test compounds were given orally at a dose of 20 mg/kg, given for 7 days. On the 8<sup>th</sup> day, the bleeding time test was carried out in the test animals. Bleeding time measurement was done by lying down the mice on the test table. The tip of the tail was cleaned with alcohol 70% then the tail of the mice was slashed along 2 mm. The dripping blood was placed on a filter paper every 15 seconds to observe whether there is still blood coming out. The duration, started from the first drop of the blood until the blood stops dripping, was determined as the bleeding time. The positive control was given oral aspirin dose of 20 mg/kg body weight (BW) in CMC-Na 0.5%. The negative control group was given oral CMC-Na 0.5%. Other two groups were given oral FA and MF in the same way and dose as that of aspirin. At the same time, the

clotting time was determined from the duration of the first blood drop until the clot formed. This antiplatelet assay method was approved by the Ethical Commission of Airlangga University.

### ***In silico* test**

The *in silico* test was carried out by docking the molecules against COX-1 (PDB ID 4O1Z) as the target enzyme. 4O1Z is COX-1 crystal structure binding MXM\_807 [A] as a native ligand downloaded in a resolution of 2.90 Å from the RCSB Protein Data Bank ([www.rcsb.org](http://www.rcsb.org)). The 3D structure of the molecules was imported into workspace and positioned into the cavity aligning to MXM\_807 [A]. The molecular docking was performed against the cavity-x using the MolDock SE algorithm with maximum 1500 iteration. The enzyme-ligand affinity was determined from the docking score that expressed as the Moldock Score (MDS). The best docking results must meet the criteria of the lowest energy pose, and the docked molecule was in the same place as the native ligand.

## **RESULTS AND DISCUSSION**

### **Synthesis**

Methyl ferulate (MF) was obtained as white solids, and its melting point was 58 °C. The yield obtained from 55 °C and 65 °C reactions were 50.3% and 67.1%, respectively. The  $R_f$  values are shown in Table 1 which display that  $R_f$  value of MF is higher than FA. This condition occurred because of a change in lipophilicity of compounds from a hydrophilic acid (FA) to a more lipophilic ester (MF). According to ChemBioUltra software, Log P of FA is 1.42 while log P of MF is 1.68.


**Table 1.**  $R_f$  of Methyl ferulate

Eluent	$R_f$ MF	$R_f$ FA
Ethyl acetate : <i>n</i> -hexane (1:1)	0.63	0.43
Chloroform : dichlorometane : methanol (4:3:0.2)	0.68	0.31
Acetone : chloroform (1:4)	0.74	0.51

The melting point of methyl ferulate was lower than ferulic acid, which melted at 171 °C. This phenomenon is caused by changes in the hydrogen bonds in these molecules, wherein the molecule of ferulic acid there are hydrogen bond dimers, while in the molecule methyl ferulate there is no hydrogen bond due to the loss of the hydroxy group of the carboxylate in ferulic acid which is replaced by the methoxy group in methyl ferulate.

The spectroscopic data of methyl ferulate were as follow:  $\lambda_{\max}$  (nm): 326 nm; IR ( $\nu$ ;  $\text{cm}^{-1}$ ): 1718 (C=O ester), 3406 (O-H), 1642 (-C=C- alkene), 1590 and 1436 (-C=C- aromatic), 1264 (-C-O).  $^1\text{H-NMR}$  ( $\delta$ : ppm) (400 MHz,  $\text{CDCl}_3$ ): 3.83 (s, 3H), 3.88 (s, 3H), 6.25 (1H, d,  $J = 16.0$  Hz), 6.40 (1H, s), 6.88 (1H, d,  $J = 8.4$  Hz), 7.00 (2H, d,  $J = 2.0$  Hz), 7.60 (1H, d,  $J = 16.0$  Hz).  $^{13}\text{C-NMR}$  ( $\delta$ : ppm) (100 MHz,  $\text{CDCl}_3$ ): 51.74 (1C), 55.91 (1C), 109.75 (1C), 114.72 (1C), 150.74 (1C), 123.05 (2C), 126.79 (1C), 147.13 (1C), 148.33 (1C), 168.13 (1C).

The profile of IR spectra of the product from the reactions at 65 °C and 55 °C showed that the designed ester was successfully synthesized, based on the stretching band of C=O ester at wavenumber 1718  $\text{cm}^{-1}$  that is different from C=O ferulic acid band at 1691  $\text{cm}^{-1}$ . In addition, the widening band around 3000  $\text{cm}^{-1}$  originating from the OH group in the starting material was no longer found. Comparison between the IR spectra of the product from reactions at 65 °C and 55 °C showed the correlation value of the two compounds was 0.9949. This result shows that the synthesis at both temperatures could produce the same compound of methyl ferulate (Figure 2).



**Figure 2.** The comparison of IR spectra of methyl ferulate synthesized at 55 °C and 65 °C.


Based on the  $^1\text{H-NMR}$  spectra, it was known that there were three additional proton from methoxy groups at chemical shift ( $\delta$ ) 3.84 ppm (3H, singlet). There was one proton at  $\delta_{\text{H}}$  6.24 ppm (1H, duplet) and one proton at  $\delta_{\text{H}}$  7.57 ppm (1H, duplet) which have the same coupling constant  $J=16$  Hz. This value indicated that protons in alkenes have a *trans* configuration (Pavia *et al.*, 2014).

From UV spectral data, it was found that  $\lambda_{\max}$  ferulic acid was 322 nm, whereas  $\lambda_{\max}$  of methyl ferulate was 326 nm. There was little difference of  $\lambda_{\max}$  between two compounds

because the methoxy group in methyl ferulate was not a chromophore group. Moreover, the electronic transition from  $\sigma \rightarrow \sigma^*$  in the methyl group required a large amount of energy. As a result, the shift of  $\lambda$  in the methyl ferulate was not significant (Pavia *et al.*, 2014).

All of the spectral data showed that the same compounds from both reactions at 55 °C and 65 °C were successfully synthesized. The higher the temperature, the higher heat energy that is received by the molecule, hence the greater the kinetic energy that can be used by molecules to collide each other in order to initiate a reaction. This theory makes it is easier for chemical reactions and higher probabilities to form new compounds.

The mechanism reaction for the synthesis of methyl ferulate using Fischer esterification method is nucleophilic addition (Figure 3). The reaction began with the protonation of the carbonyl group of ferulic acid by strong acid as catalyst. Protonated carbonyl groups would be attacked by nucleophilic groups from alcohol. The nucleophilic groups of carbonyl would take protons and formed water ( $H_2O$ ) that would be eliminated as a by-product. The protons in the carbonyl group were then released again to reform a strong acid catalyst (Solomons and Fryhle, 2011).



**Figure 3.** Reaction mechanism of methyl ferulate synthesis

The synthesis in the present study used the ultrasonic wave method, in which the ultrasonic apparatus firstly transmitted the wave through the water as the medium, then the wave propagated to all parts of the solution in the reaction flask. The mechanical activation process, called cavitation, removed the attraction between molecules in the liquid phase. Once small air bubbles were formed inside the cavity, these bubbles would absorb energy from ultrasonic waves and produce the larger cavity.

As the socket grew, the air bubbles inside could no longer absorb ultrasonic energy. Finally, the fluid around the cavity would go inside and broke the cavity. This rupture of the air cavity created favorable conditions for chemical reactions. As a result of the vibrational propagation, kinetic movements of molecules would occur followed by collisions that triggered chemical reactions. In order to obtain the best condition of reactions, the oriented



procedures were conducted at several reaction temperatures. In the condition including temperature at 35 °C and the moderate intensity of ultrasonic wave, the reaction ran slowly and the new compound formed in the 120<sup>th</sup> minutes, based on the new spot observed on TLC. The next reaction that was carried out at condition of 50 °C in temperature and moderate intensity of ultrasonic wave, the new compound was found after 90 min. This results indicated that the reaction proceeds faster by increasing the temperature.


In a further experiment, the temperature was increased to 65 °C, and the ultrasonic intensity was raised from moderate to high intensity then the new compound was formed after 60 min reaction. Based on these results, temperature of 65 °C and high ultrasonic intensity were selected as the condition of methyl ferulate's synthesis. As a comparison for the study of reaction rates, 55 °C was chosen as reaction temperature with the same ultrasonic intensity. Theoretically, the reaction rate was also influenced by the initial concentration of the reacting compound (Singh, 2006).

To find out whether the TLC-densitometry can be used to monitor the changes of ferulic acid concentration (reactant), a linearity test was carried out between the ferulic acid level (ppm) and the percentage area (%). Data processing result the line equation  $y = 0.001x + 3.4833$  with the correlation coefficient (R) = 0.9943. The  $R_{table}$  suitable for this experiment using 5 samples is 0.8054. The value of  $R_{calculated}$ , which is greater than  $R_{table}$  indicated that the two variables had significant linear relationship.

#### **Determination of Reaction Order**

Determination of the reaction order was done by the graphical method, where the concentration of reactants (ferulic acid) in three data form ( $C_t$ ,  $\log C_t$ ,  $1/C_t$ ) each was plotted against time (in minutes) to obtained a straight-line. Concentration of ferulic acid during esterification reaction at 55 °C and 65 °C were shown in Figure 4 and 5.

According to Sinko *et al.* (2006), reaction order can be determined by three methods, namely the graph method, the substitution method, and the half-time method. In this study, we used the graph method. The graph method was conducted by making some plots of  $C_t$ ,  $\log C_t$ , and  $1/C_t$  each against time (t) and then was compared which plot that produced a linear line. If the graph between concentration ( $C_t$ ) and time (t) was linear, then the reaction was zeroth order. If the graph between log concentration ( $\log C_t$ ) and time (t) was linear, then the reaction had first order. The reaction order was second if the graph between  $1/C_t$  and time (t) was linear (Singh, 2006). The linearity of the graph was determined by the value of R obtained from the regression equation of the line.



**Figure 4.** (a) Plot of ferulic acid concentration (ppm) ( $C_t$ ) vs time in minutes ( $t$ ) at 55 °C (b) plot of  $\log C_t$  vs time in minutes ( $t$ ) at 55 °C (c) Plot of  $1/C_t$  vs time in minutes ( $t$ ) at 55 °C.

Data in Fig 4 and Fig 5 (above) showed that the greatest  $R^2$  value was obtained from the plot of  $1/C_t$  vs. time ( $t$ ). This results showed that the Fischer esterification reaction of ferulic acid to methyl ferulate followed the second-order which has an equation 4.

$$\frac{1}{C_t} = \frac{1}{C_0} + kt \tag{4}$$


Notes :

$C_t$  = the remaining concentration of the reactant (ferulic acid at time=  $t$  minutes)

$C_0$  = initial concentration of reactant (ferulic acid at time= 0 minute)

$k$  = rate constant

$t$  = time of reaction



**Figure 5.** (a) plot of ferulic acid concentration in ppm ( $C_t$ ) vs time in minutes ( $t$ ) at 65 °C (b) plot of  $\log C_t$  vs time in minutes ( $t$ ) at 65 °C (c) Graph of Plot  $1/C_t$  vs time in minutes ( $t$ ) at of 65 °C.

The equation of second-order reaction (equation 4) means that if the ferulic acid concentration increased twice, then the rate of esterification reaction will rise up 4-fold or increase 2 times exponentially. The changes in reaction conditions such as temperature, solvent or initial compound concentrations will affect the rate constant ( $k$ ) for the reaction. Results of the rate constant ( $k$ ) for these experiments are in accordance with the mentioned theory.

#### Determination of Rate Constant ( $k$ ) and Energy of Activation ( $E_a$ )

After determining the rate order of the reaction and rate constant ( $k$ ) for each reactions using equation 4., the activation energy ( $E_a$ ) from the ferulic acid esterification reaction can also be calculated from the two  $k$  value of reactions at 55 °C and 65 °C (Figure 6).




Figure 6. (a) Plot of  $1/C_t$  ferulic acid versus time ( $t$ ) at 55 °C (b) Plot of  $1/C_t$  ferulic acid versus time ( $t$ ) at 65 °C.

The activation energy value ( $E_a$ ) can be resolved using the Arrhenius equation using data of absolute temperature ( $T$ ) (equation 2) and rate constant ( $k$ ) (equation 3). The results of calculation were shown in Table 2.

Table 2. Linear equation and rate constant ( $k$ ) of both temperatures and activation energy of reaction


Temperature (°C)	Temperature absolute (K)	Line Equation	$k$ (cons <sup>-1</sup> min <sup>-1</sup> )	Activation Energy ( $E_a$ ) (kcal/mol)
55 °C	328	$y = 4.10^{-5}x + 0.0120$	$4.10^{-5}$	17.9
65 °C	338	$y = 9.10^{-5}x + 0.0115$	$9.10^{-5}$	

The results of data processing denoted that the  $E_a$  of the Fischer esterification of ferulic acid was 17.9 kcal/mol. This means that each mole of ferulic acid requires 17.9 kcal energy to carry out an esterification reaction with the help of a catalyst, then the energy needed to react  $2.10^{-3}$  moles of the ferulic acid compound is 0.0358 kcal or equal to 35.8 calories.

### Antiplatelet Assay


The result of *in vivo* antiplatelet evaluation of methyl ferulate compared to the negative control, positive control (aspirin) and ferulic acid are shown in Figure 7. Based on the statistical analysis one way ANOVA, it was found that the blood clotting time and bleeding time of the ferulic acid and methyl ferulate respectively were significantly different from negative control ( $P= 0.000$  and  $0.002$ ) which means that ferulic acid and methyl ferulate had antiplatelet effects.

According to data in Figure 7, it appears that the change from carboxylic acid to ester group can increase activity 1.6 times more potent based on the blood clotting time, and 1.35 times higher based on bleeding time data. However, the antiplatelet potency of methyl ferulate was same with aspirin according to the capability to prolong clotting time and bleeding time.




**Figure 7.** Antiplatelet activity histogram of methyl ferulate (MF), ferulic acid (FA), aspirin (ASP), and control negative (CN). CT = clotting time, BT = bleeding time, (n=5).

The results of docking *in silico* against COX-1 enzyme (PDB ID 4O1Z) indicate that methyl ferulate has greater potency for COX-1 inhibition than ferulic acid. MolDock score of methyl ferulate is 112.223 kcal/mol, while ferulic acid has 106.779 kcal/mol. As observed in the Figure 8 and 9, methyl ferulate and ferulic acid showed the same mode of interaction which involved their same functional groups and the same amino acids, i.e. Ala527 and Met522, in cavity-4 (vol. 76.8; surface 239.36) of COX-1 ( $X=240.12$ ;  $Y=99.88$ ;  $Z=5.09$ ). The possible difference comes from the additional interaction of the hydrophobic and steric interaction of methyl ester moiety with Val349, which are invisible in Figure 9.



**Figure 8.** The interaction ferulic acid with amino acids residue of COX-1 in (a) 2D and (b) 3D.



**Figure 9.** The interaction methyl ferulate with amino acids residue of COX-1 in (a) 2D and (b) 3D.

From *in silico* study on P2Y<sub>12</sub> receptors, it was known that MolDock Score of methyl ferulate (-111,142 kcal/mol) was lower than ferulic acid (-100,981 kcal/mol) (Ekowati *et al.*, 2018) which means methyl ferulate was capable of building more strong ligand-receptor interaction than ferulic acid.

The P2Y<sub>12</sub> receptor that bound to the Gα<sub>i2</sub> protein would encourage platelet activation and aggregation through several intracellular pathways in the downstream receptors. The activation of the Gα<sub>i2</sub> protein caused inhibition of adenosine cyclic monophosphate (cAMP) which facilitated platelet activation, by inhibiting cAMP-dependent protein kinase. P2Y<sub>12</sub> also strengthened the platelet response by stimulating phosphatidylinositol-3 kinase (PI-3K) activity which leads to continuous aggregation. P2Y<sub>12</sub> activated glycoprotein (GP) GIIb/IIIa receptors which in turn bound fibrinogen and connected platelets (Yang *et al.*, 2013; O'Connor *et al.*, 2011). Therefore, if these receptors bind to the test compound then the Gα<sub>i2</sub>-protein bonding will be inhibited, and further, the activation and aggregation of platelets will be also prevented.

## CONCLUSIONS

The temperature effects on ultra sonic assisted synthesis of methyl ferulate are to affect the reaction rate constant and yield product. The value of reaction rate constant and yield product at temperature of 65 °C are higher than 55 °C. The presence of a methyl group of ester moiety increased the antiplatelet activity. The potency of methyl ferulate as antiplatelet agent was same as aspirin.

## ACKNOWLEDGMENTS

This research was funded by PTUPT Grant of Airlangga University in the year 2018.

## REFERENCES

- Antonopoulou, I., Leonov, L., Jütten, P., Cerullo, G., Faraco, V., Papadopoulou, A., Kletsas, D., Ralli, M., Rova, U. and Christakopoulos, P., 2017. Optimized Synthesis of Novel Prenyl Ferulate Performed by Feruloyl Esterases from *Myceliophthora Thermophila* in Microemulsions. *Applied Microbiology and Biotechnology*, 101(8), 3213-3226. doi: 10.1007/s00253-017-8089-8.
- Arani, N.M. and Safari, J., 2011. A Rapid and Efficient Ultrasound-Assisted Synthesis of 5, 5-diphenylhydantoins and 5,5-diphenyl-2-thiohydantoins. *Ultrasonics Sonochemistry*, 18(2), 640-643. doi: 10.1016/j.ultsonch.2010.09.001.
- Cravotto, G., Boffa, L., Turello, M., Parenti, M. and Barge, A., 2005. Chemical Modifications of Bile Acids Under High-Intensity Ultrasound or Microwave Irradiation. *Steroids*, 70(2), 77-83. doi: 10.1016/j.steroids.2004.09.007.
- Demirbas, A., 2008. Comparison of Transesterification Methods for Production of Biodiesel from Vegetable Oils and Fats. *Energy Conversion and Management*, 49(1), 125-130. doi: 10.1016/j.enconman.2007.05.002.
- Ekowati, J., Diyah, N.W., Nofianti, K.A., Hamid, I.S. and Siswandono, S., 2018. Molecular Docking of Ferulic Acid Derivatives on P2Y<sub>12</sub> Receptor and Their ADMET Prediction. *Journal of Mathematical and Fundamental Sciences*, 50(2), 203-219. doi: 10.5614/2Fj.math.fund.sci.2018.50.2.8.
- Fauchier, L., Clementy, N., Pelade, C., Collignon, C., Nicolle, E. and Lip, G.Y., 2015. Patients with Ischemic Stroke and Incident Atrial Fibrillation: A Nationwide Cohort Study. *Stroke*, 46(9), 2432-2437. doi: 10.1161/STROKEAHA.115.010270.
- Lamoureux, G. and Agüero, C., 2009. A Comparison of Several Modern Alkylating Agents. *Arkivoc*, 1, 251-264.
- Mega, J.L. and Simon, T., 2015. Pharmacology of Antithrombotic Drugs: An Assessment of Oral Antiplatelet and Anticoagulant Treatments. *The Lancet*, 386(9990), 281-291. doi: 10.1016/S0140-6736(15)60243-4.
- O'Connor, S., Montalescot, G. and Collet, J.P., 2011. The P2Y<sub>12</sub> Receptor as A Target of Antithrombotic Drugs. *Purinergic Signalling*, 7(3), 325. doi: 10.1007/s11302-011-9241-z.

- Pacheco, B.S., Nunes, C.F., Rockembach, C.T., Bertelli, P., Mesko, M.F., Roesch-Ely, M., Moura, S. and Pereira, C.M., 2014. Eco-Friendly Synthesis of Esters Under Ultrasound with p-Toluenesulfonic Acid as Catalyst. *Green Chemistry Letters and Reviews*, 7(3), 265-270. doi: 10.1080/17518253.2014.941950.
- Pavia, D.L., Lampman, G.M., Kriz, G.S. and Vyvyan, J.A., 2014. *Introduction to Spectroscopy*. Cengage Learning, Stamford.
- Shieh, W.C., Dell, S. and Repič, O., 2002. Large Scale Microwave-Accelerated Esterification of Carboxylic Acids with Dimethyl Carbonate. *Tetrahedron Letters*, 43(32), 5607-5609. doi: 10.1016/S0040-4039(02)01116-4.
- Singh, Y., 2006. *Martin's Physical Pharmacy and Pharmaceutical Sciences*. New Jersey: Department of Pharmaceutics Ernest Mario School of Pharmacy Rutgers, The State University of New Jersey.
- Solomons, T. W. G. & Fryhle, C. B., 2011. *Organic Chemistry*, 10th edition. John Wiley & Sons, Inc., New Jersey.
- Yang, X.Z., Diao, X.J., Yang, W.H., Li, F., He, G.W., Gong, G.Q. and Xu, Y.G., 2013. Design, Synthesis and Antithrombotic Evaluation of Novel Dabigatran Prodrugs Containing Methyl Ferulate. *Bioorganic & Medicinal Chemistry Letters*, 23(7), 2089-2092. doi: 10.1016/j.bmcl.2013.01.126.
- Zhang, P.X., Lin, H., Qu, C., Tang, Y.P., Li, N.G., Kai, J., Shang, G., Li, B., Zhang, L., Yan, H. and Liu, P., 2015. Design, Synthesis, and In Vitro Antiplatelet Aggregation Activities of Ferulic Acid Derivatives. *Journal of Chemistry*, 2015, 1-7. doi: <http://dx.doi.org/10.1155/2015/376527>.

---

**Proof Alchemy\_29914**

4 messages

---

**alchemy** <alchemy@mipa.uns.ac.id>  
To: juni-e@ff.unair.ac.id

Sat, Sep 28, 2019 at 2:38 PM


Ykh Author,

Berikut kami kirimkan proof artikel anda yang akan diterbitkan dalam Alchemy jurnal penelitian Kimia dalam bentuk word (mohon diaktifkan "Final showing Markup" pada tab "Review" MS Office anda untuk melihat komentar dari kami). Mohon kesediaannya untuk mengecek ulang dan memberikan revisi atas masukan dan komentar dari editor (jika ada). Bagian yang dirubah silakan ditandai (misal dengan font merah atau highlight).

Agar dapat dipublikasikan dibulan September 2019, kami menunggu respon Anda dalam hingga 30 September 2019 pukul 11.00 WIB. Terimakasih.

Salam,  
Alchemy

---

 **Proof\_29914-85981-1-CE-ED.doc**  
996K


---

**juni ekowati** <juni-e@ff.unair.ac.id>  
To: kholis amalia nofianti <kholis-a-n@ff.unair.ac.id>

Sun, Sep 29, 2019 at 5:36 AM

[Quoted text hidden]

---

 **Proof\_29914-85981-1-CE-ED.doc**  
996K

---

**juni ekowati** <juni-e@ff.unair.ac.id>  
To: alchemy <alchemy@mipa.uns.ac.id>

Mon, Sep 30, 2019 at 2:45 AM

Yth. Editor Alchemy,

Terimakasih atas masukan yang diberikan. Untuk tanggapan dan proof atas usulan tersebut kami lampirkan pada attachment. Kami sertakan juga hasil revisi kami dengan nama file yang sama Proof\_29914-85981-1-CE-ED. Perubahan kami beri tanda warna stabilo biru dan merah muda.

Kami berharap paper bisa dipublikasi pada bulan September 2019.

Demikian, mohon bantuan untuk proses selanjutnya.


Terima kasih  
Juni Ekowati

[Quoted text hidden]

---

**2 attachments**

 **Revisi file ALCHEMY.docx**  
16K

 **Proof\_29914-85981-1-CE-ED.doc**  
998K

---

**juni ekowati** <juni-e@ff.unair.ac.id>  
To: alchemy <alchemy@mipa.uns.ac.id>

Tue, May 12, 2020 at 1:47 PM



Yth. Ibu Fitria Rahmawati  
Editor in Chief ALCHEMY Jurnal Penelitian Kimia

Sehubungan dengan keperluan untuk persyaratan kenaikan pangkat, kami mohon bantuan untuk mendapatkan email korespondensi awal submission manuscript dengan judul : The temperature effect on ultrasonic-assisted of synthesis methyl ferulate and its antiplatelet assay. (*Proof\_29914-85981-1-CE-ED*) dan pemberitahuan accepted maupun published untuk paper kami tersebut.

Atas perhatian dan bantuan yang diberikan kami sampaikan terimakasih sebesar-besarnya.

Salam hormat  
Juni Ekowati

On Sat, Sep 28, 2019 at 2:38 PM alchemy <[alchemy@mipa.uns.ac.id](mailto:alchemy@mipa.uns.ac.id)> wrote:

[Quoted text hidden]

---

**Pembayaran artikel Juni Ekowati Fakultas Farmasi Unair, no 29914-71914-1-SM**

4 messages

---

**juni ekowati** <juni-e@ff.unair.ac.id>

Thu, Sep 19, 2019 at 12:27 PM

To: alchemy@mipa.uns.ac.id

Cc: alchemymipauns@gmail.com

Yth. Dewan Redaksi Alchemy

Sehubungan dengan SDH terbitnya keterangan Pay Now pd sistem online utk paper kami, maka bersama ini kami kirimkan biaya penerbitan sebesar Rp 1.000.000.

Mhn bantuan utk proses selanjutnya.

Terimaksh

Salam

Juni Ekowati

**Mandiri Online1568870469.png**

26K

---

**alchemy mipa uns** <alchemy@mipa.uns.ac.id>

Thu, Sep 19, 2019 at 1:57 PM

To: juni ekowati &lt;juni-e@ff.unair.ac.id&gt;

Dear Author,

Bukti pembayaran sudah diterima redaksi Alchemy. Berikut kami kirimkan kuitansi pembayarannya.

Terimakasih atas kepercayaannya kepada Alchemy.

Salam,

Khoirina DN

ALCHEMY: Jurnal Penelitian Kimia

[Quoted text hidden]

**Payment Receipt\_PR 33756 Juni Ekowati001.pdf**

39K

---

**juni ekowati** <juni-e@ff.unair.ac.id>

Thu, Sep 19, 2019 at 2:10 PM

To: alchemy mipa uns &lt;alchemy@mipa.uns.ac.id&gt;

Dear Bu Khoirina,

mhn info utk paper sy lihat sampai tahap editing...apakah itu berarti SDH diterima atau bgmn. Mhn arahannya.

Terimaksh

Salam

Juni Ekowati

[Quoted text hidden]

7/31/2020

Airlangga University Mail - Pembayaran artikel Juni Ekowati Fakultas Farmasi Unair, no 29914-71914-1-SM

**alchemy mipa uns** <alchemy@mipa.uns.ac.id>

Fri, Sep 27, 2019 at 9:56 AM

To: juni ekowati <juni-e@ff.unair.ac.id>

Dear Ibu Juni Ekowati

Paper Ibu dengan no 29914-71914-1-SM sudah dinyatakan diterima dan layak untuk diterbitkan di jurnal Alchemy vol 15 No 2.

Terimakasih,

Salam,

Khoirina Dwi N

Journal manager ALCHEMY: Jurnal Penelitian Kimia

[Quoted text hidden]