

# 12. Nystatin Profile on Candida Species in HIVAIDS

*by* Muhammad Yulianto Listiawan

---

**Submission date:** 21-Oct-2022 09:43AM (UTC+0800)

**Submission ID:** 1931099005

**File name:** 12.\_Nystatin\_Profile\_on\_Candida\_Species\_in\_HIVAIDS.pdf (514.37K)

**Word count:** 4031

**Character count:** 21937

1

## Nystatin Profile on *Candida* Species in HIV/AIDS Patients with Oral Candidiasis: A Phenomenology Study

Dwi Murtiastutik\*<sup>id</sup>, Cut Shelma Maharani, Rahmadewi and Muhammad Yulianto Listiawan

Department of Dermatology and Venereology, Faculty of Medicine, Universitas Airlangga, Dr. Soetomo General Hospital, Surabaya - 60285, Indonesia.

### Abstract

Oral candidiasis is the main symptom that often appears in patients with Human Immunodeficiency Virus (HIV) / Acquired Immune Syndrome (AIDS). Recent studies reported that some bacteria causing oral candidiasis are resistant to antifungal drugs. Describing nystatin profile against candida species in HIV / AIDS patients with oral candidiasis. Twenty-nine subjects were divided into 2 groups based on sex (23 male subjects and 6 female subjects). Subjects carried out tissue culture procedure and were tested for sensitivity to fluconazole and nystatin. The analysis was conducted by comparing sex and type of infecting bacteria. Statistical analysis used chi-square test, fisher, or ANOVA with 95% CI with  $p < 0.05$ . The average age of male and female subjects was  $43.15 \pm 3.67$  years and  $40.02 \pm 10.23$  years, respectively, with age range of 18-65 years. Recurrent oral candidiasis in male and female patients was 65.22% and 83.33%, respectively ( $p = 0.079$ ). Subjects were resistant to fluconazole as much as 77.50% in men and 61.54% in women ( $p = 0.823$ ). On the other hand, subjects sensitive to nystatin were 92.50% in men and 92.31% in women ( $p = 0.167$ ). Fluconazole was resistant to *Candida albicans* (68.00%) and non-*Candida albicans* (78.57%) ( $p = 0.048$ ), while nystatin was sensitive to *Candida albicans* (92.00%) and non-*Candida albicans* (92.86%) ( $p = 0.791$ ). Most subjects were resistant to fluconazole, while the majority of subjects were sensitive to nystatin.

**Keywords:** Nystatin, fluconazole, oral candidiasis, *Candida albicans*, HIV/AIDS.

\*Correspondence: [dwimurtiastutik@yahoo.co.id](mailto:dwimurtiastutik@yahoo.co.id); (+6231) 5020251

(Received: 29 July 2019; accepted: 11 October 2019)

**Abbreviation:** HIV: human immunodeficiency virus; CA=Candida albicans; NCA=Non-candida albicans; AIDS: acquired immunodeficiency syndrome; WHO: world health organization; ARV: antiretrovirals; and SDA: Sabouraud Dextrose Agar.

**Citation:** Dwi Murtiastutik, Cut Shelma Maharani, Rahmadewi and Muhammad Yulianto Listiawan, Nystatin Profile on *Candida* Species in HIV/AIDS Patients with Oral Candidiasis: A Phenomenology Study, *J Pure Appl Microbiol.*, 2019; **13**(4):2013-2019. <https://doi.org/10.22207/JPAM.13.4.12>

© The Author(s) 2019. **Open Access.** This article is distributed under the terms of the [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) which permits unrestricted use, sharing, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## INTRODUCTION

Oral candidiasis is one of the first clinical signs of AIDS found in 50% to 95% of HIV/AIDS patients. The condition is mostly caused by *Candida albicans*, which number is around 2-69.1% found in adult's oral cavity<sup>1,2</sup>. *Candida albicans* is not the only species causing candidiasis, but also other species including *Candida glabrata*, *Candida krusei*, *Candida tropicalis*, *Candida parapsilosis* and *Candida dubliniensis*. *Candida* species is a commensal microorganism found in oral mucosa. However, this species becomes a predisposition factor causing oral candidiasis<sup>3,4</sup>. Early treatment of oral candidiasis, according to WHO, includes administration of topical antifungal agents, such as nystatin, amphotericin B, miconazole, and clotrimazole. Those agents can be given in oral candidiasis case without complication<sup>2,5</sup>.

In Indonesia, nystatin is an effective and affordable choice of antifungal for oral candidiasis<sup>3</sup>. The available doses of nystatin are 100.000 U/mL, and 400.000 – 600.000 U/mL for adults for 4 times a day for 7 – 14 days<sup>6,5</sup>. In 2017, there were 261 out of 1020 patients with HIV/AIDS treated in Dr. Soetomo General Hospital, Surabaya, Indonesia, who experienced oral candidiasis. The number increased to 273 patients in 2018. From June – July 2018, there were 20 oral candidiasis patients treated with oral nystatin, but 30% of which returned to the hospital with the similar case.

Based on the above explanation, the authors conducted an in vitro test to measure nystatin resistance in oral candidiasis patients with HIV/AIDS.

## METHODS

The subjects of this research were HIV/AIDS patients treated in Dr. Soetomo General Hospital Surabaya, Indonesia. The inclusion criteria were: HIV/AIDS patients diagnosed with rapid test/3 HIV testing methods<sup>7,8</sup>, having diagnosed with oral candidiasis by clinical examination and 10-20% KOH test<sup>9,10</sup>, and male or female patients aged >18 years. This study excluded subjects taking antifungal drugs in 2 weeks before test, and no colony growth found on culture examination with candida Sabouraud Dextrose Agar (SDA). The subjects must fulfill the informed consent.

This study employed an observational descriptive design carried out from August 2018 to

February 2019. The process of culture extraction was conducted in Dr. Soetomo General Hospital, Surabaya, Indonesia, and followed by culture examination that was carried out in Surabaya Health Laboratory, Surabaya, Indonesia. There were 29 patients who were consecutively sampled for the subjects in this research (Fig. 1). We also obtained 53 *Candida* isolates. The study protocol was in accordance with ethical procedure (0231/KEPK/IV/2018).

We first examined the subject's culture<sup>9,10</sup> that was taken from oral tissue swab. The positive *Candida* was grown in SDA at 37°C for 48 hours<sup>11</sup>. We used CHROMagar *Candida* (CHROMagar *Candida*, France) as the SDA medium. The growing *Candida* specimen were identified using cornmeal agar and tween 80 that were incubated at 42-45°C<sup>12,13</sup>. We also conducted carbohydrate test to identify *Candida* species<sup>14</sup>. Furthermore, we conducted resistance test using disc diffusion method on Mueller Hinton agar with 2% glucose and methylen blue. The isolate of *Candida* species was implanted on the agar, then a paper disk containing nystatin or fluzonazole was placed on top of it. We made a 24-48-hour-observation to look for an inhibition zone around the paper disc. The diameter of inhibition zone was measured using caliper (Rosco Diagnostica, Taastrup, Denmark). We interpreted the inhibition zone diameter using CLSI<sup>15</sup>. This study used nystatin with a dose of 100,000 UI/ml (pharma chemistry Ltd, Bekasi, Indonesia) and fluconazole at a dose of 2 mg/ml (pharma chemistry Ltd, Bekasi, Indonesia)

We assessed demographic and clinical data of patients. The collected data were then processed using IBM SPSS Statistics software version 23.0 (IBM Corp., Armonk, NY, USA). The statistical analysis used chi-square, fisher, or ANOVA with 95% CI ( $p < 0.05$ ).

## RESULTS

Twenty-nine patients were divided into two groups based on their sex that consisted of 23 male subjects and 6 female subjects. The average age of male and female patients was  $43.15 \pm 3.67$  years and  $40.02 \pm 10.23$ , respectively. They were divided into several age groups, where most subjects were found in the age range of 36 – 45 years (8 subjects; 27.59%), and followed by age group of 56 – 65 years (7 subjects; 24.13%). Most

subjects had high school education background (48.27%), and unemployed (12 subjects; 41.37%) (Table 1). Most patients were Javanese (48.28%), and followed by Madurese (44.83%).

The subject's clinical condition was as follows: 27 subjects (93.10%) had lesions on the tongue, 1 subject (3.45%) in the mucous membrane, and 1 subject (3.45%) in the corner of the lips (Figure 2). Most subjects were recurrent

oral candidiasis patients (male = 65.22% and female = 83.33%) ( $p = 0.079$ ). Some subjects had a history of systematic antifungal treatment (male = 17.39% and female = 16.67%) and topical antifungal (male = 34.78% male and female = 50.00%), with  $p = 0.002$ . Most subjects used antiretroviral (ARV) as much as 86.96% in men and 10.34% in women ( $p = 0.518$ ; Table 1).

**Table 1.** Demographic and Clinic Characteristics of patient Gender

Variable	Sex		p
	Male (n=23)	Female (n=6)	
Age (mean $\pm$ SD)	43.15 $\pm$ 3.67	40.02 $\pm$ 10.23	-
Education (%)			
Not attending school	2 (8.70)	2 (33.33)	-
Junior high school	7 (30.44)	1 (16.67)	
Senior high school	11 (47.83)	3 (50.00)	
Undergraduate/Diploma	3 (13.04)	0 (0.00)	
Ethnic (%)			
Java	10 (43.48)	4 (66.67)	-
Madura	11 (47.83)	2 (33.33)	
Other	2 (8.70)	0 (0.00)	
Oral candidiasis (%)			
Recurrent	15 (65.22)	5 (83.33)	0.079
First infection	8 (34.78)	1 (16.67)	
Treatment history (%)			
Systemic antifungal	4 (17.39)	1 (16.67)	0.002*
Topical antifungal	8 (34.78)	3 (50.00)	
ARV treatment (%)	20 (86.96)	3 (50.00)	0.518

SD=standard deviation; ARV=antiretroviral; \*significant 0.05

**Table 2.** Comparison of culture results on male and female subjects

Variable	Sex		p
	Male (n=40)	Female (n=13)	
Bacterium (%)			
Candida albicans	18 (45.00)	7 (53.85)	0.035*
Non-Candida albicans	22 (55.00)	6 (46.15)	
Fluconazole (%)			
Sensitive	9 (22.50)	5 (38.46)	0.823
Resistance	31 (77.50)	8 (61.54)	
Nystatin (%)			
Sensitive	37 (92.50)	12 (92.31)	0.167
Resistance	3 (7.50)	1 (7.69)	

\*significant 0.05

**Table 3.** Comparison of fluconazole and nystatin sensitivity tests in the *Candida albicans* and non-candida albicans groups

Variable	Bacterium		p
	CA (n=25)	NCA (n=28)	
Fluconazole (%)			
Sensitive	8 (32.00)	6 (21.43)	0.048*
Resistance	17 (68.00)	22 (78.57)	
Nystatin (%)			
Sensitive	23 (92.00)	26 (92.86)	0.791
Resistance	2 (8.00)	2 (7.14)	

CA=Candida albicans; NCA=Non-candida albicans; \*significant 0.05




Fig. 1. Flowchart Diagram of Subject Sampling



Fig. 2. Clinical picture of the subject on the first day of hospital admission

Culture comparison based on the sex found that most subjects were infected with non-*Candida albicans* bacteria as much as 55.00% in male subjects, while most female subjects were infected with *Candida albicans* bacteria as much as 53.85% ( $p = 0.035$ ). Most subjects were resistant to fluconazole as much as 77.50% in men and 61.54% in women ( $p = 0.823$ ). The majority of subjects were sensitive to nystatin as much as 92.50% in men and 92.31% in women ( $p = 0.167$ ; Table 2).

Most *Candida albicans* bacteria were resistant to fluconazole (68.00%), and most non-*Candida albicans* bacteria were also resistant to



fluconazole (78.57%) ( $p = 0.048$ ). Most *Candida albicans* bacteria were sensitive to nystatin drugs (92.00%), while non-candida albicans bacteria were mostly sensitive to nystatin drugs as much as 92.86% ( $p = 0.791$ ; Table 3).

## DISCUSSION

The demographic data of this research included age, educational level and occupation. The highest age range was found in the adult group (63%). According to the data of Directorate General PPM & PL of the Ministry of Health in 2016, there were more than 50% HIV/AIDS patients were young adults and productive age groups with age group of 25-49 years old<sup>16</sup>. The finding of this study supported that adults included in productive and sexually active groups were more likely to engage in unprotected sexual behavior that was prone to HIV transmission<sup>17</sup>.

HIV/AIDS infection is a disease that has a huge social impact. Ninety percent of HIV/AIDS patients are likely to have oral cavity diseases that will have impact on the life quality, including occupational sector. This study found that 45% of the subjects were unemployed, and 9 subjects (45%) were high school graduates. HIV/AIDS patients with a low educational and socio-economic background have a bad oral health that makes them prone to oral cavity diseases<sup>18</sup>.

This study found that most patients were infected HIV due to heterosexual behavior (65%). This finding reflected the general condition of HIV/AIDS in East Java, in which the virus is mostly transmitted through heterosexual behavior (69.6%), followed by narcotics (21.9%)<sup>19</sup>. All patients in this study had white patches in their oral cavity, and being diagnosed with pseudomembranous oral candidiasis. This finding was consistent with a study conducted in India in 2013, in which pseudomembranous candidiasis was found in 72% of the subjects<sup>20</sup>. The ARV administration to HIV/AIDS patients could significantly reduce oral candidiasis. Some patients in this study previously had ARV therapy, while the new HIV/AIDS patients had not received the therapy yet<sup>21,22</sup>. Fungal infection was still found in most patients with antifungal therapy history, both systemic and topical<sup>23</sup>.

A disc diffusion method was conducted to measure sensitivity of all *Candida* species to

nystatin. The results of sensitivity test were in the form of inhibition zone diameter. The criteria of susceptibility and resistance to antifungal agents were determined according to the interpretation of inhibition zone diameter for fungi by Rosco Diagnostica Company<sup>24</sup>. This study found that neither *Candida albicans* nor *Candida non-albicans* species that resisted to nystatin. Nystatin currently becomes the primary therapy for oral candidiasis in patients with HIV/AIDS.

The resistance against nystatin was divided into two groups, namely intrinsic and extrinsic sensitivity. The extrinsic sensitivity shows a change in sensitivity pattern of *Candida* species, from sensitive to resistance against antifungal therapy. On the other hand, intrinsic sensitivity has occurred early on antifungal therapy. This study found some intrinsic sensitivities in a form of infection caused by *Candida krusei*, which occurred in 8% of subjects<sup>25</sup>.

Some literatures stated that nystatin resistance is very minimum, but the therapy has side effects and toxicity. Nystatin generally works by distracting fungal cytoplasmic membrane and interacting with ergosterol. Ergosterol is important to maintain integrity and function of the enzymes of fungal membrane. Nystatin produces holes in cell membrane that becomes a way out for potassium ion and magnesium cellular components. This causes damages in proton gradient of cell membrane that leads to fungal cell death. Nystatin has a high binding capacity to ergosterol and low binding capacity to 3 hydroxy or oxysterol, such as fecosterol and episterol that becomes an important reason for nystatin resistance<sup>26</sup>.

Although there is an increased in nystatin resistance, this remains a rare occurrence in fungal pathogenic isolates since nystatin could not be used for systemic fungal infection. Therefore, the indications are not as much as the azole group. The incident of nystatin-resistant strains may be largely not considered. Most fungal species are considered susceptible to nystatin. However, some of which intrinsically less susceptible to this antifungal agent, such as *Candida glabrata*, *Scedosporium prolificans* and *Aspergillus terreus*. Some species are also more susceptible to nystatin resistance, including *Candida lusitanae*, *Candida guilliermondii* and *Candida krusei*<sup>27,25</sup>.

Factors such as recurrent oral candidiasis and history of antifungal usage are considered to cause differences of *Candida* species. Those factors are thought to be a predisposing factor that changes the type of *Candida* into *Candida* non-albicans. This may occur in patients with recurrent oral candidiasis as they are also exposed to antifungal medication thus supports the previous hypothesis. The characteristics of an antifungal drug are also factors that play a role in a difficult-to-treat infection. Fungistatic drugs will further encourage the formation of resistance compared to fungicidal drugs. Absorption, distribution and metabolism of a drug also contribute to the overall effectiveness of treatment based on the location of an infection. Antifungal drug dosages, including quantity, frequency, administration schedule and cumulative doses, can also play a role in treating a fungal infection. The administration of antifungal medication along with other prescription can also change the effectiveness of antifungal drugs. In addition, in the course of advanced HIV/AIDS, extensive fungal colonization was also found<sup>27</sup>.

This study found a change in the spectrum causing oral candidiasis, since *Candida albicans* species were mostly found, while the number of *Candida* non-albicans was increasing.

## CONCLUSION

In the invitro test, no *Candida* isolate was found to be resistant to nystatin. Therefore, oral nystatin is still effectively used as a standard therapy for HIV/AIDS patients with oral candidiasis.

## ACKNOWLEDGEMENTS

Special thanks to Hastika Saraswati, MD who has assisted in our paper translate and Bernadya Yogatri Anjuwita Saputri, MD who helped us in the research. In addition, we would like to thank the Surabaya Health Laboratory, Indonesia, for helping us to carry out fluconazole resistance tests.

## CONFLICTS OF INTEREST

The authors declare that there is no conflict of interest.

## AUTHORS' CONTRIBUTION

DM conceived the study. R contributed in study design. DM and YL collected data. CSM

participated in data analysis and interpretation. DM drafted the manuscript. R and YL revised the manuscript. All authors read and approved the manuscript for publication.

## FUNDING

None.

## DATA AVAILABILITY

The dataset used and/or analyzed during the current study are available from corresponding author on reasonable request.

## ETHICS STATEMENT

The study protocol was in accordance with ethical procedure (0231/KEPK/IV/2018). All subject was received consent forms.

## REFERENCES

1. Patil S, Rao RS, Majumdar B, Anil S. Clinical Appearance of Oral Candida Infection and Therapeutic Strategies. *Frontiers in Microbiology*, 2015; 6: 1391-1391. <https://doi.org/10.3389/fmicb.2015.01391>
2. Goulart LS, Souza WWRd, Vieira CA, Lima JSd, Olinda RAd, Araujo Cd. Oral colonization by *Candida* species in HIV-positive patients: association and antifungal susceptibility study. *Einstein (Sao Paulo, Brazil)*, 2018; 16(3): eAO4224-eAO4224. <https://doi.org/10.1590/s1679-45082018ao4224>
3. Anwar KP, Malik A, Subhan KH, Profile of candidiasis in HIV infected patients. *Iranian journal of microbiology*, 2012; 4(4): 204-209.
4. Ribeiro Ribeiro AL, de Alencar Menezes TO, de Melo Alves-Junior S, de Menezes SAF, Marques-da-Silva SH, Rosario Vallinoto AC. Oral carriage of *Candida* species in HIV-infected patients during highly active antiretroviral therapy (HAART) in Belem, Brazil. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, 2015; 120(1): 29-33. <https://doi.org/10.1016/j.oooo.2015.03.008>
5. Lyu X, Zhao C, Yan Z-M, Hua H, Efficacy of nystatin for the treatment of oral candidiasis: a systematic review and meta-analysis. *Drug Design, Development and Therapy*, 2016; 10: 1161-1171. <https://doi.org/10.2147/DDDT.S100795>
6. Patil S, Majumdar B, Sarode SC, Sarode GS, Awan KH, Oropharyngeal Candidosis in HIV-Infected Patients-An Update. *Frontiers in Microbiology*, 2018; 9: 980-980. <https://doi.org/10.3389/fmicb.2018.00980>
7. Mungrue K, Sahadool S, Evans R, Boochay S, Ragoobar F, Maharaj K, Green S, Pennerman T, Tayopa O, Assessing the HIV rapid test in the fight against the HIV/AIDS epidemic in Trinidad. *HIV/AIDS (Auckland, NZ)*, 2013; 5: 191-198. <https://doi.org/10.2147/HIV.S30432>
8. Huang X, Liu X, Chen J, Bao Y, Hou J, Lu X, Xia W, Xia H, Song A, Liu Z, Su B, Chen H, Chen Y, Wu H, Evaluation

- of Blood-Based Antibody Rapid Testing for HIV Early Therapy: A Meta-Analysis of the Evidence. *Frontiers in Immunology*, 2018; **9**: 1458-1458. <https://doi.org/10.3389/fimmu.2018.01458>
9. Byadarahally Raju S, Rajappa S, Isolation and identification of *Candida* from the oral cavity. *ISRN dentistry*, 2011; **2011**: 487921-487921. <https://doi.org/10.5402/2011/487921>
10. Coronado-Castellote L, Jimenez-Soriano Y, Clinical and microbiological diagnosis of oral candidiasis. *Journal of Clinical and Experimental Dentistry*, 2013; **5**(5):e279-e286. <https://doi.org/10.4317/jced.51242>
11. Saigal S, Bhargava A, Mehra SK, Dakwala F, Identification of *Candida albicans* by using different culture medias and its association in potentially malignant and malignant lesions. *Contemporary Clinical Dentistry*, 2011; **2**(3): 188-193. <https://doi.org/10.4103/0976-237X.86454>
12. Nadeem SG, Hakim ST, Kazmi SU, Use of CHROMagar *Candida* for the presumptive identification of *Candida* species directly from clinical specimens in resource-limited settings. *The Libyan Journal of Medicine*, 2010; **5**: 10.3402/ljm.v3405i3400.2144. <https://doi.org/10.3402/ljm.v5i0.2144>
13. Baradkar V, Mathur M, Kumar S, Hichrom *Candida* agar for identification of *Candida* species. *Indian Journal of Pathology and Microbiology*, 2010; **53**(1): 93-95. <https://doi.org/10.4103/0377-4929.59192>
14. Kali A, Srirangaraj S, Charles P, A cost-effective carbohydrate fermentation test for yeast using microtitre plate. *Indian Journal of Medical Microbiology*, 2015; **33**(2): 293. <https://doi.org/10.4103/0255-0857.154884>
15. Sahu C, Jain V, Mishra P, Prasad KN, Clinical and laboratory standards institute versus European committee for antimicrobial susceptibility testing guidelines for interpretation of carbapenem antimicrobial susceptibility results for *Escherichia coli* in urinary tract infection (UTI). *Journal of Laboratory Physicians*, 2018; **10**(3):289-293. [https://doi.org/10.4103/JLP.JLP\\_176\\_17](https://doi.org/10.4103/JLP.JLP_176_17)
16. Lattif AA, Banerjee U, Prasad R, Biswas A, Wig N, Sharma N, Haque A, Gupta N, Baquer NZ, Mukhopadhyay G, Susceptibility pattern and molecular type of species-specific *Candida* in oropharyngeal lesions of Indian human immunodeficiency virus-positive patients. *J. Clin. Microb.*, 2004; **42**(3): 1260-1262. <https://doi.org/10.1128/JCM.42.3.1260-1262.2004>
17. Ishii H, Matano T, Development of an AIDS vaccine using Sendai virus vectors. *Vaccine*, 2015; **33**(45): 6061-6065. <https://doi.org/10.1016/j.vaccine.2015.06.114>
18. Kambu Y, Waluyo A, Kuntarti K, Umur orang dengan HIV AIDS (ODHA) berhubungan dengan tindakan pencegahan penularan HIV. *Jurnal Keperawatan Indonesia*, 2016; **19**(3): 200-207. <https://doi.org/10.7454/jki.v19i3.473>
19. Bajomo AS, Ayo-Yusuf OA, Rudolph MJ, Tsotsi NM, Impact of oral lesions among South African adults with HIV/AIDS on oral health-related quality of life. *Journal of Dental Sciences*, 2013; **8**(4): 412-417. <https://doi.org/10.1016/j.jds.2013.04.011>
20. Oktafiani D, Megasari NLA, Fitriana E, Nasronudin, Lusida MI, Soetjipto, Detection of Human Herpesvirus-8 Antigen In Hiv-Infected Patients In East Java, Indonesia. *African Journal of Infectious Diseases*, 2018; **12**(2): 43-46. <https://doi.org/10.21010/ajid.v12i2.7>
21. Fauk NK, Kustanti CY, Wulandari R, Damayani AD, Mwanri L, Societal determinants of HIV vulnerability among clients of female commercial sex workers in Indonesia. *PLoS one*, 2018; **13**(11): e0207647-e0207647. <https://doi.org/10.1371/journal.pone.0207647>
22. Prabawanti C, Dijkstra A, Riono P, Hartana G, A survey on HIV-related health-seeking behaviors among transgender individuals in Jakarta, based on the theory of planned behavior. *BMC Public Health*, 2015; **15**: 1138-1138. <https://doi.org/10.1186/s12889-015-2480-0>
23. Altman K, Vanness E, Westergaard RP, Cutaneous manifestations of human immunodeficiency virus: a clinical update. *Current Infectious Disease Reports*, 2015; **17**(3): 464-464. <https://doi.org/10.1007/s11908-015-0464-y>
24. Patel PK, Erlandsen JE, Kirkpatrick WR, Berg DK, Westbrook SD, Loudon C, Cornell JE, Thompson GR, Vallor AC, Wickes BL, Wiederhold NP, Redding SW, Patterson TF, The Changing Epidemiology of Oropharyngeal Candidiasis in Patients with HIV/AIDS in the Era of Antiretroviral Therapy. *AIDS Research and Treatment*, 2012; **2012**: 262471-262471. <https://doi.org/10.1155/2012/262471>
25. Sariguzel F, Berk E, Koc A, Sav H, Aydemir G, Evaluation of CHROMagar *Candida*, VITEK2 YST and VITEK® MS for identification of *Candida* strains isolated from blood cultures. *Infez. Med.*, 2015; **23**: 318-322
26. Mohamadi J, Motaghi M, Panahi J, Havasian MR, Delpisheh A, Azizian M, Pakzad I, Anti-fungal resistance in *Candida* isolated from oral and diaper rash candidiasis in neonates. *Bioinformatics*, 2014; **10**(11): 667-670. <https://doi.org/10.6026/97320630010667>
27. Moges B, Bitew A, Shewaamare A, Spectrum and the *In vitro* Antifungal Susceptibility Pattern of Yeast Isolates in Ethiopian HIV Patients with Oropharyngeal Candidiasis. *International Journal of Microbiology*, 2016; **2016**: 8. <https://doi.org/10.1155/2016/3037817>



## 12. Nystatin Profile on Candida Species in HIV/AIDS

---

### ORIGINALITY REPORT

---

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

10%

PUBLICATIONS

9%

STUDENT PAPERS

---

### PRIMARY SOURCES

---

1	Submitted to University of Nebraska, Lincoln Student Paper	3%
2	academica-e.unavarra.es Internet Source	2%
3	www.hindawi.com Internet Source	1%
4	Tong Yin, Maria Jaeger, Carsten Scheper, Gregorz Grodkowski, Tomasz Sakowski, Marija Klopčič, Beat Bapst, Sven König. "Multi-breed genome-wide association studies across countries for electronically recorded behavior traits in local dual-purpose cows", PLOS ONE, 2019 Publication	1%
5	www.scielo.br Internet Source	1%
6	rjptonline.org Internet Source	1%
7	Submitted to South Dakota Board of Regents Student Paper	1%

---

8	<a href="http://repository.ubaya.ac.id">repository.ubaya.ac.id</a> Internet Source	1 %
9	Wita Saraswati, Ratih Wardani, Suhatno, Pudjo Hartono, Ario Imandiri. "The Effect of Electroacupuncture Therapy on Pain, Plasma $\beta$ -Endorphin, and Quality of Life of Stage III Cervical Cancer Patients: A Randomized Control Trial", <i>Journal of Acupuncture and Meridian Studies</i> , 2021 Publication	<1 %
10	Submitted to International Health Sciences University Student Paper	<1 %
11	<a href="http://www.gutnliver.org">www.gutnliver.org</a> Internet Source	<1 %
12	Xin Lyu, Chen Zhao, Hong Hua, Zhimin Yan. "Efficacy of nystatin for the treatment of oral candidiasis: a systematic review and meta-analysis", <i>Drug Design, Development and Therapy</i> , 2016 Publication	<1 %
13	<a href="http://www.nfid.org">www.nfid.org</a> Internet Source	<1 %
14	Raphaël Gaisne, Morgane Péré, Victorio Menoyo, Maryvonne Hourmant, David Larmet-Burgeot. "Calciphylaxis epidemiology, risk factors, treatment and survival among	<1 %

# French chronic kidney disease patients: a case-control study", BMC Nephrology, 2020

Publication

---

15 [e-journal.unair.ac.id](http://e-journal.unair.ac.id) <1 %  
Internet Source

---

16 [www.dovepress.com](http://www.dovepress.com) <1 %  
Internet Source

---

17 Abirami Lakshmy Jayachandran, Radhika Katragadda, Ravinder Thyagarajan, Leela Vajravelu et al. "Oral Candidiasis among Cancer Patients Attending a Tertiary Care Hospital in Chennai, South India: An Evaluation of Clinicomycological Association and Antifungal Susceptibility Pattern", Canadian Journal of Infectious Diseases and Medical Microbiology, 2016 <1 %  
Publication

---

18 [journalofinequalitiesandapplications.springeropen.com](http://journalofinequalitiesandapplications.springeropen.com) <1 %  
Internet Source

---

19 [www.ie-ei.eu](http://www.ie-ei.eu) <1 %  
Internet Source

---

20 [www.tandfonline.com](http://www.tandfonline.com) <1 %  
Internet Source

---

21 Abbas Hosain Pour, Samira Salari, Pooya Ghasemi Nejad Almani. "Oropharyngeal candidiasis in HIV/AIDS patients and HIVfree

subjects in the Southeast of Iran", Current Medical Mycology, 2019

Publication

---

22

Kátia Torres, Sônia Lima, Suely Ueda. "Activity of the aqueous extract of Schinus terebinthifolius Raddi on strains of the Candida genus", Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics, 2016

Publication

---

<1 %

23

Maryam Erfaninejad, Ali Zarei Mahmoudabadi, Elham Maraghi, Mohammad Hashemzadeh, Mahnaz Fatahinia. "Epidemiology, prevalence, and associated factors of oral candidiasis in HIV patients from southwest Iran in post-highly active antiretroviral therapy era", Frontiers in Microbiology, 2022

Publication

---

<1 %

24

[hdl.handle.net](https://hdl.handle.net)

Internet Source

---

<1 %

25

Siqi Sheng, Yixian Ma, Yue Zou, Fangyuan Hu, Ling Chen. "PD-L1/PD-L2 deficiency alters numbers and types of microglia and transcriptomic landscape of neural retina", Research Square Platform LLC, 2022

Publication

---

<1 %

26

Zahra Rafat, Elahe Sasani, Yahya Salimi, Samaneh Hajimohammadi, Mohammad Shenagari, Davoud Roostaei. "The Prevalence, Etiological Agents, Clinical Features, Treatment, and Diagnosis of HIV-Associated Oral Candidiasis in Pediatrics Across the World: A Systematic Review and Meta-Analysis", *Frontiers in Pediatrics*, 2021

Publication

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On



# 12. Nystatin Profile on Candida Species in HIV/AIDS

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---