

Effect of Lead Acetate on the Concentration and Sperm Morphology in mice Balb/c (*Mus musculus*)

Yeti Mareta Undaryati*, Sri Agus Sudjarwo, Reny I'tishom****

*STIKES Insan Cendekia Medika **Airlangga University

Abstract

Air pollution is mostly dominated by motor vehicle exhaust gas. Lead is one of motor vehicle exhaust gas that is one of toxic heavy metal, one of which is a toxicant for sperm. This study was aimed to demonstrate effect of lead acetate on the concentration and sperm morphology in mice Mice (*Mus musculus*) Balb/c divided into 2 groups. K group was the control group, P group was given with lead acetate 0.075 g/Kg BW 1x/day orally for 35 days. Result, all data were normally distributed and homogeneous, then analyzed using One Way Anova test. Result showed significant difference in sperm concentration and normal-shaped sperm morphology. In conclusion, lead acetate can significant reduce sperm concentration and normal-shaped sperm morphology in mice Balb/c (*Mus musculus*).

Keywords: Lead acetate, sperm concentration, sperm morphology

Introduction

Air pollution in Indonesia is increasing because with increasing traffic volume growth, which reached 15% per year [1]. The largest air pollution is dominated by motor vehicle exhaust. Lead is one of the results of motor vehicle exhaust [2] that are toxic to the male reproductive system, especially in spermatozoa [3].

Several studies on the toxicity of lead has been done. Common effects seen in men include: reduced libido, abnormal spermatogenesis (reduced motility and number), chromosomal damage, infertility, abnormal prostatic function and changes in serum testosterone [4]. Animal studies, administration of lead acetate in mice decreased the number of spermatozoa in the epididymis (dosage 0:25%) and lower sperm concentration at high doses (doses 0.5%) [5], lower testosterone levels, lower levels of LH and FSH [6,7], even the administration of lead

acetate study in female rats who are breastfeeding decreased testicular weight and volume, as well as a decrease in the diameter of the seminiferous tubules and germinal epithelium in children are breast-fed male rats [8].

Research conducted in humans, men are exposed to the exhaust emissions of motor vehicles obtained an average morphology and sperm concentration is lower if dibandingkan with men who are not exposed [9,10].

This study was aimed to demonstrate effect of lead acetate on the concentration and sperm morphology in mice.

Materials and Method

Animal

Three months old mice (*Mus musculus*) Balb/c were used for this study. Mice were housed in controlled rooms temperature and all standard procedures.

Experimental design

Mice (*Mus musculus*) Balb/c were divided into 2 groups. K group was the control group, P group was given with lead acetate 0.075 g/Kg BW 1x/day orally for 35 days.

At the end of the entire study, the mice were sacrificed. Vas deferens were taken for examination of concentration and morphology of the sperm.

Sperm Concentration

Sperm concentration examination steps are 3-5 sperm density estimates in the field of view (magnification 400x), samples sucked with the leukocytes pipette and counted in a Neubauer counting chamber.

Sperm Morphology

The steps of sperm morphology examination are make a smear on a glass object, soak in methanol, safranin solution, wash immersion in phosphate buffer solution, flushing with a phosphate buffer solution, soak in a solution of crystal violet, rinse in distilled water, observe under the microscope with 1000x magnification and oil immersion, calculate sperm in normal-shaped and abnormal-shaped.

Statistical analysis

The results obtained from control and lead-induced animals are expressed as means \pm SD. Statistical analysis was use one way Anova test with p value <0.05 was considered significant.

Result

Sperm Concentration

Administration of lead acetate reduce mean of sperm concentration. The result of one way Anova test showed significant decrease ($p<0.05$) sperm concentration (table 1).

Table 1 Effect of lead acetate on sperm concentration

Variable	K group (Mean \pm SD)	P group (Mean \pm SD)
Sperm Concentration	14.45 \pm 4.6	7.77 \pm 3.58*

* $p<0.05$

Sperm Morphology

Administration of lead acetate also reduce mean of normal-shaped sperm morphology. The result of one way Anova test showed significant decrease ($p<0.05$) normal-shaped sperm morphology (table 2)

Table 2 Effect of lead acetate on sperm morphology

Variable	K group (Mean \pm SD)	P group (Mean \pm SD)
Normal-shaped Sperm Morphology	86.33 \pm 3.08	61.67 \pm 7.81**

** $p<0.01$

Discussion

Induction of lead acetate in this study led to a decrease in sperm concentration and a decrease in normal-shaped sperm morphology in lead acetate-induced group compared with the control group. The same results lead acetate administration to lower the concentration of spermatozoa of mice significantly [11]. The high concentration of lead in the cement can reduce sperm count significantly [12].

These results indicate that the induction of lead acetate also cause a decrease in normal-shaped sperm morphology significantly. Granting 0.5% lead acetate in drinking water for 6 weeks also showed a decrease in the percentage of motile spermatozoa, increase the percentage of abnormal sperm in the epididymis [13]. The same result, namely the provision of lead acetate 100 mg / KgBW increased abnormal sperm shape [14].

Lead is a toxin that is common and persistent environment that can damage cell components and alter the cell's genetic. Lead can increase the production of free radicals and lower the antioxidant reserves in response to damage [16]. Lead in the blood of approximately 95-99% is carried by red blood cells, and binds to hemoglobin. Heaps of lead in the red blood cells and soft tissue allegedly most responsible for the toxic effects to the body [16].

Lead toxicity is able to increase the production of reactive oxygen species (ROS). The mechanism of the toxicity of lead in an increase in ROS through two ways: increase the production of ROS and lower antioxidant reserves. Increased ROS will cause lipid peroxidation, protein peroxidation and DNA oxidation nucleic acids [17]. Oxidative stress with the accumulation of ROS, especially O_2^- and H_2O_2 thought to contribute to cell damage, apoptosis, and cell death [18]. Apoptosis that occurs in germ cells in the testes resulting in reduced germ cells resulting in a decrease in the concentration of sperm. As well as the resulting cell damage sperm abnormalities in cell shapes.

Lead can get into the brain resulting in decreased brain weight of mice significantly, increased residual lead in the rat brain, degenerative changes and hypertrophy of the endocrine cells of the pituitary gland, atrophy of the seminiferous tubules, decreased the number of Sertoli cells and Leydig, and a decrease in LH and testosterone in significant [19] which resulted in the disruption of the process of spermatogenesis and steroidogenesis in the testes.

Conclusion

Lead acetate can significant reduce sperm concentration and normal-shaped sperm morphology.

References

1. The World Bank Country Study, 1994. *Indonesia Environment and Development*. Washinton DC
2. Kusminingrum N, Gunawan G, 2008. *Polusi Udara Akibat Aktivitas Kendaraan Bermotor Di Jalan Perkotaan Pulau Jawa Dan Bali*. Pusat Litbang Jalan dan Jembatan
3. Sokol RZ, Wang S, Wan Y, Stanczyk FZ, Gentzschlein E, Chapin RE, 2002. Long Term, Low Dose Lead Exposure Alters the Gonadotrophine-Releasing Hormone System in the Male Rat. *Environmental Health Perspectives*, Vol. 110, no. 9
4. Flora G, Gupta D, Tiwari A, 2012. Toxicity of lead: A review with recent updates. *Interdiscip Toxicol*. Vol. 5(2): 47-58
5. Wadi SA, Ahmad G, 1999. Effects of lead on the male reproductive system in mice. *J Toxicol Environ Health A*. Apr 9;56(7):513-21
6. Hamadouche N Ait, Nesrine S, Abdelkeder A, 2013. Lead Toxicity and the Hypothalamic-Pituitary-Testicular Axis. *Not Sci Biol*. 5(1):1-6
7. Lamia H, Mounir T, Adel C, Abdelhamid K, Fadhel NM, Rachid S, 2008. Effects of oral intoxication by lead acetate on pituitary-testicular axis in the pubertal rat. *Italian Journal of Public Health (IJPH)* - Year 6, Volume 5, Number 4
8. Dorostghoal M, Dezfoolian A, Sorooshnia F, 2011. Effects of Maternal Lead Acetate Exposure during Lactation on Postnatal Development of Testis in Offspring Wistar Rats. *Iranian Journal of Basic Medical Sciences* Vol. 14, No. 2, Mar-Apr, 122-131

9. I'tishom R, Lubis HS, Sono OP, 2008. Perubahan morfologi spermatozoa pada pria terpapar emisi gas buang kendaraan bermotor. *Majalah Ilmu Faal Indonesia*, Juni Vol 7, No. 3. Hal 174-178
10. I'tishom R, Lunardhi H, Lubis HS, Sono OP, 2011. Perubahan Konsentrasi Spermatozoa Pada Pria Terpapar Emisi Gas Buang Kendaraan Bermotor Di Surabaya Tahun 2007. *Jurnal kesehatan reproduksi* 1 (2):67-72 among lead exposed men. *Toxicology and Applied Pharmacology* 228, 378–384
11. Wang L, Xun P, Zhao Y, Wang X, Qian L, Chen F, 2008. Effects of Lead Exposure on Sperm Concentrations and Testes Weight in Male Rats: A Meta-regression Analysis. *Journal of Toxicology and Environmental Health*. Part A, 71:7, 454 — 463
12. Wu HM, Lin-Tan DT, Wang ML, Huang HY, Lee CL, Wang HS, Soong YK, Lin JL, 2012. Lead level in seminal plasma may affect semen quality for men without occupational exposure to lead. *Reproductive Biology and Endocrinology*. 10:91
13. Wadi SA, Ahmad G, 1999. Effects of lead on the male reproductive system in mice. *J Toxicol Environ Health A*. Apr 9;56(7):513-21
14. Acharya UR, Rathore RM, Mishra M, 2003. Role of Vitamin C On Lead Acetate Induced Spermatogenesis In Swiss Mice. *Environmental Toxicology and Pharmacology* 13: 9-14
15. Patrick LND, 2006. Lead Toxicity Part II: The Role of Free Radical Damage and The Use of Antioxidant In The Pathology and Treatment of Lead Toxicity. *Alternative Medicine Review* Vol 11, no. 2
16. Patocka J, Cerny K, 2003. Inorganic Lead Toxicity: *Acta Medica The Seventh Scientific Conference of the Charles University Faculty of Medicine and Hospital Hospital*, Vol. 46, no. 2, pp. 65-72
17. Ercal N, Gurer-Orhan H, Aykin-Burns N, 2001. Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem*. Dec;1(6):529-39.
18. Handy DE, Lubos E, Yang Y, Galbraith JD, Kelly N, Zhang YY, Leopold JA, Loscalzo J, 2009. Glutathione Peroxidase-1 Regulates Mitochondrial Function to Modulate Redox-dependent Cellular Responses. *J Biol Chem*. May 1; 284(18): 11913–11921
19. Hamadouche N Ait, Nesrine S, Abdelkeder A, 2013. Lead Toxicity and the Hypothalamic-Pituitary-Testicular Axis. *Not Sci Biol*. 5(1):1-6