Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (Allium Sativum) Extract, Purple Sweet Potato

by Yudi Her Oktaviono

Submission date: 11-Apr-2020 04:13PM (UTC+0800)

Submission ID: 1295060053

File name: ated with Garlic Allium Sativum Extract, Purple Sweet Potato.pdf (898.19K)

Word count: 3615

Character count: 20621

Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (Allium sativum) Extract, Purple Sweet Potato (Ipomoea batatas) Extract, and Vitamin C

Yudi Her Oktaviono^{1,*}, Alisia Yuana Putri¹, Makhyan Jibril Al-Farabi^{1,2}, Yesita Rizky Firmansyah³, Ferry Sandra^{4,5}

Yudi Her Oktaviono^{1,*}, Alisia Yuana Putri¹, Makhyan Jibril Al-Farabi^{1,2}, Yesita Rizky Firmansyah³, Ferry Sandra^{4,5}

³Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Prof Moestopo Street 6-8, Surabaya, INDONESIA.

²School of Management, Healthcare Entrepreneurship Division, University College London, Gower St, Bloomsbury, WC1E6BT London, UK.

³Postgraduate School of Biomedicine, Faculty of Medicine, Brawijaya University, INDONESIA.

⁴Prodia Stem Cell Laboratory, Kramat 7 No. 11 Street, Jakarta, INDONESIA. ⁵Department of Biochemistry and Molecular

Department of Biochemistry and Molecula Biology, Faculty of Dentistry, Universitas Trisakti, Yyai Tapa Street No.260, Jakarta, INDONESIA

Correspondence

Yudi Her Oktaviono

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Prof Moestopo Street 6-8, Surabaya, INDONESIA.

E-mail: yoktaviono@gmail.com

History

- Submission Date: 18-1-2020;
- Review completed: 10-02-2020;
- Accepted Date: 26-02-2020.

DOI: 10.5530/pj.2020.12.

Article Available online

http://www.phcogj.com/v12/i3

Copyright

© 2020 Phcogj.Com. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.



The endothelial progenitor cell (EPCs) proliferation capability is reduced in the patient with stable coronary artery disease (SCAD). Garlic (Allium sativum), purple sweet potato (Ipomoea batatas), and vitamin C are proven antioxidant which potentially improve EPCs proliferation ability. **Objective**: To investigate the effect of garlic (Allium sativum), purple sweet potato (Ipomoea batatas), and vitamin C in EPCs proliferation from CAD patients and identify the involvement of Extracellular-Signal Regulated Kinase (ERK) Signalling Pathway.

Material and Method: Mononuclear cells were isolated from SCAD patients and cultivated with colony-forming unit (CFU)-Hill medium and divided into untreated (control), garlic extract (10 mcg/ml and 100 mcg/ml), purple sweet potato extract (1 mcg/ml and 25 μg/ml), and vitamin C (10 μg/ml and 250 μg/ml). EPCs proliferation was measured using the MTT Assay. **Results**: This research shows that EPCs proliferation was increased in the treatment with garlic extract at 10 mcg/ml and 100 mcg/ml dose (0.267 ± 0,003 and 0.391 ± 0.008 ; p < 0.05), purple sweet potato extract at 1 mcg/ml and 25 μg/ml dose (0.250 ± 0.005 and 0.3562 ± 0.023 ; p < 0.001), and vitamin C at 10 μg/ml and 250 μg/ml dose (0.259 ± 0.016 and 0.306 ± 0.022 ; p < 0.001). Increased ERK expression was found in the treatment with garlic extract, purple sweet potato extract and vitamin C. **Conclusion**: Garlic extract, purple sweet potato extract, and vitamin C can increase EPC proliferation through the ERK signaling pathway. **Key words**: Antioxidant, ERK, Endothelial Progenitor, Proliferation.

INTRODUCTION

Coronary artery disease is a major health problem that causes mortality and reduction of life quality worldwide.¹ Endothelial progenitor cells (EPCs) from the patients with stable coronary artery disease (SCAD) had a progressive reduction of proliferation abilities, which worsen as the disease progressed¹. Impaired EPCs proliferation will reduce vascular damage repair.² Lower EPCs proliferation in the patient with SCAD is also associated with a higher incidence of cardiovascular events, mortality, and morbidity.³

Multiple pathways were suggested to be responsible for EPCs impairment in SCAD patients. It is suggested that oxidative stress play significant roles in EPCs impairment through intracellular damage and balance disruption which will alter the control of apoptosis, proliferation, self-renewal, senescence, and differentiation of EPCs.⁴⁻⁶ Oxidative stress may disrupt the ERK signaling pathway, which is important in EPCs proliferation.²⁷ the antioxidant is suggested to have a beneficial effect on impaired EPCs proliferation from patients with cardiovascular disease.^{7,8}

Plants with antioxidant properties such as Chokeberry (Aronia melanocarpa),9 potato shoot (Solanum tuberosum) and Marigold (Calendula officinalis) has been proven to improve impaired EPCs proliferation.¹⁰ Similarly, vitamin C also able to prevent lowering EPCs proliferation caused by TNF-a.10 Garlic (Allium sativum), purple sweet potato (Ipomoea batatas) extract and vitamin C have potent antioxidant capabilities.8,11 Previous studies also have shown purple sweet potato extract and vitamin C on the EPCs from SCAD patients and identify its mechanism.8 Hence, this research aims to identify the effect of garlic extract, purple sweet potato extract and vitamin C treatment on the EPCs and identify the involvement of ERK phosphorylation in the SCAD patient.

MATERIAL AND METHODS

Garlic extract, purple sweet potato extract, and vitamin C preparation

Garlic and Purple sweet potato were obtained from UPT Materia Medica Batu, Indonesia and vitamin C powder was obtained from Sigma-Aldrich, USA. Purple sweet potato and garlic extract were produced with aqueous extraction method as described



Cite this article: Oktaviono YH, Putri AY, Al-Farabi MJ, Firmansyah YR, Sandra F. Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (*Allium sativum*) Extract, Purple Sweet Potato (*Ipomoea batatas*) Extract, and Vitamin C. Pharmacog J. 2020;12(3):

previously.¹² The L-ascorbic acid dose referred to previous research that use dose 250 mcg/mL to improve adipocyte stem cell proliferation.¹³ Briefly, PSP chunks were mixed in water with a 1:1 ratio and blended. The mixture was filtered then boiled for 30 min and dried up using a rotary evaporator. PSP extract was diluted with the culture medium to achieve a concentration of 1 mcg/mL and 25 mcg/mL. Vitamin C powder was suspended in double-distilled water and diluted with culture medium to obtain a concentration of 10 mcg/mL and 250 mcg/ml.

Subject recruitment and sample collection

The blood sample was obtained from eight patients with SCAD in Dr. Soetomo General Hospital with inclusion criteria as follows: male, aged 40-59, stable angina, and coronary angiography showed >50% stenosis of left main coronary artery or >70% of other coronary arteries. Subjects with a history of percutaneous coronary intervention, coronary artery bypass grafting, acute myocardial infarct, diabetes, smoking, and anemia were excluded. The study protocol was approved by the Health Research Ethics Committee of Dr. Soetomo General Hospital, Surabaya (No.292/Panke.KKE/IV/2016). Each subject has signed informed consent before subject recruitment.

EPCs isolation and culture

Peripheral blood mononuclear cells (PBMCs) were isolated from the blood sample by Ficoll Histopaque 1077 (Sigma-Aldrich, USA). To isolate EPCs from PBMCs, a standard protocol was conducted as described previously.² Briefly, 5x10⁵ cells/mL PBMCs were cultured in the fibronectin-coated 6-well plate with basal stemline II hematopoietic stem cell expansion medium (Sigma-Aldrich, USA) supplemented with 15% fetal bovine serum and 40 ng/mL vascular endothelial growth factor. The culture was maintained at 37°C with 5% CO₂ in a humidified atmosphere. Two days after, non-adherent cells were discarded, and fresh medium was added. Two weeks after, cultured cells were stained FITC-labeled anti-human CD34 antibody clone 581 (Biolegend, USA) and documented with an inverted immunofluorescence microscope. EPCs were confirmed from CD 34 expression.

EPCs proliferation assay

MTT cell proliferation assay kit (Sigma-Aldrich, USA) was used to measure EPCs proliferation as described previously. Treated EPCs were added with MTT reagent and incubated in a 37°C incubator with 5% CO₂ for 4 hours. Proliferation was determined from the reduction of tetrazolium (MTT) into insoluble formazan product by viable EPCs mitochondria. Absorbance was measured with a microplate reader at 595 nm wavelength.

Western blot analysis

The protein samples from EPCs were loaded into sodium dodecyl sulfate-polyacrylamide gel (SDS-PAGE) for electrophoresis. The electroblotting apparatus was used to transfer the protein onto the nitrocellulose membrane. The membranes were incubated with primary antibodies rabbit anti-human ERK (Sigma-Aldrich, USA) for overnight. The membrane then washed and incubated with conjugated secondary anti-rabbit IgG antibody. The result was visualized with an image analyzer.

Statistical analysis

Statistical analyses were carried out using IBM SPSS Statistics 25.0 (IBM Corp, USA), Data were compared using the ANOVA test and considered to be significantly different if p < 0.05 or p < 0.001.

RESULTS

Demography of the subjects

The blood samples were obtained from eight patients with SCAD and history of antihypertension and statin treatment. Demography of the subjects is shown in Table 1.

EPCs confirmation

EPCs were confirmed by CD34 expression, the most commonly-used surface marker to identify EPCs (Benjamin et al. 2013). EPCs colony and CD34 expression can be seen in Figure 1.

The effect of garlic (*Allium sativum*) extract, purple sweet potato (*Ipomoea batatas L.*), and vitamin C on the EPC proliferation

As shown in Figure 2, under the treatment with garlic extract, purple sweet potato and vitamin C at all doses, the proliferation capability of EPCs was significantly higher compared to the negative control (p < 0.05, ANOVA). EPCs proliferation capability was increased in a dose-dependent manner with the treatment using garlic extract, purple sweet potato extract and vitamin C. Highest EPCs proliferation capability were observed on the EPCs treated with 100 mg/ml garlic extract.

ERK expression on the EPCs treated with garlic extract, purple sweet potato extract, and vitamin C

Western blot examination showed positive expression of ERK phosphorylation at the higher dose of garlic extract, purple sweet potato extract and vitamin C (Figure 3). This suggests that a higher dose of garlic extract, purple sweet potato extract, and vitamin C will increase the EPCs proliferation capability followed by increased ERK phosphorylation.

DISCUSSION

Patient with SCAD has impaired EPCs proliferation and migration capabilities. ^{1,2} This research showed that garlic extract, purple sweet potato extract, and vitamin C had beneficial effects to improve the proliferation of the EPCs derived from peripheral blood of SCAD patients in a concentration-dependent manner. This finding is consistent with previous researches, which showed the ability of garlic extract, purple sweet extract benefit to improve the proliferation of several stem cell lines. Garlic extract was proven to improve neovasculogenesis in a dose-dependent manner for an animal model of neovacsculogenesis. ¹⁴ Garlic extract contains a high level of allicin, which has been proven to

Table 1: Subjects demography.

Variable	Mean± SD
Age (year)	54.5 ± 4.31
Height (cm)	168.0 ± 1.3
Body Mass Index (kg/m²)	25.39 ± 2.13
Systolic Blood Pressure (mmHg)	137.5 ± 24.35
Diastolic Blood Pressure (mmHg)	80.0 ± 7.56
Heart rate (beats/min)	86 ± 8.68
Total Cholesterol (mg/dL)	200.5 ± 74.75
Triglyceride (mg/dL)	97 ± 11.64
LDL (mg/dL)	145 ± 61.11
HDL (mg/dL)	35 ±7.64
LVEF (%)	53.5 ± 4.11

LDL: Low-Density Lipoprotein, HDL: High-Density Lipoprotein, LVEF: Left Ventricle Ejection Fraction.




Figure 1: a) EPCs colony under 100x magnification in the inverted microscope; b) Immunofluorescence view of FITC-labelled CD34 expression which confirms EPCs; White bar represents 100 μm.




Figure 2: Garlic extract, purple sweet potato extract, and vitamin C improve EPCs proliferation in a dose-dependent manner. EPCs proliferation after treated with a) 10 mcg/mL and 100 mcg/mL garlic extract; b) 1 mcg/mL and 25 mcg/mL purple sweet potato extract; c) 10 mcg/mL and 250 mcg/mL Vitamin C for 48 h. EPCs proliferation was measured using MTT proliferation assay and statistically analyzed, as described in Materials and Methods. Sextuplicate was performed for each group. * significant difference at p < 0.05; **: significant difference at p < 0.001.



Figure 3: Garlic extract, purple sweet potato extract, and vitamin C increase ERK expression with higher doses. ERK phosporylation in the electrophoresis gel after treated with a) 10 mcg/mL garlic extract; b) 100 mcg/mL garlic extract; c) 1 mcg/mL purple sweet potato extract; d) 25 mcg/mL purple sweet potato extract; e) 10 mcg/mL vitamin C; and f) 250 mcg/mL vitamin C for 48 h.

improve EPCs proliferation and migration.¹⁵ It is also suggested that garlic extract improves EPCs' neovasculogenesis capability through modulation of the PI3/Akt pathway.¹⁴

Purple sweet potato contains a high amount of anthocyanins, which was proven to improve impaired EPCs proliferation and migration in vivo¹⁶. High-level of anthocyanin also has been proven to increase EPCs proliferation capability in a dose-dependent manner through the reduction of intracellular ROS.⁹ This suggested that the benefit of purple sweet potato extract on the EPCs proliferation might involve the

ROS pathway. Interestingly, purple sweet potato showed an inhibitory effect on breast cancer, gastric cancer, bladder cancer cell, and colon adenocarcinoma proliferation1. 17-19 This suggests that purple sweet potato might both stimulate or inhibit cell proliferation depending on the type of cells.

Vitamin C treatment has been proven to improve the proliferation of adipocyte stem cells, cardiac progenitor cells, and intestinal stem cells. ^{13,20,21} Similar to this research, vitamin C at the dose of 10 mcg/mL was shown to prevent the impairment of EPCs proliferation

caused by TNF- α through P38 inhibition.¹⁰ Interestingly, vitamin C at the dose of 100 mg/dl can reduce the proliferation of the EPCs and vasculogenesis.²² This suggests that dose-dependent effect of vitamin C to the EPCs proliferation.

In this research, the higher dose of garlic extract (100 mcg/mL), purple sweet potato extract (25 mcg/mL), and vitamin C (100 mcg/mL) were proven to increase phosphorylated ERK of the EPCs compared to the lower dose. While the exact mechanism of impaired proliferation in the EPCs in CHD patients remains unclear, it is suggested that oxidative stress may be involved in the EPCs impairment.5 Several antioxidants such as vitamin E, resveratrol, and L-arginine have proven to improve EPCs proliferation in-vivo and in vitro. 10,16 Oxidative stress downregulates ERK signaling pathway in the embryonic stem cell.6 While ERK signal transduction pathway is responsible for promoting cell proliferation, nutrient uptake, and cell survival.23 Reduced ERK signaling pathway also has proven to reduce EPCs proliferation in patients with stable angina.2 Hence, It is speculated that the beneficial effect of the garlic extract, purple sweet extract, and vitamin C extract to improve EPCs proliferation may involve its antioxidant capability which increases ERK phosphorylation.

CONCLUSION

Treatment with garlic extract, purple sweet potato extract and vitamin C increase EPC proliferation which might involve ERK signaling pathway.

ACKNOWLEDGMENT

The authors are grateful to the staff of the Surabaya Regenerative and <u>Stem</u> Cell Centre who assist the laboratory works.

CONFLICTS OF INTEREST


The authors declare no conflict of interest.

REFERENCES

- Vasa M, Ficht L, Scherer S, Aicher A, Adler K, Urbich C, et al. Number and migratory activity of circulating endothelial progenitor cells inversely correlate with risk factors for coronary artery disease. Circ Res. 2001:89:E1-E7.
- Oktaviono YH, Sargowo D, Widodo MA, Dirgantara Y, Chouw A, Sandra F. Proliferation of Peripheral Blood-derived Endothelial Progenitor Cells from Stable Angina Subjects. Indones Biomed J. 2014;6:91-6.
- Werner N, Wassmann S, Ahlers P, Schiegl T, Kosiol S, Link A, Walenta K, Nickenig G. Endothelial progenitor cells correlate with endothelial function in patients with coronary artery disease. Basic Res Cardiol. 2007:102(6):565-71.
- Rehman J, Li J, Orschell CM, March KL. Peripheral blood endothelial progenitor cells are derived from monocyte/macrophages and secrete angiogenic growth factors. Circulation. 2003;107:1164-9.

- Loomans CJ, De Koning EJ, Staal FJ, Rabelink TJ, Zonneveld AJ. Endothelial progenitor cell dysfunction in type 1 diabetes: another consequence of oxidative stress? Antioxid Redox Signal. 2005;7(11-12):1468-75.
- Cieślar PA, Yue J, Lee HC, Skonieczna M, Wei YH. ROS and Oxidative Stress in Stem Cells. Oxid Med Cell Longev. 2017:5047168.
- Lucchesi D, Russo R, Gabriele M, Longo V, Del Prato S, Penno G, et al. Grain and bean lysates improve function of endothelial progenitor cells from human peripheral blood: involvement of the endogenous antioxidant defenses. PLoS One. 2014:9(10).
- Bong JK, Sung SH, Ji HH, Soo NP. Antioxidant Activities of *Ipomoea batatas* L. (Purple Sweet Potato) Extracts Cultured in Korea. Journal of the Society of Cosmetic Scientists of Korea. 2014;40:423-30.
- Parzonko A, Oświt A, Bazylko A, Naruszewicz M. Anthocyans-rich Aronia melanocarpa extract possesses ability to protect endothelial progenitor cells against angiotensin II induced dysfunction. Phytomedicine. 2015;15;22(14):1238-46.
- Fiorito C, Rienzo M, Crimi E, Rossiello R, Balestrieri ML, Casamassimi A, et al. Antioxidants increase number of progenitor endothelial cells through multiple gene expression pathways. Free Radic Res. 2008:42 (8):754-62.
- Hao C, Gangliang H. Extraction, characterisation and antioxidant activity of Allium sativum polysaccharide. International Journal of Biological Macromolecules. 2018:114:415-9.
- Jawi I, Indrayani W, Arijana I, Subawa A, Suprapta D. Aqueous extratc of purole sweet potato increased SOD-2 and SOD-3 expression on human umbilical vein endothelial cells in vitro. J Biol Agric Health. 2016;6(2):103-10.
- 13. Yu J, Tu YK, Tang YB, Cheng NC. Stemness and transdifferentiation of adipose-derived stem cells using L-ascorbic acid 2-phosphate-induced cell sheet formation. Biomaterials. 2014;35(11):3516-26.
- Chiang EP, Chiu SC, Pai MH, Wang YC, Wang FY, Kuo YH, Tang FY. Organosulfur garlic compounds induce neovasculogenesis in human endothelial progenitor cells through a modulation of MicroRNA 221 and the PI3-K/Akt signaling pathways. J Agric Food Chem. 2013;22;61(20):4839-49.
- Zhang FX, He CYL, Cheng GL, Wang SG. Effect of allicin on proliferation activity and migration ability of human peripheral blood endothelial progenitor cells. Chinese Journal of Biologicals. 2014;27(2):202-6.
- 16. Balestrieri M., Schiano C, Felice F, Casamassimi A, Balestrieri A, Milone L, et al. Effect of low doses of red wine and pure resveratrol on circulating endothelial progenitor cells. J Biochem. 2008;143:179-86.
- Sugata M, Lin CY, Shih YC. Anti-inflammatory and anticancer activities of Taiwanese purple-fleshed sweet potatoes (*Ipomoea batatas*) extracts. Biomed Res Int. 2015;768093.
- Xiaoling L, Min S, Lei H, Tao W, Huanjiao Z, Chao W. Purple Sweet Potato Anthocyanin Inhibits the Proliferation of Human Retinal Pigment Epithelial Cell by Blocking Cell Cycle and Inducing Apoptosis. Advance Journal of Food Science and Technology. 2016:11(8):561-9.
- Li WL, Yu HY, Zhang XJ, Ke M, Hong T. Purple sweet potato anthocyanin exerts antitumor effect in bladder cancer. Oncol Rep. 2018:40(1):73-82.
- Cao N, Liu Z, Chen Z, Wang J, Chen T, Zhao X, et al. Ascorbic acid enhances the cardiac differentiation of induced pluripotent stem cells through promoting the proliferation of cardiac progenitor cells. Cell Research. 2012;22(1):219-36.
- Mohamed MS, Chen Y, Yao CL. A serum-free medium developed for in vitro expansion of murine intestinal stem cells. Biotechnology Journal. 2014;9(7):962-70.
- Mikirova NA, Ichim TE, Riordan NH. Anti-angiogenic effect of high doses of ascorbic acid. J Transl Med. 2005:6:50.
- 23. Cantley LC. The phosphoinositide 3-kinase pathway. Science. 2002:296:1655-7.

GRAPHICAL ABSTRACT



ABOUT AUTHORS



Yudi Her Oktaviono work as an interventional cardiologist in the Department of Cardiology and Vascular Medicine Soetomo General Hospital and Faculty of Medicine, University of Airlangga. Previously, He led the Indonesian Cardiologist association in Surabaya and currently he is the vice president of the Indonesian Cardiologist association. His research interests include complex interventional cardiology, endothelial progenitor stem cells, usage of natural compound for cardiac disease and detection of early cardiac markers.



Alisia Yuana Putri work as a cardiologist in the Department of Cardiology and Vascular Medicine Soetomo General Hospital and Faculty of Medicine, University of Airlangga and Manyar Medical Center. Her research interests endothelial progenitor stem cells and cardiology risk in surgery.



Yesita Rizky Firmansyah Putri is a medical doctor which currently work as Trust Grade Doctor of Neurosciences at University Hospital Coventry and Warwickshire, UK. She had postgraduate degree in Biomedicine from Brawijaya University. Her research interests are Clinical Cardiology and Neurology.



Makhyan Jibril Al-Farabi is a cardiology resident in the Department of Cardiology and Vascular Medicine Soetomo General Hospital and Faculty of Medicine, University of Airlangga. He also have postgraduate degree from University College London, Quantics School of Business and Technology and Brawijaya University. His research interests includes stem cells, clinical cardiology, cardiometabolic syndrome, healthcare entrepreneurship and AI in cardiology.

Oktaviono, et al.: Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (Allium sativum) Extract, Purple Sweet Potato (Ipomoea batatas) Extract, and Vitamin C



Ferry Sandra is a professor in the Department of Biochemistry and Molecular Biology Dental Faculty, Trisakti University Jakarta. He is also responsible to led the the BioCORE Laboratory. Previously, he was the director of Stem Cells and Cancer Institute (SCI). His research interests includes stem cells, cancer, inflammation and dentistry.

Cite this article: Oktaviono YH, Putri AY, Al-Farabi MJ, Firmansyah YR, Sandra F. Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (*Allium sativum*) Extract, Purple Sweet Potato (*Ipomoea batatas*) Extract, and Vitamin C. Pharmacog J. 2020;12(3):

Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (Allium Sativum) Extract, Purple Sweet Potato

ORIGIN	ALITY REPORT			
2 SIMILA	1% ARITY INDEX	13% INTERNET SOURCES	19% PUBLICATIONS	1% STUDENT PAPERS
PRIMAR	RY SOURCES			
1	Chusri T Extract f	Suchantabud, Tedalubmook. "Chro rom Sphagnetico cognosy Journal,	nic Toxicity of la trilobata (L.)	Leaf 2%
2	Mohamn Angliana Peripher Cells fro	Oktaviono, Djan nad Aris Widodo, n Chouw, Ferry S al Blood-derived m Stable Angina an Biomedical Jo	Yanni Dirgant andra. "Prolife Endothelial Pr Subjects", The	tara, ration of ogenitor
3	doaj.org	е		1%
4	WWW.jCC Internet Source			1%
5	WWW.WjC			1%

6	Internet Source	1%
7	Savneet Kaur. "Genetic engineering with endothelial nitric oxide synthase improves functional properties of endothelial progenitor cells from patients with coronary artery disease: an in vitro study", Basic Research in Cardiology, 05/29/2009 Publication	1%
8	Submitted to iGroup Student Paper	1%
9	A Y Putri, B S Pikir, Y H Oktaviono, F Alzahra'. "Effects of Garlic Extract (allicin) on Proliferation of Endothelial Progenitor Cells (EPC) in Patients with Stable Coronary Artery Disease", IOP Conference Series: Earth and Environmental Science, 2020 Publication	1%
10	www.id-press.eu Internet Source	1%
11	kb.psu.ac.th Internet Source	1%
12	En-Pei Isabel Chiang, Shao-Chih Chiu, Man-Hui Pai, Yi-Cheng Wang, Fu-Yu Wang, Yueh- Hsiung Kuo, Feng-Yao Tang. "Organosulfur Garlic Compounds Induce Neovasculogenesis	<1%


in Human Endothelial Progenitor Cells through a Modulation of MicroRNA 221 and the PI3-K/Akt Signaling Pathways", Journal of Agricultural and Food Chemistry, 2013

Publication

Internet Source

www.spandidos-publications.com <1% 13 Internet Source Song, Brian J., Teryn N. Sapper, Claire E. 14 Burtch, Karen Brimmer, Mark Goldschmidt, and Mario G. Ferruzzi. "Photo- and Thermodegradation of Anthocyanins from Grape and Purple Sweet Potato in Model Beverage Systems", Journal of Agricultural and Food Chemistry, 2013. Publication "American Transplant Congress 2007 Executive <1% and Program Planning Committees and Abstract Review Committees", American Journal of Transplantation, 5/2007 Publication e-sciencecentral.org 16 Internet Source www.nature.com 17 Internet Source clinicaltrials.gov 18

19	repository.tudelft.nl Internet Source	<1%
20	www.ncbi.nlm.nih.gov Internet Source	<1%
21	homepage.ntu.edu.tw Internet Source	<1%
22	www.kevinmd.com Internet Source	<1%
23	rd.springer.com Internet Source	<1%
24	Kim, Ok-Kyung, Da-Eun Nam, Ho-Geun Yoon, Sun Jung Baek, Woojin Jun, and Jeongmin Lee. "Immunomodulatory and Antioxidant Effects of Purple Sweet Potato Extract in LP-BM5 Murine Leukemia Virus-Induced Murine Acquired Immune Deficiency Syndrome", Journal of Medicinal Food, 2015. Publication	<1%
25	www.dovepress.com Internet Source	<1%
26	T. Umemura, J. Soga, T. Hidaka, H. Takemoto et al. "Aging and Hypertension Are Independent Risk Factors for Reduced Number of Circulating Endothelial Progenitor Cells", American Journal of Hypertension, 2008 Publication	<1%



polysaccharide from Medemia argun fruit", International Journal of Biological Macromolecules, 2019

Publication

Chuanguo Liu, Kun Huang, Guorong Li, 33 Pingping Wang, Chang Liu, Congcong Guo, Zongguo Sun, Jie Pan. "Ascorbic acid promotes 3T3-L1 cells adipogenesis by attenuating ERK signaling to upregulate the collagen VI", Nutrition & Metabolism, 2017

Publication

hippokratia.gr 34 Internet Source

<1%

<1%

Hui Yang. "Adiponectin promotes endothelial 35 cell differentiation from human peripheral CD14⁺ monocytes in vitro", Journal of Cellular and Molecular Medicine, 4/2006

Publication

Cheng-Hsien Lu, Hsin-Ching Lin, Chih-Cheng 36 Huang, Wei-Che Lin et al. "Increased circulating endothelial progenitor cells and anti-oxidant capacity in obstructive sleep apnea after surgical treatment", Clinica Chimica Acta, 2015 Publication

<1%

Vijayakumar R. Vishnu, Raveendran S. Renjith, 37 Archana Mukherjee, Shirly Raichal Anil et al. " Comparative Study on the Chemical Structure

<1%

and Antiproliferative Activity of Anthocyanins in Purple Root Tubers and Leaves of Sweet Potato ()", Journal of Agricultural and Food Chemistry, 2019

Publication

Sunan Wang, Shaoping Nie, Fan Zhu.
"Chemical constituents and health effects of sweet potato", Food Research International, 2016

<1%

Publication

Nai-Chen Cheng, Tsung-Yu Hsieh, Hong-Shiee Lai, Tai-Horng Young. "High glucose-induced reactive oxygen species generation promotes stemness in human adipose-derived stem cells", Cytotherapy, 2016

<1%

Publication

Exclude quotes

On

Exclude matches

Off

Exclude bibliography

-

Extracellular-Signal Regulated Kinase Signalling Pathway Mediates the Increased Proliferation of EPCs Treated with Garlic (Allium Sativum) Extract, Purple Sweet Potato

GRADEMARK REPORT	
final grade /O	GENERAL COMMENTS Instructor
PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	
PAGE 6	