

Human Umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration

by Yudi Her Oktaviono

Submission date: 07-Jun-2020 08:42PM (UTC+0800)

Submission ID: 1339322860

File name: ces_endothelial_progenitor_cells_proliferation_and_migration.pdf (1.12M)

Word count: 5895

Character count: 34179



RESEARCH ARTICLE

Human umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration [version 1; peer review: awaiting peer review]

Yudi Her Oktaviono ¹, Suryo Ardi Hutomo ¹, Makhyan Jibril Al-Farabi ¹, Angliana Chouw ², Ferry Sandra ³

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga, Soetomo General Academic Hospital, Surabaya, Indonesia

²Stem Cell Division, Prodia Laboratory, Jakarta, Indonesia

³Department of Biochemistry and Molecular Biology, Faculty of Dentistry, Universitas Trisakti, Jakarta, Indonesia

V1 First published: 05 Jun 2020, 9:537
<https://doi.org/10.12688/f1000research.23547.1>

Latest published: 05 Jun 2020, 9:537
<https://doi.org/10.12688/f1000research.23547.1>

Open Peer Review

Reviewer Status AWAITING PEER REVIEW

Any reports and responses or comments on the article can be found at the end of the article.

Abstract

Background: Human umbilical cord blood-mesenchymal stem cell (hUCB-MSC)-derived secretome is known to be able to promote neovascularization and angiogenesis, so it is also thought to have a capability to modulate endothelial progenitor cell (EPC) functions. Atorvastatin is the cornerstone of coronary artery disease (CAD) treatment which can enhance EPCs proliferation and migration. This study aims to analyze the effect of the hUCB-MSC-derived secretome and its combination with atorvastatin toward EPCs proliferation and migration.

Methods: EPCs were isolated from a CAD patient's peripheral blood. Cultured EPCs were divided into a control group and treatment group of 2.5 μ M atorvastatin, hUCB-MSC-derived secretome (2%, 10%, and 20% concentration) and its combination. EPCs proliferation was evaluated using an MTT cell proliferation assay, and EPC migration was evaluated using a Transwell migration assay kit.

Results: This research showed that hUCB-MSC-derived secretomes significantly increase EPC proliferation and migration in a dose-dependent manner. The high concentration of hUCB-MSC-derived secretome were shown to be superior to atorvastatin in inducing EPC proliferation and migration ($p < 0.001$). A combination of the hUCB-MSC-derived secretome and atorvastatin shown to improve EPCs proliferation and migration compared to hUCB-MSC-derived secretome treatment or atorvastatin alone ($p < 0.001$).

Conclusions: This study concluded that the hUCB-MSC-derived secretome work synergistically with atorvastatin treatment in improving EPCs proliferation and migration.

Keywords

coronary artery disease, endothelial progenitor cells, mesenchymal stem cells, secretome, statins

Corresponding author: Yudi Her Oktaviono (yoktaviono@gmail.com)

Author roles: **Oktaviono YH:** Conceptualization, Formal Analysis, Investigation, Methodology, Resources, Validation, Writing – Review & Editing; **Hutomo SA:** Investigation, Methodology, Project Administration, Visualization, Writing – Original Draft Preparation; **Al-Farabi MJ:** Project Administration, Writing – Review & Editing; **Chouw A:** Investigation, Software; **Sandra F:** Conceptualization, Data Curation, Funding Acquisition, Methodology, Resources, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This research was funded by the Prodia Education and Research Institute.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Oktaviono YH *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Oktaviono YH, Hutomo SA, Al-Farabi MJ *et al.* **Human umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration [version 1; peer review: awaiting peer review]** F1000Research 2020, 9:537 <https://doi.org/10.12688/f1000research.23547.1>

First published: 05 Jun 2020, 9:537 <https://doi.org/10.12688/f1000research.23547.1>

Introduction

Coronary artery disease (CAD) is the leading cause of mortality and morbidity worldwide¹. It is responsible for the deaths of 7.2 million people or 12.2% of total deaths per year worldwide. Despite advancement in CAD management (e.g. novel antiplatelet therapy, coronary stents, percutaneous coronary intervention techniques and devices, and coronary artery bypass surgery), there are some clinical subsets of CAD which remain untreatable such as ischemic cardiomyopathy, refractory angina, and patient which cannot undergo revascularization due to clinical and anatomical complexity^{2,3}.

It is already known that CAD is caused by atherosclerosis, which is followed by reduced levels of circulating endothelial progenitor cells (EPCs)⁴. EPCs can differentiate into mature endothelial cells and also promote endothelial repair. Hence, increasing circulating EPC levels is proven to improve endothelial function⁵. EPCs also had a critical role in the stimulation of angiogenesis and vasculogenesis. Hence, increasing EPC proliferation and migration may reduce ischemia and improve myocardial performance^{4,6}.

Regenerative treatment for CAD using stem cells has been extensively studied in the last decade⁷. However, these treatments faced challenges of low engraftment, poor survival, and low differentiation of the transplanted cells. Despite regenerative treatment shown to be promising *in vitro*, clinical studies showed unsatisfying results⁸. Hence, the researcher started to shift regenerative treatment from cell based-treatment into cell-free treatment using paracrine stimulation⁹. Nowadays, the usage of cell-free therapeutics as a regenerative therapy in cardiovascular diseases also started to be emerged⁹.

The secretome is the wide array variety of paracrine factors produced by mesenchymal stem cells (MSCs). Human umbilical cord blood mesenchymal stem cells (hUCB-MSCs) derived secretome was proven could promote neovascularization, angiogenesis¹⁰⁻¹³ and improved cardiac systolic function by protecting myocardial cells from apoptosis¹⁴. However, using this approach to improve neovascularization is yet to be investigated. Hence, it is hypothesized that increasing EPCs proliferation and migration by the hUCB-MSC-derived secretome may be responsible for this effect. Statins, through their pleiotropic effect, are the cornerstone of CAD treatment. Atorvastatin is one of the most prescribed statins, whose ability to modulate EPCs proliferation and migration has already been well studied in both laboratory and clinical settings¹⁵⁻¹⁷. Furthermore, this study aims to compare the effect of the hUCB-MSC-derived secretome, atorvastatin and the two in combination in modulating EPC proliferation and migration.

Methods

Sample collection

A 50–100 mL peripheral blood sample was obtained from a patient with CAD. The patient was recruited from the outpatient cardiovascular clinic at Pusat Pelayanan Jantung Terpadu, Dr. Soetomo General Hospital, Surabaya, in March 2020. The inclusion criteria were as follows: male, aged 40–59 years old,

history of chronic ischemic heart disease as proven by coronary angiography results that showed >50% stenosis of left main coronary artery or >70% of other coronary arteries. The exclusion criteria were as follows: a history of percutaneous coronary intervention procedures or coronary artery bypass grafting surgery, acute coronary syndromes, and anemia.

This study protocol has an ethical clearance from the Health Research Ethics Committee of Dr. Soetomo General Hospital, Surabaya (No.1567/KEPK/X/2019, approved on 8 October 2019). The included subjects provided written informed consent before subject recruitment. All details which include personal information were omitted.

HUCB-MSCs-derived secretome preparation

The HUCB-MSCs-derived secretome was prepared according to the previous study¹⁸. HUCB-MSCs (3H Biomedical AB, Uppsala, Sweden) was cultured in MesenCult™ MSC Basal medium, supplemented with MesenCult™ Stimulatory supplement (StemCell Technologies Inc., Vancouver, Canada), and also added with penicillin and streptomycin. Upon reaching 80% confluency, the media was replaced with MesenCult™ MSC Basal medium (supplement-free media) and incubated for 24 hours. The media was collected and centrifuged. The supernatant was used as a conditioned medium that contained hUCB-MSCs-derived secretome.

Isolation and culture of EPCs

Peripheral blood mononuclear cells (PBMCs) were isolated by density centrifugation of CAD patient's peripheral blood using Histopaque-1077 (Sigma-Aldrich, USA). After centrifugation of peripheral blood, PBMCs then cultured with STEMLINE-II hematopoietic stem cell expansion medium (Sigma-Aldrich, USA) supplemented with stem cell factor, thrombopoietin, Flt-3 ligand, vascular endothelial growth factor, and interleukin-6. A total of 5×10^6 mononuclear cells/ml were seeded into fibronectin-coated 6-well plate dish and cultured at 37°C and 5% CO₂ levels for 5 days. Non-adherent cells were then transferred for the proliferation and migration assay. After five days of culture, EPCs were confirmed using FITC-labeled anti-human CD34 antibody (animal source was mouse, 5 µL antibody was diluted at 500 µL per 1×10^6 cells; catalog number 60013FI, Gene ID: 947, by StemCell Technologies Inc., Vancouver, Canada) staining and examined with immunofluorescence microscopy.

Treatment of EPCs

Cultured EPCs were divided into eight treatment groups for each proliferation and migration assays. Those treatment include control group, 2.5 µM atorvastatin, low (2%), medium (10%) and high (20%) doses of hUCB-MSC-derived secretome, and combination of 2.5 µM atorvastatin with each dose of the hUCB-MSC-derived secretome. There were n=5 replications made from each treatment.

EPCs proliferation assay

The MTT cell proliferation assay kit (Sigma-Aldrich, St Louis, MO, USA) was used to measure EPCs proliferation as described previously¹⁹. Treated EPCs were added with MTT reagent

2 and incubated in a 37°C incubator with 5% CO₂ for 4 hours. Proliferation was determined from the reduction of tetrazolium (MTT) into insoluble formazan product by viable EPCs mitochondria. Absorbance was measured with a microplate reader at 595 nm wavelength. EPCs proliferation was measured as optical density (OD). MTT assay was measured at day 3 after reagent addition.

EPCs migration assay

EPCs migration was evaluated using the 24-mm diameter insert, 3-µm pore size, 6-well Transwell migration assay kit (Corning, USA). A total of 5×10⁵ cultured EPCs were placed in the upper part of the Transwell migration assay kit. Next, 2 mL of EPC media and each treatment were added in the lower chamber compartment and then incubated for 24 hours at 37°C. Non-migratory cells were removed manually. On the receiver plate, the new basal medium was placed and added 500 µL of trypsin + EDTA solution 0.5%, followed by 10 minutes incubation. Then, cells on the bottom surface of the membrane were stained with Giemsa and cell images were obtained on a light microscope and counted manually in n=5 random fields/sample.

6 Statistical analysis

Statistical analyses were conducted using SPSS Statistics 23.0 to detect significance level at p<0.05. One-way ANOVA was used to compare groups, with Fisher's least significant difference (LSD) post hoc test. Kruskal-Wallis test was used if there are violations to the assumption of normality and the assumption of homogeneity of variance. Correlation between variables was obtained using Spearman's correlation followed by a linear regression test.

Results

Baseline characteristics and demography of CAD patient

Clinical examination, blood sampling, electrocardiography, echocardiography and coronary angiography was conducted and evaluated in order to examine the inclusion and exclusion criteria. Our sample had a 1-year history of coronary artery disease, he suffered from refractory chest pain despite the optimum medical therapy. The coronary angiography showed complex lesion (three-vessel disease with chronic total occlusion) which was not amenable to undergo revascularization. The baseline characteristics of the patient are presented in Table 1.

Table 1. Characteristics of the patient.

Variables	Result
Sex	Male
3 Age	59 years old
Body Mass Index (BMI)	27.3
Blood pressure	140/90
Heart rate	90 beats per minute
Electrocardiography	Sinus rhythm, pathological Q-waves at V-V6 leads.
Laboratory	
20 Total cholesterol (mg/dL)	240
Triglyceride (mg/dL)	131
LDL (mg/dL)	140
HDL (mg/dL)	55
Hemoglobin (mg/dL)	14.2
Blood glucose (mg/dL)	142
Echocardiography	
Left ventricle ejection fraction	41% (teich); 36% (biplane)
Left ventricle internal diameter	5.8 cm
Wall motion	Hypokinesia at anterior, anteroseptal, inferoseptal, other segments kinetic was normal
Valves	mild mitral regurgitation
Coronary Angiography	
9 Left main coronary artery (LMCA)	normal
Left anterior descending artery (LAD)	70% stenosis at ostial, chronic total occlusion (CTO) at distal
Left circumflex coronary artery (LCX)	70% stenosis at proximal, CTO at distal, grade 2 collaterals from LCX to RCA
Right coronary artery (RCA)	CTO at proximal

EPC characteristics

EPCs were successfully isolated and cultured from the CAD patient's peripheral blood. It was confirmed through light microscopy and an immunofluorescence assay (Figure 1). Raw images are available as *Underlying data*²⁰.

hUCB-MSCs-derived secretome and atorvastatin increase EPCs proliferation

EPCs were evaluated using the MTT proliferation assay. As shown in Figure 2, both atorvastatin and hUCB-MSCs-derived secretome treatment groups at all doses increase EPCs proliferation compared to the control ($p < 0.05$, ANOVA). hUCB-MSC-derived secretome treatment showed a dose-dependent relationship with EPCs proliferation. At medium (10%) and high (20%) doses, hUCB-MSC-derived secretome was shown to elicit superior EPC proliferation than atorvastatin (OD 1.252 ± 0.104 and 1.585 ± 0.029 , respectively, vs 0.738 ± 0.025 ; $p < 0.01$). Raw absorbance data for MTT assays are available as *Underlying data*²⁰.

Pearson's correlation showed a significant and strong correlation between hUCB-MSCs-derived secretome treatment with EPC proliferation ($r = 0.954$; $p < 0.001$). The linear regression test showed an R^2 of 0.910.

Combination of hUCB-MSCs-derived secretome and atorvastatin increase EPCs proliferation compared with single treatment

Figure 2 showed the combination of atorvastatin and hUCB-MSC-derived secretome at the dose of 2%, 10% and 20%

concentration have significantly higher EPCs proliferation compared to atorvastatin alone (OD 0.803 ± 0.046 , 1.298 ± 0.075 and 1.761 ± 0.419 vs 0.738 ± 0.025 , $p < 0.05$). In addition, combination of hUCB-MSC-derived secretome at dose of 2%, 10% and 20% with atorvastatin showed higher EPCs proliferation compared to hUCB-MSC-derived secretome alone (OD 0.803 ± 0.046 vs 0.713 ± 0.049 , 1.298 ± 0.075 vs 1.252 ± 0.104 and 1.761 ± 0.419 vs 1.585 ± 0.029 , $p < 0.05$). The combination group showed a significant and very strong correlation with EPC proliferation ($r = 0.973$; $p < 0.001$). Linear regression test showed R^2 of 0.947.

hUCB-MSCs-derived secretome and atorvastatin increase EPCs migration

EPCs migration from each treatment group was analyzed using the Transwell migration assay. As shown in Figure 3, EPC treatment with atorvastatin and all doses of hUCB-MSCs-derived secretome significantly increase EPC migration compared to the control group ($p < 0.05$, ANOVA). Treatment with $2.5 \mu\text{M}$ atorvastatin has significantly higher EPCs migration than low (2%) and medium (10%) doses of hUCB-MSC-derived secretome (34.40 ± 3.05 vs 17.20 ± 1.92 and 27.00 ± 4.00 , $p < 0.05$). However, high doses (20%) of hUCB-MSC-derived secretome showed significantly higher migrated EPCs than atorvastatin (51.00 ± 5.15 vs 34.40 ± 3.05 , $p < 0.001$). Raw cell counts used to assess migration are available as *Underlying data*²⁰.

Pearson's correlation showed a significant and very strong correlation between hUCB-MSCs-derived secretome treatment with EPC migration ($r = 0.968$; $p < 0.001$). The linear regression test showed an R^2 of 0.937.




Figure 1. Immunofluorescence characterization of cultured EPCs. (A) DAPI staining of cultured EPCs showed viable cells through blue fluorescent of cells nuclei. (B) EPCs were confirmed, using FITC-labeled anti-human CD34 expression on immunofluorescence microscope. (C) Merged view of DAPI and FITC stained cells. (D) The light microscope view showed the spindle shape of EPCs. The white bar represents 50 μm .




Figure 2. Comparison of EPC proliferation effects among all treatment groups (see text). ^aSignificant difference compared to the control group ($p < 0.001$). ^bSignificant difference compared to the 2.5 µM atorvastatin group ($p < 0.001$). ^cSignificant difference compared to the 2% hUCB-MSC-derived secretome group, ($p < 0.001$). ^dSignificant difference compared to the 10% hUCB-MSC-derived secretome group ($p < 0.001$). ^eSignificant difference compared to the 20% hUCB-MSC-derived secretome group ($p < 0.001$). ^fSignificant difference compared to the combination of 2% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$). ^gSignificant difference compared to the combination of 10% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$). ^hSignificant difference compared to the combination of 20% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$)




Figure 3. Comparison of EPCs migration effects among all treatment groups (see text). ^aSignificant difference compared to the control group ($p < 0.001$). ^bSignificant difference compared to the 2.5 µM atorvastatin group ($p < 0.001$). ^cSignificant difference compared to the 2% hUCB-MSC-derived secretome group ($p < 0.001$). ^dSignificant difference compared to the 10% hUCB-MSC-derived secretome group ($p < 0.001$). ^eSignificant difference compared to the 20% hUCB-MSC-derived secretome group ($p < 0.001$). ^fSignificant difference compared to the combination of 2% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$). ^gSignificant difference compared to the combination of 10% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$). ^hSignificant difference compared to the combination of 20% hUCB-MSC-derived secretome and atorvastatin group, ($p < 0.001$)

Combination of hUCB-MSCs-derived secretome and atorvastatin increase EPCs migration compared with single treatment

In Figure 3, EPCs migration was significantly higher in combination treatment groups (atorvastatin and hUCB-MSC-derived secretome) at 2%, 10%, and 20% doses compared to the atorvastatin alone (38.20 ± 3.49 , 50.20 ± 5.31 and 76.40 ± 7.50 vs 34.40 ± 3.05 , $p < 0.001$).

Combination of hUCB-MSC-derived secretome with atorvastatin also showed higher EPCs migration than the hUCB-MSC-derived secretome-only group at 2%, 10% and 20% concentrations (38.3 ± 3.49 vs 17.20 ± 1.92 , 50.20 ± 5.31 vs 27.00 ± 4.00 and 76.40 ± 7.50 vs 51.00 ± 5.15 , respectively; all $p < 0.001$). The combination of high-dose (20%) hUCB-MSC-derived secretome and atorvastatin had the highest number of migrated EPC ($76.4 \pm 7.50 \times 10^3$ cells). The combination group had a significant and very strong correlation with EPC migration ($r = 0.970$; $p < 0.001$). The linear regression test showed an R^2 of 0.942.

Discussion

This research showed that treatment with hUCB-MSC-derived secretome, atorvastatin and a combination of the two increased the proliferation and migration of EPCs (isolated from CAD patient's peripheral blood). hUCB-MSC-derived secretome enhances EPCs proliferation and migration in a dose-dependent manner. The combination of hUCB-MSC-derived secretome and atorvastatin was shown to be superior to atorvastatin or hUCB-MSC-derived secretome alone.

In this research, hUCB-MSC-derived secretome treatment increased EPC proliferation in a dose-dependent manner, with the concentrations of 10% and 20% shown to be superior to atorvastatin. Previous studies showed that atorvastatin treatment is superior to other statins at improving EPC proliferation²¹⁻²³. The hUCB-MSC-derived secretome is also composed of cytokines, chemokines, growth factors, proteins, and extracellular vesicles which may be involved in EPCs proliferation and migration^{13,24}. Vascular endothelial growth factor (VEGF), stromal-derived factor-1 (SDF-1), insulin-like growth factor (IGF-1) are contained in the hUCB-MSC-derived secretome which may be involved in increasing EPC proliferation²⁵. VEGF has been shown to improve the proliferation and differentiation of EPCs through activation of Ras signaling, and the MAPK/ERK pathway²⁶⁻²⁸. SDF-1 and IGF 1 also been shown to increase the EPCs proliferation in response to the PI3K/protein kinase B signaling pathway and promote angiogenesis²⁹⁻³¹. Hence, it is suggested that hUCB-MSC-derived secretome treatment is beneficial to improve EPC proliferation which may involve MAPK/ERK and PI3K/protein kinase B pathway.

HUCB-MSC-derived secretome treatment shown to increase EPCs migration in a dose-dependent manner, with a concentration of 20% shown to be superior to atorvastatin. Similarly, the previous study showed secretome-derived from placental-MSCs is able to significantly increase EPCs migration³². The HUCB-MSC-derived secretome contains pro-angiogenic factors,

such as human angiopoietin-1 (Ang-1), hepatocyte growth factor (HGF), insulin-like growth factor I (IGF-I), prostaglandin E2 (PGE2), transforming growth factor-beta 1 (TGF- β 1), vascular cell adhesion protein 1 (VCAM-1) and vascular endothelial growth factor (VEGF)³³. MSCs also have immunomodulatory and anti-inflammatory properties, as it contributes to the maintenance of self-renewal capacity through E-Prostanoid 2 (EP2)³⁴ and immune cell activation and maturation, including CD4⁺ helper T cells, B cells, dendritic cells, natural killer cells, monocytes and macrophages³⁵. The HUCB-MSC-derived secretome also has a higher anti-inflammatory effect than other MSCs³⁶ and antioxidant properties, as proven in previous studies conducted in renal injury³⁷ and ischemic stroke³⁸. Inflammatory stimuli and oxidative stress are also known to impair EPCs migration³⁹. Chemoattractant gradient was an important driving factor to induce EPCs mobilization. Hence, a high concentration of growth factors in the HUCB-MSC-derived secretome increase the gradient between the top and lower parts of the Transwell migration assay may lead into an increase of EPCs migration. taken together, wide array molecules and multiple possible pathways involved in HUCB-MSCs secretome treatment seem to be responsible for its superiority against atorvastatin.

The synergistic effect of the HUCB-MSC-derived secretome with atorvastatin in enhancing EPCs proliferation and migration was demonstrated in this study. These combinations significantly increase EPCs proliferation and migratory activity by up to two-fold. Previously, The combination of MSCs with another compound, including statins, was shown to have beneficial effects in angiogenesis and neovascularization⁴⁰⁻⁴². Co-culture of MSCs and EPCs have been shown to demonstrate improved EPC proliferation and migration, and enhance their angiogenic capacity^{43,44}. However, the exact mechanism of these combinations to improve EPCs proliferation and migration is yet to be investigated. It is speculated that the involvement of multiple pathways may be responsible for its superiority against HUCB-MSCs-derived secretome or atorvastatin alone.

A mitogen-activated protein kinase (MAPK) pathway has been known to play a role in increasing EPCs proliferation⁴⁵. Cell cycle progression through increased Cyclin D1 expression mediated by PI3K/Akt and MAPK pathway also involved in EPCs proliferation⁴⁶. While increasing microRNA 221/222 expression shown to reduce EPCs proliferation capabilities⁴⁷. HUCB-MSC-derived secretome treatment was speculated to improve EPCs proliferation through MAPK/ERK and PI3K/protein kinase B pathway. While atorvastatin improves EPCs proliferation through downregulation of microRNA 221/222 expression⁴⁷. The involvement of these multiple pathways may result in higher EPCs proliferation in the combination group.

Enhanced growth factor levels through hUCB-MSC-derived secretome treatment will augment the chemoattractant gradient³⁷, thus leading to enhanced migration of EPCs. The anti-inflammatory and antioxidant properties of hUCB-MSC-derived secretome also speculated to improve EPCs migration^{35,37}. While atorvastatin can increase the production of endothelial nitric oxide synthase and nitric oxide, which reduces the

oxidative stress that impairs EPCs migration⁴². Statin also could prevent EPCs senescence by upregulating TRF2 of EPCs, hence enhance migratory capacity⁴⁸. Those facts suggested that the combination of hUCB-MSC-derived secretome and atorvastatin will have superior EPCs migration through the involvement of multiple pathways.

In summary, hUCB-MSC-derived secretome may be developed and combined with atorvastatin treatment in CAD patients to improve EPCs proliferation and migration. However, this research did not measure the exact composition of the hUCB-MSC-derived secretome. The previous study showed that the secretome from another type of MSC can increase EPCs migration but not EPCs proliferation⁴⁹. Hence, further research should be directed to identify the substance within the hUCB-MSC-derived secretome which is responsible for increasing EPC proliferation and migration, and compare it with other MSC secretomes. Further research should also verify the multiple pathways which may be responsible for the improvement of EPCs proliferation and migration in the combination group.

Conclusions

High dose hUCB-MSC-derived secretome outperforms atorvastatin to improve EPC proliferation and migration. A combination of hUCB-MSC-derived secretome with atorvastatin seems to be beneficial in promoting neovascularization through improved

EPCs proliferation and migration effect compared to hUCB-MSC-derived secretome or atorvastatin alone.

Data availability

Underlying data

Figshare: Human umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration. <https://doi.org/10.6084/m9.figshare.12186507.v20>.

This project contains the following underlying data:

- RAW Data - F1000 revisi2 by SAH (XLSX). (Raw absorbance data from MTT proliferation assay and cell counts from the Transwell migration assay.)
- RAW DATA f1000 (ZIP). (Raw images generated in this study, including images used to generate cell counts and raw immunofluorescence images.)

Data are available under the terms of the [Creative Commons Attribution 4.0 International license \(CC-BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).

Acknowledgements

The authors are thankful to Christian Pramudita, MD., Ilma Alfia Isaridha, MD., Melly Susanti MD., and Dwi Fachrul MD., for invaluable support in this project.

References

- Benjamin EJ, Muntner P, Alonso A, et al.: **Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association.** *Circulation.* 2019; **139**(10): e56–e528. [PubMed Abstract](#) | [Publisher Full Text](#)
- Sainsbury PA, Fisher M, de Silva R: **Alternative interventions for refractory angina.** *Heart.* 2017; **103**(23): 1911–1922. [PubMed Abstract](#) | [Publisher Full Text](#)
- Cheng K, Sainsbury P, Fisher M, et al.: **Management of Refractory Angina Pectoris.** *Eur Cardiol.* 2016; **11**(2): 69–76. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Siddique A, Shantsila E, Lip GY, et al.: **Endothelial progenitor cells: what use for the cardiologist?** *J Angiogenesis Res.* 2010; **2**(1): 6. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Wils J, Favre J, Bellien J: **Modulating putative endothelial progenitor cells for the treatment of endothelial dysfunction and cardiovascular complications in diabetes.** *Pharmacol Ther.* 2017; **170**: 98–115. [PubMed Abstract](#) | [Publisher Full Text](#)
- George AL, Bangalore-Prakash P, Rajoria S, et al.: **Endothelial progenitor cell biology in disease and tissue regeneration.** *J Hematol Oncol.* 2011; **4**: 24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fisher SA, Doree C, Mathur A, et al.: **Stem cell therapy for chronic ischaemic heart disease and congestive heart failure.** *Cochrane Database Syst Rev.* 2016; **12**: CD007888. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Madonna R, Van Laake LW, Davidson SM, et al.: **Position Paper of the European Society of Cardiology Working Group Cellular Biology of the Heart: cell-based therapies for myocardial repair and regeneration in ischemic heart disease and heart failure.** *Eur Heart J.* 2016; **37**(23): 1789–1798. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gnecchi M, Zhang Z, Ni A, et al.: **Paracrine mechanisms in adult stem cell signaling and therapy.** *Circ Res.* 2008; **103**(11): 1204–1219. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vasa M, Fichtschler S, Adler K, et al.: **Increase in circulating endothelial progenitor cells by statin therapy in patients with stable coronary artery disease.** *Circulation.* 2001; **103**(24): 2885–90. [PubMed Abstract](#) | [Publisher Full Text](#)
- Lievadot J, Murasawa S, Kureishi Y, et al.: **HMG-CoA reductase inhibitor mobilizes bone marrow-derived endothelial progenitor cells.** *J Clin Invest.* 2001; **108**(3): 399–405. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Dimmeler S, Aicher A, Vasa M, et al.: **HMG-CoA reductase inhibitors (statins) increase endothelial progenitor cells via the PI 3-kinase/Akt pathway.** *J Clin Invest.* 2001; **108**(3): 391–7. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gallina C, Turinetto V, Giachino C: **A New Paradigm in Cardiac Regeneration: The Mesenchymal Stem Cell Secretome.** *Stem Cells Int.* 2015; **2015**: 765846. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Konala VB, Mamidi MK, Bhonde R, et al.: **The current landscape of the mesenchymal stromal cell secretome: A new paradigm for cell-free regeneration.** *Cytotherapy.* 2016; **18**(1): 13–24. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Vizoso FJ, Eiro N, Gid S, et al.: **Mesenchymal Stem Cell Secretome: Toward Cell-Free Therapeutic Strategies in Regenerative Medicine.** *Int J Mol Sci.* 2017; **18**(9): pii: E1852. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Phelps J, Sanati-Nezhad A, Ungrin M, et al.: **Bioprocessing of Mesenchymal Stem Cells and Their Derivatives: Toward Cell-Free Therapeutics.** *Stem Cells Int.* 2018; **2018**: 9415367. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Zhao Y, Sun X, Cao W, et al.: **Exosomes Derived from Human Umbilical Cord Mesenchymal Stem Cells Relieve Acute Myocardial Ischemic Injury.** *Stem Cells Int.* 2015; **2015**: 761643. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Sandra F, Sudiono J, Sidharta EA, et al.: **Conditioned Media of Human Umbilical**

- Cord Blood Mesenchymal Stem Cell-derived Secretome Induced Apoptosis and Inhibited Growth of HeLa Cells.** *Indones Biomed J.* 2014; 6(1): 57.
[Publisher Full Text](#)
19. Oktaviono YH, Al-farabi MJ, Oliva L, *et al.*: **Preliminary Study: Purple Sweet Potato Extract Seems to Be Superior to Increase the Migration of Impaired Endothelial Progenitor Cells Compared to l-Ascorbic Acid.** *Sci Pharm.* 2017; 87(3): 16.
[Publisher Full Text](#)
 20. Al Farabi, Makhyan Jibril: **Human umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration.** *figshare.* Dataset. 2020.
<http://www.doi.org/10.6084/m9.figshare.12186507.v2>
 21. Meuthia F, Oktaviono YH, Soemantri D: **Effects of Statins on Endothelial Progenitor Cell Proliferation from Peripheral Blood of Stable Coronary Artery Disease Patient.** *J Kardiol Indones.* 2017; 38(1): 6–12.
[Publisher Full Text](#)
 22. Oktaviono YH, Savitri TVR, Soemantri D: **Rosuvastatin is Superior Compared to Simvastatin and Atorvastatin to Induce Endothelial Progenitor Cells Migration.** *J Clin Diagnostic Res.* 2019; 1077: 5–8.
[Publisher Full Text](#)
 23. Rawlings R, Nohria A, Liu PY, *et al.*: **Comparison of effects of rosuvastatin (10 mg) versus atorvastatin (40 mg) on rho kinase activity in caucasian men with a previous atherosclerotic event.** *Am J Cardiol.* 2009; 103(4): 437–441.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 24. Eleuteri S, Fierabracci A: **Insights Into the Secretome of Mesenchymal Stem Cells and Its Potential Applications.** *Int J Mol Sci.* 2019; 20(18): 4597.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 25. Kwon HM, Hur SM, Park KY, *et al.*: **Multiple paracrine factors secreted by mesenchymal stem cells contribute to angiogenesis.** *Vascul Pharmacol.* 2014; 63(1): 19–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
 26. Xu J, Liu X, Jiang Y, *et al.*: **MAPK/ERK signalling mediates VEGF-induced bone marrow stem cell differentiation into endothelial cell.** *J Cell Mol Med.* 2008; 12(6A): 2395–2406.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 27. Ge Q, Zhang H, Hou J, *et al.*: **VEGF secreted by mesenchymal stem cells mediates the differentiation of endothelial progenitor cells into endothelial cells via paracrine mechanisms.** *Mol Med Rep.* 2018; 17(1): 1667–1675.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 28. Arutyunyan I, Fatkhudinov T, Kananykhina E, *et al.*: **Role of VEGF-A in angiogenesis promoted by umbilical cord-derived mesenchymal stromal/stem cells: *in vitro* study.** *Stem Cell Res Ther.* 2016; 7: 46.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 29. Chen SL, Sheiban I, Xu BO, *et al.*: **Impact of the complexity of bifurcation lesions treated with drug-eluting stents: the DEFINITION study (Definitions and impact of complex bifurcation lesions on clinical outcomes after percutaneous coronary intervention using drug-eluting stents).** *JACC Cardiovasc Interv.* 2014; 7(11): 1266–1276.
[PubMed Abstract](#) | [Publisher Full Text](#)
 30. Yu H, Feng Y: **The potential of statin and stromal cell-derived factor-1 to promote angiogenesis.** *Cell Adh Migr.* 2008; 2(4): 254–257.
[Publisher Full Text](#)
 31. Hou J, Peng X, Wang J: **Mesenchymal stem cells promote endothelial progenitor cell proliferation by secreting insulin-like growth factor-1.** *Mol Med Rep.* 2017; 16(2): 1502–1508.
[PubMed Abstract](#) | [Publisher Full Text](#)
 32. Kamprorn W, Kheolamai P, U-Pratya Y: **Endothelial Progenitor Cell Migration-Enhancing Factors in the Secretome of Placental-Derived Mesenchymal Stem Cells.** *Stem Cells Int.* 2016; 2016: 2514326.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 33. Wu M, Zhang R, Zou Q, *et al.*: **Comparison of the Biological Characteristics of Mesenchymal Stem Cells Derived from the Human Placenta and Umbilical Cord.** *Sci Rep.* 2018; 8(1): 5014.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 34. Lee BC, Kim HS, Shin TH, *et al.*: **PG2 2 maintains self-renewal of human adult stem cells via EP2-mediated autocrine signaling and its production is regulated by cell-to-cell contact.** *Sci Rep.* 2016; 6: 26298.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 35. Murphy MB, Moncivais K, Caplan AI: **Mesenchymal stem cells: Environmentally responsive therapeutics for regenerative medicine.** *Exp Mol Med.* 2013; 45(11): e54.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 36. Jin HJ, Bae YK, Kim M, *et al.*: **Comparative analysis of human mesenchymal stem cells from bone marrow, adipose tissue, and umbilical cord blood as sources of cell therapy.** *Int J Mol Sci.* 2013; 14(9): 17986–18001.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 37. Lee KH, Tseng WC, Yang CY, *et al.*: **The Anti-Inflammatory, Anti-Oxidative, and Anti-Apoptotic Benefits of Stem Cells in Acute Ischemic Kidney Injury.** *Int J Mol Sci.* 2019; 20(14): pii: E3529.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 38. Cunningham CJ, Redondo-castro E, Allan SM: **The therapeutic potential of the mesenchymal stem cell secretome in ischaemic stroke.** *J Cereb Blood Flow Metab.* 2018; 38(8): 1276–1292.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Tousoulis D, Andreou I, Antoniadou C, *et al.*: **Role of inflammation and oxidative stress in endothelial progenitor cell function and mobilization: therapeutic implications for cardiovascular diseases.** *Atherosclerosis.* 2008; 201(2): 236–247.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Luo R, Lu Y, Liu J, *et al.*: **Enhancement of the efficacy of mesenchymal stem cells in the treatment of ischemic diseases.** *Biomed Pharmacother.* 2019; 109(1): 2022–2034.
[PubMed Abstract](#) | [Publisher Full Text](#)
 41. Kwon YW, Heo SC, Jeong GO, *et al.*: **Tumor necrosis factor- α -activated mesenchymal stem cells promote endothelial progenitor cell homing and angiogenesis.** *Biochim Biophys Acta.* 2013; 1832(12): 2136–2144.
[PubMed Abstract](#) | [Publisher Full Text](#)
 42. Song L, Yang YJ, Dong QT, *et al.*: **Atorvastatin enhance efficacy of mesenchymal stem cells treatment for swine myocardial infarction via activation of nitric oxide synthase.** *PLoS One.* 2013; 8(5): 1–12.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 43. Liang T, Zhu L, Gao W, *et al.*: **Coculture of endothelial progenitor cells and mesenchymal stem cells enhanced their proliferation and angiogenesis through PDGF and Notch signaling.** *FEBS Open Bio.* 2017; 7: 1722–1736.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Li Z, Yang A, Yin X, *et al.*: **Mesenchymal stem cells promote endothelial progenitor cell migration, vascularization, and bone repair in tissue-engineered constructs via activating CXCR2-Src-PKL/Vav2-Rac1.** *FASEB J.* 2014; 400(1–2): 201–206.
[PubMed Abstract](#) | [Publisher Full Text](#)
 45. Sandra F, Oktaviono YH, Widodo MA: **Endothelial progenitor cells proliferated via MEK-dependent p42 MAPK signaling pathway.** *Mol Cell Biochem.* 2015; 400(1–2): 201–6.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Qiu C, Xie Q, Zhang D, *et al.*: **GM-CSF induces cyclin D1 expression and proliferation of endothelial progenitor cells via PI3K and MAPK signaling.** *Cell Physiol Biochem.* 2014; 63(1): 19–28.
[PubMed Abstract](#) | [Publisher Full Text](#)
 47. Minami Y, Satoh M, Maesawa C, *et al.*: **Effect of atorvastatin on microRNA 221 / 222 expression in endothelial progenitor cells obtained from patients with coronary artery disease.** *Eur J Clin Invest.* 2009; 39(5): 359–67.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Assmus B, Urbich C, Aicher A, *et al.*: **HMG-CoA reductase inhibitors reduce senescence and increase proliferation of endothelial progenitor cells via regulation of cell cycle regulatory genes.** *Circ Res.* 2003; 92(9): 1049–55.
[PubMed Abstract](#) | [Publisher Full Text](#)
 49. Kamprorn W, Kheolamai P, U-Pratya Y, *et al.*: **Effects of mesenchymal stem cell-derived cytokines on the functional properties of endothelial progenitor cells.** *Eur J Cell Biol.* 2018; 95(3–5): 153–63.
[PubMed Abstract](#) | [Publisher Full Text](#)

The benefits of publishing with F1000Research:

- Your article is published within days, with no editorial bias
- You can publish traditional articles, null/negative results, case reports, data notes and more
- The peer review process is transparent and collaborative
- Your article is indexed in PubMed after passing peer review
- Dedicated customer support at every stage

For pre-submission enquiries, contact research@f1000.com

F1000Research

Human Umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration

ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Douglas Kell, Marnie Potgieter, Etheresia Pretorius. "Individuality, phenotypic differentiation, dormancy and 'persistence' in culturable bacterial systems: commonalities shared by environmental, laboratory, and clinical microbiology", F1000Research, 2015
Publication 1%
- 2 Yudi Her Oktaviono, Djanggan Sargowo, Mohammad Aris Widodo, Yanni Dirgantara, Angliana Chouw, Ferry Sandra. "Proliferation of Peripheral Blood-derived Endothelial Progenitor Cells from Stable Angina Subjects", The Indonesian Biomedical Journal, 2014
Publication 1%
- 3 www.postershop.co.uk
Internet Source 1%
- 4 Yong, P.F.K.. "Selective deficits in blood dendritic cell subsets in common variable immunodeficiency and X-linked 1%

agammaglobulinaemia but not specific
polysaccharide antibody deficiency", Clinical
Immunology, 200804

Publication

5	pesquisa.bvsalud.org Internet Source	1 %
6	lipidworld.biomedcentral.com Internet Source	<1 %
7	menssanaresearch.com Internet Source	<1 %
8	www.nature.com Internet Source	<1 %
9	dict.cnki.net Internet Source	<1 %
10	"Tuesday, 30 August 2011", European Heart Journal, 08/02/2011 Publication	<1 %
11	content.iospress.com Internet Source	<1 %
12	journals.lww.com Internet Source	<1 %
13	www.ncbi.nlm.nih.gov Internet Source	<1 %
14	techscience.com Internet Source	<1 %

- 15 www.cordbloodresearchnews.com Internet Source <1 %
-
- 16 www.hindawi.com Internet Source <1 %
-
- 17 P. Habertzettl, D.J. Conklin, T.E. O'Toole. "Endothelial Progenitor Cells: Properties, Function, and Response to Toxicological Stimuli", Elsevier BV, 2018 Publication <1 %
-
- 18 Rachael A Panizzo, David G Gadian, Jane C Sowden, Jack A Wells, Mark F Lythgoe, Patrizia Ferretti. "Monitoring ferumoxide-labelled neural progenitor cells and lesion evolution by magnetic resonance imaging in a model of cell transplantation in cerebral ischaemia", F1000Research, 2013 Publication <1 %
-
- 19 Takayasu Ohtake, Shuzo Kobayashi, Shimon Slavin, Yasuhiro Mochida et al. "Human Peripheral Blood Mononuclear Cells Incubated in Vasculogenic Conditioning Medium Dramatically Improve Ischemia/Reperfusion Acute Kidney Injury in Mice", Cell Transplantation, 2018 Publication <1 %
-
- 20 O SANCHEZ. "Family history of early cardiovascular disease: diastolic blood <1 %

pressure as a main target", American Journal of Hypertension, 2004

Publication

21

Panagiotis V Samelis. "Transosseous Suture Fixation of Avulsion Fracture of the Ulnar Collateral Ligament of the Thumb Metacarpophalangeal Joint", Cureus, 2020

Publication

22

www.oncotarget.com

Internet Source

23

Erdal Karaöz, Pinar Cetinalp Demircan, Özlem Sağlam, Ayca Aksoy, Figen Kaymaz, Gökhan Duruksu. "Human dental pulp stem cells demonstrate better neural and epithelial stem cell properties than bone marrow-derived mesenchymal stem cells", Histochemistry and Cell Biology, 2011

Publication

24

Steinbrecher, K.A.. "Increases in Guanylin and Uroguanylin in a Mouse Model of Osmotic Diarrhea Are Guanylate Cyclase C-Independent", Gastroenterology, 200111

Publication

25

Eliana Baez-Jurado, Oscar Hidalgo-Lanussa, Biviana Barrera-Bailón, Amirhossein Sahebkar et al. "Secretome of Mesenchymal Stem Cells and Its Potential Protective Effects on Brain Pathologies", Molecular Neurobiology, 2019

<1 %

<1 %

<1 %

<1 %

<1 %

26

Gilbert, A., A. Wyczalkowska-Tomasik, M. Zendzian-Piotrowska, and B. Czarkowska-Paczek. "Training differentially regulates elastin level and proteolysis in skeletal and heart muscles and aorta in healthy rats", *Biology Open*, 2016.

Publication

<1 %

27

H. Shmilovich. "Circulating endothelial progenitor cells in patients with cardiac syndrome X", *Heart*, 9/1/2007

Publication

<1 %

28

Shin Ishikane, Shunsuke Ohnishi, Kenichi Yamahara, Masaharu Sada et al. "Allogeneic Injection of Fetal Membrane-Derived Mesenchymal Stem Cells Induces Therapeutic Angiogenesis in a Rat Model of Hind Limb Ischemia", *STEM CELLS*, 2008

Publication

<1 %

29

bmcmmedresmethodol.biomedcentral.com

Internet Source

<1 %

30

Dominique Manikowski, Birgit Andrée, Esther Samper, Clémence Saint-Marc et al. "Human adipose tissue-derived stromal cells in combination with exogenous stimuli facilitate three-dimensional network formation of human endothelial cells derived from various sources", *Vascular Pharmacology*, 2018

<1 %

31

Francisco J Azuaje. "Predictive integration of gene functional similarity and co-expression defines treatment response of endothelial progenitor cells", BMC Systems Biology, 2011
Publication

<1 %

32

Xiao Bai, Jie Xi, Yanwen Bi, Xin Zhao, Weidong Bing, Xiangbin Meng, Yimin Liu, Zhonglai Zhu, Guangmin Song. " TNF- α promotes survival and migration of MSCs under oxidative stress NF- κ B pathway to attenuate intimal hyperplasia in vein grafts ", Journal of Cellular and Molecular Medicine, 2017
Publication

<1 %

33

breast-cancer-research.biomedcentral.com
Internet Source

<1 %

34

repository.kulib.kyoto-u.ac.jp
Internet Source

<1 %

35

www.pubfacts.com
Internet Source

<1 %

36

www.spandidos-publications.com
Internet Source

<1 %

37

Evelyn Lara, Nathaly Rivera, Joel Cabezas, Felipe Navarrete, Fernando Saravia, Lleretny Rodríguez-Alvarez, Fidel Castro. "Endometrial Stem Cells in Farm Animals: Potential Role in

<1 %

Uterine Physiology and Pathology", Bioengineering, 2018

Publication

38

Gilang Mauladi Rahman, Muhammad Aminuddin, Budi S Pikir. "LEFT ATRIAL STRAIN BY SPECKLE TRACKING ECHOCARDIOGRAPHY IS NOT PREDICTED FUNCTIONAL CAPACITY IN HYPERTENSIVE POPULATION WITH PRESERVED SYSTOLIC FUNCTION AND NORMAL RESTING LEFT ATRIAL PRESSURE", Indonesian Journal of Cardiology, 2019

Publication

<1 %

39

Jin, Hye, Yun Bae, Miyeon Kim, Soon-Jae Kwon, Hong Jeon, Soo Choi, Seong Kim, Yoon Yang, Wonil Oh, and Jong Chang. "Comparative Analysis of Human Mesenchymal Stem Cells from Bone Marrow, Adipose Tissue, and Umbilical Cord Blood as Sources of Cell Therapy", International Journal of Molecular Sciences, 2013.

Publication

<1 %

40

Mirjam Groger, Hannah S. Fischer, Luzia Veletzky, Albert Lalremruata, Michael Ramharter. "A systematic review of the clinical presentation, treatment and relapse characteristics of human Plasmodium ovale malaria", Malaria Journal, 2017

Publication

<1 %

41	<p>Zhang, Qian, Hong Wang, Yue-Jin Yang, Qiu-Ting Dong, Tian-Jie Wang, Hai-Yan Qian, Na Li, Xi-Mei Wang, and Chen Jin. "Atorvastatin treatment improves the effects of mesenchymal stem cell transplantation on acute myocardial infarction: The role of the RhoA/ROCK/ERK pathway", International Journal of Cardiology, 2014.</p> <p>Publication</p>	<1 %
42	<p>circ.ahajournals.org</p> <p>Internet Source</p>	<1 %
43	<p>repository.unair.ac.id</p> <p>Internet Source</p>	<1 %
44	<p>www.thieme-connect.com</p> <p>Internet Source</p>	<1 %
45	<p>www.translational-medicine.com</p> <p>Internet Source</p>	<1 %
46	<p>Chiang, Kuang-Hsing, Wan-Li Cheng, Chun-Ming Shih, Yi-Wen Lin, Nai-Wen Tsao, Yung-Ta Kao, Chih-Ting Lin, Shinn-Chih Wu, Chun-Yao Huang, and Feng-Yen Lin. "Statins, HMG-CoA Reductase Inhibitors, Improve Neovascularization by Increasing the Expression Density of CXCR4 in Endothelial Progenitor Cells", PLoS ONE, 2015.</p> <p>Publication</p>	<1 %

47

Hui Jing, Xiaomin He, Jinghao Zheng.
"Exosomes and regenerative medicine: state
of the art and perspectives", Translational
Research, 2018

Publication

<1 %

48

MOHSEN EMADEDIN, SHAHEDEH KARIMI,
ALIASGHAR KARIMI, NARGES LABIBZADEH,
MARYAM NIKNEJADI, HOSSEIN BAHARVAND,
NASSER AGHDAMI. "Autologous bone
marrow-derived CD133 cells with core
decompression as a novel treatment method
for femoral head osteonecrosis: a pilot study",
Cytotherapy, 2019

Publication

<1 %

49

PING ZHANG. "Endothelial Differentiation of
Adipose-Derived Stem Cells from Elderly
Patients with Cardiovascular Disease", Stem
Cells and Development, 09/29/2010

Publication

<1 %

50

Sharon Eleuteri, Alessandra Fierabracci.
"Insights into the Secretome of Mesenchymal
Stem Cells and Its Potential Applications",
International Journal of Molecular Sciences,
2019

Publication

<1 %

51

Xiao-Ling Ma, Xiao-Cui Li, Fu-Ju Tian, Si-Ming
Zhang, Xiao-Rui Liu, Yan Zhang, Jian-Xia Fan, Yi
Lin. "Effect of the p53-tristetraprolin-

<1 %

stathmin-1 pathway on trophoblasts at maternal-fetal interface", PLOS ONE, 2017

Publication

52

doaj.org
Internet Source

<1 %

53

proteomesci.biomedcentral.com
Internet Source

<1 %

54

worldwidescience.org
Internet Source

<1 %

55

"Monday, 31 August 2009", European Heart Journal, 09/02/2009
Publication

<1 %

56

Andreas Deckert, Florian Neuhann, Christina Klose, Thomas Bruckner et al. "Assessment of renal function in routine care of people living with HIV on ART in a resource-limited setting in urban Zambia", PLOS ONE, 2017
Publication

<1 %

57

Carl Randall Harrell, Crissy Fellabaum, Nemanja Jovicic, Valentin Djonov, Nebojsa Arsenijevic, Vladislav Volarevic. "Molecular Mechanisms Responsible for Therapeutic Potential of Mesenchymal Stem Cell-Derived Secretome", Cells, 2019
Publication

<1 %

58

Wu, Yangxiao, Li Li, Wen Chen, Wen Zeng, Lingqin Zeng, Can Wen, and Chuhong Zhu.

<1 %

"Maintaining moderate platelet aggregation and improving metabolism of endothelial progenitor cell increase the patency rate of tissue engineered blood vessels", Tissue Engineering Part A, 2015.

Publication

59

Chuma, H.. "One day exposure to FGF-2 was sufficient for the regenerative repair of full-thickness defects of articular cartilage in rabbits", Osteoarthritis and Cartilage, 200410

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Human Umbilical cord blood-mesenchymal stem cell-derived secretome in combination with atorvastatin enhances endothelial progenitor cells proliferation and migration

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10