

Management of Dyslipidemia Associated with Anti-Retroviral Therapy in HIV/AIDS Patients

by Erwin Astha Triyono

Submission date: 06-Dec-2019 04:51PM (UTC+0800)

Submission ID: 1228470587

File name: Associated_with_Anti-Retroviral_Therapy_in_HIVAIDS_Patients.pdf (434.86K)

Word count: 2858

Character count: 17278

Management of Dyslipidemia Associated with Anti-Retroviral Therapy in HIV/AIDS Patients

48 Debrina Kusuma Devi, Erwin Astha Triyono, Musofa Rusli
Division of Tropical Medicine, Department of Internal Medicine, Faculty of Medicine, University of Airlangga, Dr.
Soetomo General Hospital, Jl Prof Dr Moestopo 47 Surabaya 60132, Indonesia
apji@fk.umair.ac.id

47
Keywords: Dyslipidemia, Low-Density Lipoprotein, Human Immunodeficiency Virus, Acquired Immunodeficiency Syndrome, Anti-retroviral

Abstract: Dyslipidemia is a lipid metabolism disorder characterized by increased and decreased lipid fractions in plasma, including elevated total cholesterol, low-density lipoprotein (LDL) cholesterol, TG, and decreased HDL cholesterol. Dyslipidemia often occurs in patients with Human Immunodeficiency Virus/Acquired Immunodeficiency Syndrome (HIV/AIDS), especially those who have received anti-retroviral therapy (ARV). Various ARVs can cause dyslipidemia and the most common is the Protease Inhibitor group. Dyslipidemia is an important risk factor for cardiovascular events. Management of dyslipidemia in HIV-infected patients receiving antiretroviral therapy includes lifestyle modification, switching of antiretroviral drugs, and the use of lipid-lowering drugs as targeted by the recommendations of the NCEP ATP III.

1 INTRODUCTION

18 Human Immunodeficiency Virus (HIV) is a virus that causes Acquired Immunodeficiency Syndrome (AIDS) by attacking white blood cells called CD4 and it can damage the human immune system. According to the CDC (Centers for Disease Control and Prevention), a person is diagnosed with AIDS if it has a CD4 cell count of than 200 cells/mm³ and suffers one or more opportunistic infections or cancer caused by HIV (Departemen Kesehatan RI, 2006).

21 A survey conducted by UNAIDS (Joint United Nations Program on HIV AIDS) stated that there were about 35.3 million people worldwide with HIV and 2.3 million of them were new infections in 2012 (UNAIDS, 2013). Directorate General of Disease Control and Environmental Health-Ministry of Health, Republic of Indonesia has conducted a HIV/AIDS case survey in Indonesia until 2014 that showed 150,296 cases of HIV and 55,799 cases of AIDS with deaths in 9,796 cases (Kementerian Kesehatan RI, 2014).

HIV/AIDS has a high mortality rate; however, the mortality of this disease has dropped dramatically since the use of Highly Active Anti-Retroviral Therapy (HAART). There was a 4%

reduction in the mortality rate of HIV/AIDS from 2000-2001 in the United States and it continued to decline by 70% in 2005. There was a 50% mortality rate from 2003-2007 in HIV/AIDS patients receiving anti-retroviral therapy in Indonesia. The decrease in HIV/AIDS morbidity and mortality significantly improves the quality of life of people with HIV and AIDS. However, long-term adverse effects of antiretroviral therapy (ARV) are emerging, including dyslipidemia, which is a risk factor for cardiovascular disease (HIV-CASUAL Collaboration, 2010; Bavinger et al., 2013).

Dyslipidemia is common in people with HIV who are receiving anti-retroviral therapy. The prevalence of dyslipidemia in HIV patients receiving antiretroviral therapy varies from 30% to 80%, depending on the type of drug used. The most common types are hypertriglyceridemia (40-80%) and hypercholesterolemia (10-50%) (Sprinz et al., 2010).

A study conducted at Dr. Soetomo General Hospital Surabaya, Indonesia in 2009 on 42 HIV patients receiving ARV treatment showed that serum lipid concentration for total cholesterol was 210.5 ± 68.96 mg/dL, triglyceride level of 216.14 ± 134.21 mg/dL, High Density Lipoprotein (HDL) levels of 42.76 ± 12.75 mg/dL, and Low-Density Lipoprotein

(LDL) levels of 144 ± 71.81 mg/dL (Rosita et al., 2009).

Many references correlate dyslipidemia in HIV patients with antiretroviral therapy and its effects on the cardiovascular system. Therefore, we aim to further discuss its management because good management is expected to reduce cardiovascular incidences. This study discussed the management of antiretroviral-related dyslipidemia in HIV/AIDS patients that focused on a selection of lipid-lowering drugs.

2 ANTI-RETROVIRAL THERAPY TYPES

There are six types of ARV drugs, namely:

1. Nucleoside (NRTI) and nucleotide (NtRTI) reverse transcriptase inhibitors: Zidovudine, lamivudine, emtricitabine, stavudine, didanosine, zalcitabine, abacavir, tenovufir.
2. Non-nucleoside reverse transcriptase inhibitor (NNRTI). Efavirenz, nevirapine, delavirdine
3. Protease inhibitor (PI): Amprenavir/ fosamprenavir, atazanavir, indinavir, lopinavir/ ritonavir, nelfinavir, saquinavir, tipranavir, ritonavir
4. Fusion inhibitor (FI): Enfuvirtide
5. Co-receptor antagonist: Maraviroc, vicriviroc.
6. Integrase inhibitor: Raltegravir (Table 1).

The antiretroviral drugs that are available in Indonesia are Lamivudine, Zidovudine, Tenovufir, Stavudine, Emtricitabine, Efavirenz, Nevirapine, Lopinavir, and Ritonavir. Standard antiretroviral therapy at least uses three types of two different drug types. Treatment using three types of drugs is called HAART which aims to suppress HIV viral replication. WHO recommends the use of two NRTI + one NNRTIs as first-line ARV therapy and two NRTI + boosted PIs as the second line. The first-line and second-line ARV combinations recommended in Indonesia are:

1. First line
 - AZT + 3TC + NVP (Zidovudine + Lamivudine + Nevirapine)
 - AZT + 3TC + EFV (Zidovudine + Lamivudine + Efavirenz)
 - TDF + 3TC (or FTC) + NVP (Tenofovir + Lamivudine or Emtricitabine + Nevirapine)
 - TDF + 3TC (or FTC) + EFV (Tenofovir + Lamivudine or Emtricitabine + Efavirenz)
2. Second line
 - TDF + 3TC (or FTC) + LPV/r (Tenofovir + Lamivudine or Emtricitabine + Lopinavir/ritonavir)
 - AZT + 3TC + LPV/r (Zidovudine + Lamivudine + Lopinavir/ritonavir)

Those two lines are a combination therapy commonly used in adolescent and adult patients. Selection of drugs and combinations can be adjusted to the condition of each patient (Kementerian Kesehatan RI, 2011).

Table 1: Available Antiretroviral Therapies Based on Types.

ARV Types		Examples
Nucleoside and Nucleotide Reverse Transcriptase Inhibitors (NRTIs)	Abacavir (ABC) Didanosine (ddI) Zalcitabine (ddC)	Stavudine (d4T) Zidovudine (AZT) Lamivudine (3TC) Tenofovir (TDF) Emtricitabine (FTC)
Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) Protease Inhibitor	Delavirdine (DLV) Etravirine (ETV) Indinavir (IDV) Nelfinavir (NFV) Saquinavir (SQV) Fosamprenavir (FPV) Amprenavir (APV) Tipranavir (TPV) Darunavir (DRV)	Nevirapine (NVP) Efavirenz (EFV) Lopinavir (LPV) Ritonavir (RTV)
Fusion Inhibitor Co-receptor antagonist- CCR5 antagonist	Enfuvirtide Maraviroc	
Integrase Inhibitor	Raltegravir	

Table 2: Therapeutic Effect of ARVs to Lipid Profile

Currently Recommended First-Line Agents are in Bold				
Antiretroviral	Total Cholesterol	LDL-C	HDL-C	Tnglyceride
PLs (boosted)				
Lopinavir	↑↑	↑↑	↔ / ↓	↑↑↑
Atazanavir	↑	↔ / ↑	↔ / ↓	↔
Fosamprenavir	↑	↑	↔ / ↓	↑↑
Saquinavir	↑↑	↑↑	↔ / ↓	↑
Darunavir	↑	↑	↔ / ↓	↑
Tipranavir	↑↑	↑↑	↔ / ↓	↑↑↑
NNRTIs				
Efavirenz	↑	↑	↑	↑
Nevirapine	↑	↑	↑↑	↔ / ↑
NRTIs				
Tenofovir	↔ / ↑	↔ / ↑	↔ / ↑	↔ / ↑
Abacavir	↔ / ↑	↑	↑	↑
Lamivudine	↔	↔	↔	↔
Zidovudine	↑	↑	↑	↑↑
Stavudine	↑↑	↑↑	↑	↑↑
CCR5 Inhibitors				
Maraviroc	↔	↔	↔ / ↑	↔
Integrase Inhibitors				
Raltegravir	↔ / ↑	↔ / ↑	↔ / ↑	↔

Table 3: Category of cardiovascular risk, LDL target, and lipid-lowering drug limits (NCEP ATP III).

Risk Category	LDL-C Goal	Initiate TLC	Consider Drug Therapy
High risk: CHD or CHD risk equivalent (10-years risk > 20%)	< 100 mg/dL (optional goal: < 70 mg/dL)	≥ 100 mg/dL	≥ 100 mg/dL (< 100 mg/dL: consider drug options)
Moderately high risk: 2+ risk (10-year risk 10%)	< 130 mg/dL	≥ 130 mg/dL	≥ 160 mg/dL
Lower risk: 0 – 1 risk factor	< 130 mg/dL	≥ 160 mg/dL	≥ 190 mg/dL (160-189: LDL-lowering drug optional)

CHD: Coronary Heart Disease. Major risk factors: cigarette smoking, hypertension (BP > 140/90 mmHg or on antihypertensive medication); low HDL cholesterol (< 40 mg/dl); family history of premature CHD (CHD in male first-degree relative < 55 years; CHD in female first-degree relative < 65 years, and age (men > 45 years, women > 55 years). TLC: Therapeutic Lifestyle Changes. LDL: Low-Density Lipoprotein (Grundy et al., 2004)

3 DISLIPIDEMIA IN HIV INFECTION

Dyslipidemia is defined as a lipoprotein metabolism abnormality characterized by an increase or decrease in plasma lipoprotein fraction. The major lipoprotein fraction abnormalities are the increase in total cholesterol, LDL cholesterol, triglycerides, and decreased HDL cholesterol. Dyslipidemia is a major risk factor for cardiovascular disease.

Factors that cause dyslipidemia in HIV patients include age, male sex, family history of cardiovascular disease, poor diet, obesity, and poor lifestyle (Aberg and Malvestutto, 2011).

Dyslipidemia in HIV infection is characterized by a decrease in HDL cholesterol, LDL, and total cholesterol, and an increase in TG. Hypertriglyceridemia that occurs in HIV infection is a response to the chronic inflammatory process caused by the HIV virus. The inflammatory mediators involved are interferon- α (IFN- α), interleukin-6 (IL-6) which can increase lipogenesis. There is an increased activity of protein transfer cholesterol ester (CETP) in HIV infection in which CETP plays a role in the transfer of cholesterol ester from HDL to apolipoprotein-B. This explains the reason why HDL cholesterol levels are low in HIV infection.

Dyslipidemia in HIV patients may also be part of the lipodystrophy syndrome. The lipodystrophy syndrome is a cluster of clinical symptoms due to abnormal fat distribution in HIV patients receiving antiretroviral therapy. Lipodystrophy is characterized by elevated triglycerides, total cholesterol, apoprotein B, or hyperinsulinemia. The lipodystrophy syndrome is affected by the length of treatment with ARVs and the age of the patient. This syndrome is characterized by changes in body fat composition in the form of lipoatrophy (decreased subcutaneous fat tissue due to an inability to store peripheral fat with increased distribution to visceral fat) that often occurs in the face, legs and buttocks and lipohypertrophy (accumulation of visceral fat tissue) in the form of gynecomastia and "buffalo hump".

There is an increased release of cytokines in lipodystrophy such as Tumor Necrosis Factor (TNF) and interleukin-6 (IL-6) as well as free fatty acids. Those with a decrease of adipose tissue could lead to lipotoxicity and fat deposition in the liver and muscle. The deposition of fat in the liver causes steatohepatitis and fat deposition in muscles plays a role in metabolic changes, especially insulin resistance. The presence of lipotoxicity also affects pancreatic function and leads to insulin resistance, glucose tolerance disorder, and dyslipidemia. These

metabolic changes increase the risk of the occurrence of cardiovascular disease (Caron et al., 2010; Estrada and Portilla, 2011; Feeney and Patrick, 2011). Detailed informations can be seen on table 1 and figure1.

Several PI mechanisms are involved in the process of dyslipidemia, including:

1. Barriers to proteasome in hepatocytes stimulate TG synthesis and cholesterol.
2. Barriers to Glucose Transporter-4 (GLUT-4) in skeletal muscle and adipocyte tissue result in impaired glucose absorption and lipid metabolic disorders.
3. Barriers to the regulatory sterol translocation process in the process of glucose synthesis and adipocyte differentiation reduce adipocyte differentiation which affects fat distribution and peripheral lipoatrophy.
4. Barriers to SREBP-1 can increase lipogenesis and VLDL resulting in dyslipidemia. Enzyme Binding Protein-1 (SREBP-1), an NRTI class depleting mitochondrial DNA of peripheral adipocyte tissue, accelerates apoptosis and reduces peripheral lipid storage capacity resulting in peripheral lipoatrophy and dyslipidemia. Efavirenz, an NNRTI, can directly induce the occurrence of dyslipidemia (Andrew, 2003) (Figure 2).





Figure 1. Metabolic changes due to lipodystrophy (Caron et al., 2010) VAT = visceral adipose tissue; SAT = subcutaneous adipose tissue; TNF = tumor necrosis factor, IL-6 = interleukin-6; FFAs = free fatty acids; NASH = Non-alcoholic steatohepatitis.



Nature Reviews | Drug Discovery

Figure 2: Pathogenesis of lipodystrophy, dyslipidemia, insulin resistance (Andrew, 2003).




Figure 3: Algorithm for managing dyslipidemia in HIV-infected patients.

	ATV/r	DRV/r	FPV/r	IDV/r	LPV/r	SQV/r	EFV	ETV	NVP	RPV	MVC	DTG	EVG/c	RAL	ABC	FTC	3TC	TDF	ZDV	
Statins																				
Atorvastatin	↑	↑	↑153%	↑	↑490%	↑	↓43%	↓37%	↓	---	---	---	---	---	---	---	---	---	---	---
Fluvastatin	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Lovastatin	↑	↑	↑	↑	↑	↑	↓	↓	↓	---	---	---	---	---	---	---	---	---	---	---
Pravastatin	---	---	↑81%	---	↑	---	↓50%	↓44%	↓	---	---	---	---	---	---	---	---	---	---	---
Rosuvastatin	↑213%	↑48%	↑8%	↑	↑107%	↑	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Simvastatin	↑	↑	↑	↑	↑	↑	↓68%	↓	↓	---	---	---	---	---	---	---	---	---	---	---
Fibrates																				
Bezafibrate	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Clofibrate	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Fenofibrate	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Gemfibrozil	↓	↓	↓	↓	↓	↓41%	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Ezetimibe	↑*	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Colour Legend

- No clinically significant interaction expected.
- These drugs should not be coadministered.
- Potential interaction which may require a dosage adjustment or dose monitoring.
- Potential interaction predicted to be of weak intensity (<2 fdd ↑AUC or <50% ↓AUC). No *a priori* dosage adjustment is recommended.

Text Legend

- ↑ Potential increased exposure of the lipid-lowering drug
- ↓ Potential decreased exposure of the lipid-lowering drug
-

- ↑ Potential increased exposure of HIV drug
- ↓ Potential decreased exposure of HIV drug

* Unboosted atazanavir

Numbers refer to increased or decreased AUC of the lipid-lowering drug as observed in drug-drug interaction studies.

Figure 4: Interaction of ARV with lipid-lowering drugs.

4 DYSLIPIDEMIA MANAGEMENT

The American College of Cardiology and The American Heart Association (ACC/AHA) in 2013 released a recommendation regarding the management of dyslipidemia. However, the recommendations issued by the ACC/AHA do not include the management of dyslipidemia in patients with HIV infection; thus, the management of dyslipidemia in HIV patients continues to use recommendations in accordance with NCEP ATP III (Coffey, 2014; Stone et al., 2013).

Evaluation and management of dyslipidemia in HIV-infected patients receiving antiretroviral therapy include lifestyle modification, switching of antiretroviral drugs, and the use of lipid-lowering drugs as targeted by the recommendations of NCEP ATP III. The examination of total cholesterol, LDL, HDL, and TG should be checked before starting antiretroviral therapy and should be evaluated periodically. Lifestyle modifications such as

smoking cessation, diet, and exercise are the initial recommendations. If the lipid target is still not reached then it is necessary to replace antiretroviral drugs or use lipid-lowering drugs (Estrada and Portilla, 2011; Dube et al., 2003; Grundy et al., 2004) (Table 3).

The use of lipid-lowering drugs may interact pharmacokinetically with antiretroviral drugs; thus, the selection of lipid-lowering drugs is important. Low-dose atorvastatin and rosuvastatin may be used with protease inhibitors with monitoring of side-effects. If there are no side-effects and lipid levels have not reached the target after 6-8 weeks of usage, then the doses of atorvastatin and rosuvastatin may be increased. Use of simvastatin and lovastatin is not recommended when combined with protease inhibitors. Pravastatin levels decrease by about 50% when used with protease inhibitors; thus, it is necessary to increase the dose to improve the efficacy.

Atorvastatin, lovastatin, pravastatin, and simvastatin levels decrease when used with NNRTIs

(EFV and NVP). Ezetimibe can reduce LDL levels by 10-20% and it has synergistic effects when combined with a statin class. Fibrates (gemfibrozil) or fenofibrate is useful for lowering TG levels. Fibrate can reduce TG levels by 50% and increase HDL by 5-10%.

Fibrates generally have little interaction with protease inhibitors because the fibrates are not metabolized in the cytochrome P450. Fibrates can be combined with statins to increase the effect on decreased TG up to normal levels. Fenofibrate can reduce TG levels by more than 50%. If combined with pravastatin, it can achieve lipid targets according to NCEP ATP III in patients with HIV (Estrada and Portilla, 2011; Calza et al., 2003; Wohl et al., 2008) (Figure 3 & 4).

5 PROGNOSIS

The chronic inflammatory process in HIV patients plays a role in increasing the risk of coronary artery disease. The incidence of acute myocardial infarction is quite high in HIV patients, both ARV and non-ARV. The risk of acute myocardial infarction associated with dyslipidemia in HIV patients receiving PI therapy increases by 2.13 times, NRTI increases by 1.21 times in Stavudine use, the use of Lamivudine and Zidovudine increases by 1.05 times, and the NNRTI group of Efavirenz increases by 1.02 times compared to HIV patients that do not receive antiretroviral therapy (Bavinger et al., 2013).

6 CONCLUSION

HIV/AIDS has a high mortality rate; however, the mortality of this disease has dropped dramatically since the use of Highly Active Anti-Retroviral Therapy (HAART). Dyslipidemia is common in people with HIV who are receiving anti-retroviral therapy. The incidence of acute myocardial infarction is quite high in HIV patients, both ARV and non-ARV.

REFERENCES

Aberg, J., & Malvestutto, C. D. 2011. Management Of Dyslipidemia In Hiv-Infected Patients. *Clin Lipidol*, 6, 447-462.

- Andrew, C. 2003. . Toxicity Of Antiretroviral Therapy And Implications For Drug Development. . *Nature Reviews Drug Discovery*, 2, : 624-634.
- Bavinger, C., , Bendavid, E., Niehaus, K., Olshen, R. A., . & Olkin, I. 2013. Risk Of Cardiovascular Disease From Antiretroviral Therapy For Hiv: . *A Systematic Review. Plos One*, 8.
- Calza, L., Manfredi, R. & Chiodo, F. 2003. . Statins And Fibrates For The Treatment Of Hyperlipidaemia In Hiv-Infected Patients Receiving Haart. *Aids*, 17, : 851-9.
- Caron, D., M., , Lagathu C. & Boccara, F. 2010. . Hiv-Associated Lipodystrophy: From Fat Injury To Premature Aging. *Trends Mol Med*, 16, :218-229.
- Coffey, S., , 2014. 2014. Dyslipidemia. . *In Guide For Hiv/Aids Clinical Care. Rockville: U.S.Department Of Health And Human Services*, 339-349.
- Departemen Kesehatan Ri 2006. . Situasi Hiv/Aids Di Indonesia. . Jakarta: : Pusat Data Dan Informasi Departemen Kesehatan Ri.
- Dube, M. P., Stein, J. H., Aberg, J. A., Fichtenbaum, C. J. & Gerber, J. G. 2003. . Guidelines For The Evaluation And Management Of Dyslipidemia In Hiv-Infected Adults Receiving Antiretroviral Therapy: Recommendation Of The Hiv Medical Association Of The Infectious Disease Society Of America And The Adult Aids Clinical Trials Group. *Clin Infect Dis*, 37:613-627.
- Estrada, C. & Portilla, J. 2011. . Dyslipidemia Related Antiretroviral Therapy. . *Aids Rev*, 13, 49-56. .
- Feeney, E. R. & Patrick, W. 2011. Hiv And Haart-Associated Dyslipidemia. . *Cardiovasc Med J*, 49-63.
- Grundy, S., J., C., Merz, C., Brewer, H. B. & Clark, L. T., . 2004. Implications Of Recent Clinical Trials For The National Cholesterol Education Program Adult Treatment Panel Iii Guidelines. . *Circulation*, 110, 227-239.
- Hiv-Casual Collaboration 2010. . The Effect Of Combined Antiretroviral Therapy Overall Mortality In Hiv-Infected Individuals. *Aids*, 24, : 123-37.
- Kementerian Kesehatan Ri 2011. Pedomn Nasional Tatalaksana Klinis Infeksi Hiv Dan Terapi Antiretroviral Pada Orang Dewasa Dan Remaja. . Jakarta: Kementerian Kesehatan Ri.
- Kementerian Kesehatan Ri 2014. . Statistik Kasus Hiv/Aids Di Indonesia Dilapor S/D September 2014. . Sumber : Direktorat Jenderal Pengendalian Penyakit Dan Penyehatan Lingkungan Kementerian Kesehatan Republik Indonesia.
- Rosita, C., Hans, L. & Desy, H., 2009. . Profil Lipid Penderita Hiv/Aids Yang Mendapat Pengobatan Arv. . *Media Jurnal Berkala Ilmu Kesehatan Kulit&Kelamin*, Vol 21.
- Sprinz, E., , Rosmeri, K. & Regina, K. 2010. . Dyslipidemia In Hiv-Infected Individuals. . *Braz J Infect Dis*, 14.
- Stone, N. J., Robinson, J., Lichtenstein, A. H., Bairey, M. & Blum, J. B. 2013. . Acc/Aha Guideline On The Treatment Of Blood Cholesterol To Reduce Atherosclerotic Cardiovascular Risk In Adults: A

- Report Of The American College Of Cardiology/American Heart Association Task Force On Practice Guidelines. . *J Am Coll Cardiol* 2013 Nov 7.
- Unaid 2013. . Global Report Unaid Report On The Global Aids Epidemic 2013. Switzerland:Unaid.
- Wohl, D. A., Waters, D. & Simpson, R. J. 2008. . Ezetimibe Alone Reduces Low-Density Lipoprotein Cholesterol In Hiv-Infected Patients Receiving Combination Antiretroviral Therapy. . *Clin Infect Dis*, 47: , 1105-8.



Management of Dyslipidemia Associated with Anti-Retroviral Therapy in HIV/AIDS Patients

ORIGINALITY REPORT

29%

SIMILARITY INDEX

20%

INTERNET SOURCES

27%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1

www.pitt.edu

Internet Source

2%

2

Jeroen P. H. van Wijk, Manuel Castro Cabezas. "Hypertriglyceridemia, Metabolic Syndrome, and Cardiovascular Disease in HIV-Infected Patients: Effects of Antiretroviral Therapy and Adipose Tissue Distribution", International Journal of Vascular Medicine, 2012

Publication

1%

3

www.tachc.org

Internet Source

1%

4

M. Tungsiripat. "Dyslipidemia in HIV patients.", Cleveland Clinic Journal of Medicine, 12/01/2005

Publication

1%

5

T Marie Rossouw, M E Botes, F Conradie. "Overview of HIV-related lipodystrophy", Southern African Journal of HIV Medicine, 2013

Publication

1%

6

Barbara J. Zarowitz, Wilbert S. Aronow, Kelly Hollenack, Terry O'Shea. "The Application of Evidence-Based Principles of Care in Older Persons (Issue 2): Management of Lipid Disorders", Journal of the American Medical Directors Association, 2006

Publication

1%

7

avpivnik.ru

Internet Source

1%

8

r-libre.telug.ca

Internet Source

1%

9

www.pharmacy.umn.edu

Internet Source

1%

10

Edna Maria Vissoci Reiche, Andrea Name Colado Simao. "Chapter 2 The Role of Human Immunodeficiency Virus Type 1 (HIV-1) Proteins and Antiretroviral Drug Therapy in HIV-1-Induced Oxidative Stress", InTech, 2011

Publication

1%

11

Lamarca, Karuna, Ana García Sarasola, Francesc Vidal, and Pere Domingo. "Drug therapies for HIV-related metabolic disorders", Expert Opinion on Pharmacotherapy, 2016.

Publication

1%

12

www.kebijakanaidssindonesia.net

Internet Source

1%

13

J. Phua. "Management of severe sepsis in patients admitted to Asian intensive care units: prospective cohort study", BMJ, 06/13/2011

Publication

1%

14

www.lexi.com

Internet Source

1%

15

Domingos, Hamilton, Rivaldo Venãçncio da Cunha, Anamaria Mello Miranda Paniago, Diego Mira Martins, Eduardo Brandã£o Elkhoury, and Albert Schiaveto de Souza. "Metabolic effects associated to the highly active antiretroviral therapy (HAART) in AIDS patients", Brazilian Journal of Infectious Diseases, 2009.

Publication

1%

16

Tochi Okwuosa. "HIV with Dyslipidemia", Hyperlipidemia in Primary Care, 2011

Publication

1%

17

P Widyaningsih, U U Zahra, V Y Kurniawan, Sutanto, D R S Saputro. "Susceptible infected AIDS treatment (SIAT) model", IOP Conference Series: Earth and Environmental Science, 2019

Publication

1%

18

www.completewomencare.com

Internet Source

1%

19

S B Gumede, A Fischer, W D F Venter, S T Lalla-Edward. "Descriptive analysis of World

1%

Health Organization-recommended second-line antiretroviral treatment: A retrospective cohort data analysis", South African Medical Journal, 2019

Publication

20

Tegene Legese Dadi, Adane Teshome Kefale, Teshale Ayele Mega, Muktar Sano Kedir, Habtamu Acho Addo, Tessema Tsehay Biru. "Efficacy and Tolerability of Tenofovir Disoproxil Fumarate Based Regimen as Compared to Zidovudine Based Regimens: A Systematic Review and Meta-Analysis", AIDS Research and Treatment, 2017

Publication

<1%

21

S Bakri, H Rasyid, H Kasim, S Katu. "Chronic kidney disease in HIV patients", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1%

22

editorial.glowm.com

Internet Source

<1%

23

www.ncbi.nlm.nih.gov

Internet Source

<1%

24

www.jmir.org

Internet Source

<1%

25

jamanetwork.com

Internet Source

<1%

- | | | |
|----|---|-----|
| 26 | C MERRY. "Pharmacology of Antiretroviral Drugs", Global HIV/AIDS Medicine, 2008
Publication | <1% |
| 27 | Teresa M. Seccia, Gian Paolo Rossi. "Clinical Use and Pathogenetic Basis of Laboratory Tests for the Evaluation of Primary Arterial Hypertension", Critical Reviews in Clinical Laboratory Sciences, 2008
Publication | <1% |
| 28 | Panjaitan, Epi, Satiti Retno Pudjiati, and Agnes Sri Siswati. "Low CD4+ T cell counts are not risk factor for Malassezia species infection in HIV/AIDS patients", Journal of the Medical Sciences (Berkala Ilmu Kedokteran), 2015.
Publication | <1% |
| 29 | WK Tim Wong, Jane M Ussher. "Life with HIV and AIDS in the Era of Effective Treatments: 'It's Not Just about Living Longer!'", Social Theory & Health, 2008
Publication | <1% |
| 30 | www.nejm.org
Internet Source | <1% |
| 31 | www.sajhivmed.org.za
Internet Source | <1% |
| 32 | Dewi Kurnia Putri, Hema Malini, Basmanelly Basmanelly. "Living with HIV/AIDS: Stigma and | <1% |

Social Discrimination", Jurnal The Messenger, 2019

Publication

33

pure.uva.nl

Internet Source

<1%

34

Anna Coutsooudis, Leith Kwaan, Mairi Thomson. "Prevention of vertical transmission of HIV-1 in resource-limited settings", Expert Review of Anti-infective Therapy, 2014

Publication

<1%

35

www.thebodypro.com

Internet Source

<1%

36

Madhu N. Rao. "HIV Infection and Diabetes", Principles of Diabetes Mellitus, 2010

Publication

<1%

37

www.dovepress.com

Internet Source

<1%

38

Leonardo Calza, Roberto Manfredi, Daria Pocaterra, Francesco Chiodo. "Risk of premature atherosclerosis and ischemic heart disease associated with HIV infection and antiretroviral therapy", Journal of Infection, 2008

Publication

<1%

39

William Wester, C., Sophie Limou, and Cheryl A. Winkler. "HIV Pharmacogenetics and Pharmacogenomics", Genomic and

<1%

Personalized Medicine, 2013.

Publication

40

"Minutes of The 43rd General Assembly of The European Association for The Study of Diabetes", Diabetologia, 2008

Publication

<1%

41

Physiology and Physiopathology of Adipose Tissue, 2013.

Publication

<1%

42

Marcelo C. Batiste, Ernst J. Schaefer. "Diagnosis and Management of Lipoprotein Abnormalities", Nutrition in Clinical Care, 2002

Publication

<1%

43

discovery.ucl.ac.uk

Internet Source

<1%

44

Adriana Cadilla, Nadia Qureshi, Daniel C Johnson. "Pediatric antiretroviral therapy", Expert Review of Anti-infective Therapy, 2014

Publication

<1%

45

www.yumpu.com

Internet Source

<1%

46

"Comprehensive Cardiovascular Medicine in the Primary Care Setting", Springer Science and Business Media LLC, 2011

Publication

<1%

47

ir.jkuat.ac.ke

Internet Source

<1%

48

jurnal.ugm.ac.id

Internet Source

<1%

49

Willard, S.. "The Nurse Practitioner's Role in Managing Dyslipidemia and Other Cardiovascular Risk Factors in HIV-Infected Patients: Impact of Antiretroviral Therapy", *Journal of the Association of Nurses in AIDS Care*, 200601/02

Publication

<1%

50

C BAIREYMERZ. "Treatment Guidelines Overview", *Clinical Lipidology*, 2009

Publication

<1%

51

Julian Falutz. "Therapy Insight: body-shape changes and metabolic complications associated with HIV and highly active antiretroviral therapy", *Nature Clinical Practice Endocrinology & Metabolism*, 09/2007

Publication

<1%

52

LECLERCQ, Pascale, Cecile M GOUJARD, Martin Duracinsky, Francois Andre ALLAERT, Marianne L'HENAFF, Maeva HELLET, Jean Pierre MEUNIER, Sophie CARRET, Jacques THEVENON, Philippe NGO VAN, and Gilles PIALoux. "High prevalence and impact on the quality of life of facial lipoatrophy and other

<1%

abnormalities in fat tissue distribution in HIV-infected patients treated with antiretroviral therapy", AIDS Research and Human Retroviruses, 2012.

Publication

53

Leonardo Calza. "Myocardial infarction risk in HIV-infected patients: epidemiology, pathogenesis, and clinical management :", AIDS, 03/2010

Publication

<1%

54

"Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism", Wiley, 2018

Publication

<1%

55

Scott M. Hammer. "A Controlled Trial of Two Nucleoside Analogues plus Indinavir in Persons with Human Immunodeficiency Virus Infection and CD4 Cell Counts of 200 per Cubic Millimeter or Less", New England Journal of Medicine, 09/11/1997

Publication

<1%

56

HISPALIPID Study Group Investigators. "The gap between dyslipidemia control perceived by physicians and objective control patterns in Spain", Atherosclerosis, 200610

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Management of Dyslipidemia Associated with Anti-Retroviral Therapy in HIV/AIDS Patients

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
