

DAFTAR PUSTAKA

- American Diabetes Association (2014) ‘Diagnosis and classification of diabetes mellitus’, *Diabetes Care*, 37(SUPPL.1), pp. 81–90. doi: 10.2337/dc14-S081.
- Aminoff, M., Simon, R. and Greenberg, D. (2018) *Clinical Neurology*. United State of America: McGraw-Hill Education.
- Andersen, K. K. et al. (2009) ‘Hemorrhagic and ischemic strokes compared: Stroke severity, mortality, and risk factors’, *Stroke*, 40(6), pp. 2068–2072. doi: 10.1161/STROKEAHA.108.540112.
- Arsana, P. M. et al. (2015) ‘Panduan pengelolaan dislipidemia di Indonesia’, *Pb. Perkeni*, p. 4. doi: 10.1002/bit.22430.
- Badan Pusat Statistik (2017) ‘Potret Pendidikan Indonesia Tahun 2017’, in. Available at: <https://www.bps.go.id/publication/download.html?nrbvfeve=ETY1YjUyNmMxMTIjZThmNzk5ZTV1YTYz&xzmn=aHR0cHM6Ly93d3cuYnBzLmdvLmlkL3B1YmxpY2F0aW9uLzIwMTgvMTIvMDYvPTY1YjUyNmMxMTIjZThmNzk5ZTV1YTYzL3N0YXRpc3Rpay1wZW5kaWRpa2FuLTIwMTguaHRtbA%3D%3D&twoadfnoarfeauf=MjA>.
- Badan Pusat Statistik (2018) ‘Penduduk 15 Tahun Ke Atas Menurut Status Pekerjaan Utama 1986 – 2018’. Available at: <https://www.bps.go.id/statictable/2009/04/16/971/penduduk-15-tahun-ke-atas-menurut-status-pekerjaan-utama-1986--2018.html>.
- Baehr, M. and Frotscher, M. (2012) *Duis' Topical Diagnosis in Neurology*. 5th edn. Germany: Thieme Publishing Group.
- Batubara, S. and Putri, C. (2018) ‘Jurnal Kedokteran STM (Sains dan Teknologi Medik) Januari – Juni 2018 P - ISSN 2614 – 610X E – ISSN 2614 - 8218 PERBEDAAN PENURUNAN FUNGSI KOGNITIF PADA PASIEN STROKE ANTARA LESI HEMISFER KANAN DAN KIRI DI POLIKLINIK SARAF RSUD EMBUNG FATIMAH KOTA BATAM’, pp. 16–21.
- Bozluolcay, M. et al. (2013) ‘Hypercholesterolemia as one of the risk factors of intracerebral hemorrhage’, *Acta Neurologica Belgica*, 113(4), pp. 459–462. doi: 10.1007/s13760-013-0222-6.
- Brito-Marques, P. R. de and Cabral-Filho, J. E. (2005) ‘Influence of age and schooling on the performance in a modified Mini-Mental State Examination version: a study in Brazil Northeast’, *Arquivos de Neuro-Psiquiatria. Associação Arquivos de Neuro-Psiquiatria*, 63(3a), pp. 583–587. doi: 10.1590/S0004-282X2005000400005.
- Campbell, W. (2005) *DeJong's The Neurologic Examination*. 6th edn. Philadelphia: Lippincott, Williams, & Wilkins.
- Chopra, H. K. and Ram, C. V. S. (2019) ‘Recent Guidelines for Hypertension: A Clarion Call for Blood Pressure Control in India’, *Circulation Research*, 124(7), pp. 984–986. doi: 10.1161/CIRCRESAHA.119.314789.
- Erkinjuntti, T. and Gauthier, S. (2009) ‘The concept of vascular cognitive impairment’, *Frontiers of Neurology and Neuroscience*, 24(Mid), pp. 79–85. doi: 10.1159/000197886.
- Ferreira, M. G. R., Moro, C. H. C. and Franco, S. C. (2015) ‘Cognitive performance after ischaemic stroke’, *Dementia e Neuropsychologia*, 9(2), pp. 165–175. doi: 10.1590/1980-57642015DN92000011.
- Ghani, L., Mihardja, L. K. and Delima, D. (2016) ‘Faktor Risiko Dominan Penderita

- Stroke di Indonesia', *Buletin Penelitian Kesehatan*, 44(1), pp. 49–58. doi: 10.22435/bpk.v44i1.4949.49–58.
- Hedna, V. S. et al. (2013) 'Hemispheric differences in ischemic stroke: Is left-hemisphere stroke more common?', *Journal of Clinical Neurology (Korea)*, 9(2), pp. 97–102. doi: 10.3988/jcn.2013.9.2.97.
- Hewitt, J. et al. (2012) 'Diabetes and stroke prevention: A review', *Stroke Research and Treatment*, 2012. doi: 10.1155/2012/673187.
- Hurn, P. D. and Brass, L. M. (2003) 'Estrogen and stroke: A balanced analysis', *Stroke*, 34(2), pp. 338–341. doi: 10.1161/01.STR.0000054051.88378.25.
- Kabi, G. Y. C. R., Tumewah, R. and Kembuan, M. A. H. N. (2015) 'Gambaran Faktor Risiko Pada Penderita Stroke Iskemik Yang Dirawat Inap Neurologi Rsup Prof. Dr. R. D. Kandou Manado Periode Juli 2012 - Juni 2013', *e-CliniC*, 3(1), pp. 1–6. doi: 10.35790/ecl.3.1.2015.7404.
- Khedr, E. M. et al. (2009) 'Cognitive impairment after cerebrovascular stroke: Relationship to vascular risk factors', *Neuropsychiatric Disease and Treatment*, 5(1), pp. 103–116. doi: 10.2147/ndt.s4184.
- Kim, M. and Park, J. M. (2017) 'Factors affecting cognitive function according to gender in community-dwelling elderly individuals', *Epidemiology and health*, 39, p. e2017054. doi: 10.4178/epih.e2017054.
- Krack, P. et al. (2010) 'Deep brain stimulation: From neurology to psychiatry?', *Trends in Neurosciences*. Elsevier Ltd, 33(10), pp. 474–484. doi: 10.1016/j.tins.2010.07.002.
- Lawrence, E. S. et al. (2001) 'Estimates of the prevalence of acute stroke impairments and disability in a multiethnic population', *Stroke*, 32(6), pp. 1279–1284. doi: 10.1161/01.STR.32.6.1279.
- Leisman, G., Braun-Benjamin, O. and Melillo, R. (2014) 'Cognitive-motor interactions of the basal ganglia in development', *Frontiers in Systems Neuroscience*, 8(FEB), pp. 1–18. doi: 10.3389/fnsys.2014.00016.
- Mattsson, N. et al. (2016) 'Selective vulnerability in neurodegeneration: Insights from clinical variants of Alzheimer's disease', *Journal of Neurology, Neurosurgery and Psychiatry*, 87(9), pp. 1000–1004. doi: 10.1136/jnnp-2015-311321.
- Mitrovic, I. (2014) *Pathophysiology of Disease: An Introduction to Clinical Medicine*. 7th edn. United State of America: McGraw-Hill Education.
- Mohd Zulkifly, M. F. et al. (2016) 'A Review of Risk Factors for Cognitive Impairment in Stroke Survivors', *Scientific World Journal*, 2016. doi: 10.1155/2016/3456943.
- Munsch, F. et al. (2016) 'Stroke location is an independent predictor of cognitive outcome', *Stroke*, 47(1), pp. 66–73. doi: 10.1161/STROKEAHA.115.011242.
- Murphy, S. J., McCullough, L. D. and Smith, J. M. (2004) 'Stroke in the female: role of biological sex and estrogen.', *ILAR journal / National Research Council, Institute of Laboratory Animal Resources*, 45(2), pp. 147–159. doi: 10.1093/ilar.45.2.147.
- Naço, D. et al. (2013) 'Factors influencing mini-mental state (MMSE) score in stroke patients.', *Medicinski arhiv*, 67(3), pp. 171–173. doi: 10.5455/medarh.2013.67.171-173.
- Opitz, B. (2014) 'Memory function and the hippocampus', *The Hippocampus in Clinical Neuroscience*, 34, pp. 51–59. doi: 10.1159/000356422.
- Park, J. et al. (2016) 'The impact of acute phase domain-specific cognitive function

- on post-stroke functional recovery', *Annals of Rehabilitation Medicine*, 40(2), pp. 214–222. doi: 10.5535/arm.2016.40.2.214.
- Portegies, M. L. P. et al. (2015) 'Left-sided strokes are more often recognized than right-sided strokes: The rotterdam study', *Stroke*, 46(1), pp. 252–254. doi: 10.1161/STROKEAHA.114.007385.
- Putri, M., Mutiawati, E. and Mahdani, W. (2017) 'Hubungan Derajat Stroke Terhadap Status Kognitif Pada Pasien Stroke Iskemik Di Poliklinik Saraf Rumah Sakit Umum Daerah dr . Zainoel Abidin Banda Aceh Relationship Degree Stroke on The Cognitive Status Patients Ischemic Stroke', 2, pp. 61–67.
- Ren, L. et al. (2018) 'Investigation of the prevalence of Cognitive Impairment and its risk factors within the elderly population in Shanghai, China', *Scientific Reports*. Springer US, 8(1), pp. 1–9. doi: 10.1038/s41598-018-21983-w.
- Rowland, D. T. (2012) *Aging in Asia, Population Aging*. doi: 10.1007/978-94-007-4050-1_14.
- Sacco, R. L. et al. (2013) 'An updated definition of stroke for the 21st century: A statement for healthcare professionals from the American heart association/American stroke association', *Stroke*, 44(7), pp. 2064–2089. doi: 10.1161/STR.0b013e318296aeca.
- Schmidt, R. et al. (1993) 'Cognitive impairment after acute supratentorial stroke: a 6-month follow-up clinical and computed tomographic study', *European Archives of Psychiatry and Clinical Neuroscience*, 243(1), pp. 11–15. doi: 10.1007/BF02191518.
- Tan, C. S. et al. (2015) 'Trends in stroke incidence and 28-day case fatality in a nationwide stroke registry of a multiethnic Asian population', *Stroke*, 46(10), pp. 2728–2734. doi: 10.1161/STROKEAHA.115.009797.
- Teasell, R. et al. (2019) 'Chapter 12 : Cognitive Rehabilitation', *Evidence-Based Review Stroke Rehabilitation*, (2000).
- Thrift, A. G. et al. (2017) 'Global stroke statistics', *International Journal of Stroke*, 12(1), pp. 13–32. doi: 10.1177/1747493016676285.
- Toga, A. W. and Thompson, P. M. (2003) 'Mapping brain asymmetry', *Nature Reviews Neuroscience*, 4(1), pp. 37–48. doi: 10.1038/nrn1009.
- Tziomalos, K. et al. (2009) 'Dyslipidemia as a Risk Factor for Ischemic Stroke', *Current Topics in Medicinal Chemistry*, 9(14), pp. 1291–1297. doi: 10.2174/156802609789869628.
- Voos, M. and Ribeiro do Valle, L. (2008) 'Comparative study on the relationship between stroke hemisphere and functional evolution in right-handed individuals', *Brazilian Journal of Physical Therapy*, 12(2), pp. 222–224.
- Wolff, M. and Vann, S. D. (2019) 'The cognitive thalamus as a gateway to mental representations', *Journal of Neuroscience*, 39(1), pp. 3–14. doi: 10.1523/JNEUROSCI.0479-18.2018.
- World Health Organization (2018) *Disease burden and mortality estimates*. Available at: https://www.who.int/healthinfo/%0Aglobal_burden_disease/estimates/en/index1.html%0A.
- Wray (2017) 'Stroke and Chronic Kidney Disease: Epidemiology, Pathogenesis, and Management Across Kidney Disease Stages', *Physiology & behavior*, 176(5), pp. 139–148. doi: 10.1016/j.physbeh.2017.03.040.
- Yogarajah, M. (2015) *Crash Course Neurology*. 4th edn. Elsevier.
- Yoon, J. A. et al. (2017) 'Factors associated with improvement or decline in cognitive function after an ischemic stroke in Korea: The Korean stroke cohort for

- functioning and rehabilitation (KOSCO) study', *BMC Neurology*. BMC Neurology, 17(1). doi: 10.1186/s12883-016-0780-3.
- Zhao, L. *et al.* (2018) 'Strategic infarct location for post-stroke cognitive impairment: A multivariate lesion-symptom mapping study', *Journal of Cerebral Blood Flow and Metabolism*, 38(8), pp. 1299–1311. doi: 10.1177/0271678X17728162.