# FOLIA MEDICA INDONESIA

Vol. 48 No. 1 January - March 2012

AND THE RESIDENCE OF THE PARTY OF THE PARTY

A REAL PROPERTY AND A PROPERTY OF THE PROPERTY

State and details and applicable or enterproper I not be compared to the compa THE RESERVE AND DESCRIPTION OF THE PERSON NAMED IN COLUMN TWO

Contraction of the second seco

Market Support for the State of the State of

AND AND THE RESIDENCE OF A SECOND PROPERTY OF A SEC

BETWEEN BOTH OF THE STATE OF TH

Wildler I I WAR THE REAL PROPERTY AND ADDRESS OF THE PARTY AND ADDRESS 

> terrorio detecti Dellara Cadrillo des la Praco da P are any househouse

UNIVERSITAS AIRLANGGA Unair e-journal PPJPI

# Journal Unair

MEDICINE journal category

SCIENCES journal category **ECONOMY** journal category journal category

ANIMAL-AGRICULTURE journal category

PHARMACY HEALTH journal category journal category

PSYCHOLOGY journal category


SOCIAL HUMANISM journal category











# Information Folia Medica Indonesiana

# **EDITORIAL BOARD**

### **Editor-in Chief**

Prof Dr Suhartono Taat Putra, dr, MS

# **Deputy Editor-in-Chief**

I Dewa Gede Ugrasena, dr, SpA(K)

# **Editorial Board**

Prof Bambang Permono, dr, SpA(K)

Prof Purnomo Suryohudoyo, dr

Prof Dr Doddy M Soebadi, dr, SpB, SpU(K)

Prof Soehartono DS, dr, SpOG(K)

Prof Dr Harjanto, dr, AIF

Dr FM Judajana, dr, SpPK

Prof Soetjipto, dr, MS, PhD

Prof Dr Suharto, dr, MSc, DTM&H, SpPD-KTI

Prof Dr Subijanto Marto Soedarmo, dr, SpA(K)

Prof Dr Kuntaman, dr, MS, SpMK

Dr Anang Endaryanto, dr, SpA(K)

Edhi Rianto, dr, MS

# **Managing Editor**

Dr Ahmad Suryawan, dr, SpA(K)

Dr I Ketut Sudiana, drs, MSi

Dr Nyilo Purnami, dr, SpTHT-KL(K)

Viskasari Pintoko Kalanjati, dr, MKes, PhD

Purwo Sri Rejeki, dr, MKes

Dr Gondo Mastutik, drh, MKes

Mia Ratwita Andarsini, dr, SpA

Wita Saraswati, dr, SpOG

Dr Ahmad Yudianto, dr, SpF

Linda Dewanti, dr, MKes, HMsc, PhD

Mochammad Zuhdy, S.Sos

Fitriya Niken Ariningsih, S.Sos

**Cover Media** FOLIA MEDICA INDONESIANA

Volume: 48 / No.: 1 / Pub.

2012-01

Content

- 1. Effect Of Sodium Nitrite (nano2) To Erithrocyte And Hemoglobin Profile In White Rat (rattus Norvegicus)
- 2. Synthesis Of Beta Fertilin Protein Polyclonal Antibody Of Human Sperm Membrane As A
- Candidate For Immunocontraceptive Material 3. Long-term Glucocorticoids Effects Of Bone Lining Cells Apoptosis
- 4. Zinc Sulfate Increases Lymphocyte Cd4 Count In Hiv/aids Patients At Icuid Dr. Soetomo Hospital Surabaya
- 5. Support Group Effect On Cd4 Levels In Plwha Who Recieved Art In Upipi Rsud Dr. Soetomo Surabaya
- 6. Intensity Of Swimming Exercise As Chronic Stress Induction Towards Expression Of Glucose Transporter 1 (glut1) In Brain Capillary Endothelial Of Rats (rattus Norvegicus)
- $\hbox{7. The Training Of Wiggler Monitoring Of Primary School Teachers And Students In Surabaya}\\$ To Improve Eradication Of Dengue Hemorraghic Fever
- 8. Implementation Of Safety Work For Nurse In Handling Hiv/aids Patients
- 9. Case Report: Posterior Interosseous Nerve Compression Syndrome

# Information FMI

# About FMI History **Editorial Board** Note for authors Address

# e-journal Unair



Continue »

# Licensed

All of the journals are published by Universitas Airlangga that free to access and download the articles.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

# Journal Account Users

Please enter your email GO Enter Email Here.. Saya bukan robot

reCAPTCHA

# Other Link

Official Web Unair E-journal Repository Diailib Official web PPJPI

# Statistic



# About Journal

Vision & Mission, Goals

# INTENSITY OF SWIMMING EXERCISE AS CHRONIC STRESS INDUCTION TOWARDS EXPRESSION OF GLUCOSE TRANSPORTER 1 (GLUT1) IN BRAIN CAPILLARY ENDOTHELIAL OF RATS (*Rattus norvegicus*)

Hayuris Kinandita, Lilik Herawati, Harlina

Department of Medical Physiology Faculty of Medicine, Airlangga University

# **ABSTRAK**

Stres dapat menyebabkan efek positif atau negatif pada tubuh, tergantung pada durasi dan intensitas stres. Stres berkepanjangan akan menyebabkan penyakit neurodegeneratif atau gangguan kognitif melalui gangguan dalam sistem neurotransmitter otak, tingkat biomolekuler otak, dan metabolisme otak. Regulasi potensial Glucose Transporter-1 (GLUT1) dalam respon sawar darah-otak terhadap berbagai kondisi patologis otak yang berhubungan dengan stres. Tujuan penelitian ini adalah untuk menjelaskan pengaruh intensitas stres kronis terhadap ekspresi transporter-1 glukosa pada otak tikus. Penelitian ini menggunakan 30 ekor tikus (Rattus norvegicus) galur wistar jantan yang dibagi menjadi 3 kelompok: kelompok kontrol (K0), perlakuan latihan berenang dalam intensitas berat (K1) dan latihan berenang dalam intensitas ringan (K2). Pemeriksaan dilakukan terhadap ekspresi GLUT1 pada sel endotel otak yang diamati menggunakan teknik pewarnaan imunohistokimia. Hasil penelitian menunjukkan bahwa ada penurunan ekspresi GLUT1 di endotel otak akibat latihan renang intensitas berat dan latihan renang intensitas ringan. Deplesi ekspresi GLUT1 dalam endotel otak kelompok intensitas berat lebih besar dari pada kelompok intensitas ringan dengan p <0,05. (FMI 2012;48:24-27)

Kata kunci: GLUT1 (glucose transporter-1), stres kronik, otak

# ABSTRACT

Stress can lead positive or negative effect on the body, depending on the duration and intensity of stress. Prolonged stress will cause neurodegenerative disease or cognitive impairment through disruption in brain neurotransmitter systems, biomolecular level of the brain, and brain metabolism. Potential Regulation of Glucose Transporter-1 (GLUT1) in the blood-brain barrier respons to various stress-related brain pathological conditions. The purpose of this study is to explain the influence of chronic stress intensity towards expression of glucose transporter-1 in the brains of rat. This study used 30 rats (Rattus norvegicus) male wistar strain which divided into 3 groups: control group (K0), the treatment of swimming training in severe intensity (K1) and swimming training in mild intensity (K2). Examination was conducted on GLUT1 expression in endothelial cells of the brain that be observed using immunohistochemical staining techniques. The results showed that there was depletion of GLUT1 expression in brain endothelial due to swimming training in severe intensity and swimming training in mild intensity. GLUT1 expression depletion in brain endothelial of severe intensity group was greater than in mild intensity group with p <0.05. (FMI 2012;48:24-27)

Keywords: GLUT1 (glucose transporter-1), chronic stress, brain

**Correspondence:** Hayuris Kinandita, Department of Medical Physiology, Faculty of Medicine, Airlangga University, Jl. Mayjen Prof.Dr.Moestopo 47 Surabaya 60131, Indonesia. Phone: 031-5030252-3 ext. 142. E-mail: hayuris\_k@yahoo.com

## INTRODUCTION

Stress has become a part of human and other organisms' life that can cause effects of on change in the organism, both physiological and psychological. Stress, as first proposed by Hans Selye, stress physiologist in 1976, is the body's non-specific response to various stressful or threatening situation to the organism. Prolonged stress, besides being recognized as the aetiopathogenesis of several metabolic diseases such as hypertension, coronary heart disease, and diabetes, can also cause changes in brain function (Savtchouk & Liu 2011). The

brain as one of vital organs of the body is an organ that is highly dependent on glucose as an energy provider and for the synthesis of essential amino acids as brain neurotransmitter. If the glucose levels in the brain is in a state of homeostasis and uncompensated, brain function can be impaired (Benton et al 1996).

The presence of glucose transporter proteins (GLUT) in the brain is important for ensuring glucose supply to the brain. To date, three GLUT isoforms have been identified to have significant role in brain, i.e. GLUT1, GLUT3, and GLUT5. GLUT1 has the highest concentration in the brain capillary blood vessels endothelium. Although the capillary blood vessels of the brain is only about 2% of total brain volume, the presence of GLUT1 in these capillaries remain the focus of several studies because of GLUT1 is glucose transporter that has to be passed by glucose to enter the brain (Maher et al 1994).

GLUT1 is GLUT whose expression is induced by stress state (stress-inducible (Wertheimer et al 1991). GLUT1 expression can be increased when cells are stressed (Carruthers et al 2009, Barros et al 1997). However, prolonged stress causes impaired synthesis of GLUT1 so that GLUT1 expression decreases (Duelli & Kuschinsky 2001). Physical exercise, in addition to having benefits for the body, is one of the stressors that can cause physiological changes in the body. Swimming as a form of physical exercise is a physical stressor easily used to model the stress in experimental animals. Swimming is not just a simple exercise, but the work load in swimming is high and the involvement of emotional factors in animals is difficult to eliminate that are likely to lead to a more integral stress response (Nayanatara et al 2005, Bhatia et al 2011). Level of stress such as differences in the intensity of swimming causes different physiological responses to cellular level. However, the effect of physical exercise as stress induction on the expression of GLUT1 was not yet clear. Based on the description above, the authors wanted to determine the effect of prolonged stress by inducing stress through physical exercise of swimming in different degrees of intensity on decreased expression of glucose transporter-1/GLUT1 in rats' brain.

# MATERIALS AND METHODS

The subjects were male Wistar rats aged 2-3 months, weighing 150-160 grams. Sample size was 30 rats. This study used a randomized post test only control group design. Eligible rats were divided into 3 groups: control group or no treatment (K0), the treatment group with high intensity swimming exercise (K1), and the treatment group with low intensity training exercise (K2). Physical stress used to was swimming exercise to induce stress (Forced Swimming Induced Stress) by modification of the method developed by Porsolt et al in 1977 (cited by Nayanatara et al 2005).

Swimming exercises for stress induction was done by making the animals, which had been given with a load 3% of the body weight, to swim for 70% of swimming maximum capacity for treatment group K1 and 30% of

swimming maximum capacity for treatment group K2, or until the animals experienced fatigue during the session of stress provision. The animals swam in a place that had been filled with water in a temperature of  $25 \pm 2^{\circ}$ C and a water depth of 30 cm. Swimming exercise was performed one session per day for 7 consecutive days for the effects of chronic stress (Nayanatara et al 2005). Experimental animals were sacrificed on the last day of the treatment, the seventh day, at the same time for all control and treatment groups, then we performed animal brains harvesting.

Preparation of brain tissue was made by the method of paraffinization. Then, GLUT1 expression was characterized using immunohistochemical staining. The counting of GLUT1 expressing capillary endothelium was performed quantitatively express in visual field as many as the average number of brain capillaries cut transversely on a piece of brain tissue slides.

# **RESULTS**

The study lasted for 3 months with no samples of experimental animals died or dropped out as the research proceeds. Preparation, immunohistochemical staining, and observation of GLUT1 expression in brain tissue could be performed on the entire sample. GLUT1 expression profile in brain capillary endothelium by immunohistochemical staining in control group and treatment group can be seen in Figure 1.

GLUT1 expression assessment was performed in 18 brain capillaries cut transversely, which was about the average capillary number of whole sample preparations. The descriptive analysis of GLUT1 expression in brain endothelial shows that there was a decrease in GLUT1 expression in both treatment groups compared to that in control group. The least GLUT1 expression was present in treatment group with high intensity swimming exercise followed treatment group with low intensity swimming exercise.

The results of analytical analysis using ANOVA showed significant differences (p < 0.05) of GLUT1 expression in brain capillary endothelium between groups with p = 0.0001. Least significance differences between groups was followed by Least Significance Difference (LSD) test which also showed significant values (p <0.05) with p = 0.0001 for the difference between K0 and K1, p = 0.03 for the difference between K0 and K2, and p = 0.03 for the difference between K1 and K2.




Figure 1. GLUT1 expression profile in brain capillary endothelium in control group (K0), treatment group with high intensity swimming exercise (K1), and treatment group with low intensity swimming exercise (K2)

# DISCUSSION

Stress induced by swimming exercise (Forced Swimming Induced Stress) is a type of physical stress effectively used to study the physiological changes and the capacity of organism to respond to stress using experimental animals. This is because swimmingexercise is not just a simple exercise, but it also involves emotional factors that influence the degree of stress in experimental animals (Navanatara et al 2005). Physiological stress is needed to activate various responses of the body as a defense mechanism. This is the expected positive effects of stress, the stress that can be tolerated by the body. Prolonged (chronic) stress or with excessive intensity can cause pathological negative effects on the body because the body experiences fatigue, resulting in organ dysfunction and even death (Selve 1950). The difference in chronic stress exposure intensity to an organism, such as low or high intensity, may cause different organism's response to stress.

The results showed that there is a decrease in brain endothelial GLUT1 expression in treatment group with high intensity swimming exercise (K1) with a mean of  $28.20\pm6.60$  and treatment group with low intensity swimming exercise (K2) with a mean of  $35.90\pm9.39$  when compared with the control group (K0) which had a mean of  $43.60\pm6.04$ . The results of the ANOVA analysis showed that there were significant differences in GLUT1 expression data on brain endothelium. Results of LSD between each group showed significant decrease with p < 0.05. This suggests that swimming exercise as chronic stress induction can cause changes in GLUT1 expression in brain endothelium.

Decrease in GLUT1 expression in brain endothelial due either to high or low swimming exercise intensity in this study was in accordance with some literature indicating that chronic or repeated stress is one mechanism that causes neurodegeneration through hypersecretion of glucocorticoids. Neurodegeneration can lower mRNA levels of a protein, including membrane transporters, through the inhibition of insulin secretion, an important hormone in protein synthesis (Somwar et al, 1998, Munhoz et al 2008).

GLUT1 expression in brain endothelial of treatment group with high intensity swimming exercise showed higher decrease than that in treatment group with low intensity swimming exercise with a significance level of p = 0.03. It resulted from damage of endothelial membranes due to excessive stress intensity, coupled with higher decrease in GLUT1 mRNA synthesis in the treatment group with high intensity swimming exercise than that in group with low intensity swimming exercise, so that the cell membrane GLUT1 expression would also decrease even though it received adequate activation signals (Somwar et al, 1998, Munhoz et al 2008).

Physical stress is a trigger of oxidative stress that may increase the levels of free radicals in the body. Chronic stress with tolerable intensity may reduce level of cell oxidative stress and improves cell ascorbic acid as an endogenous antioxidant in the brain. This provides the beneficial effects of physical exercise (Liu et al 2000). Other literature about the effect of free radicals on protein synthesis mentioned that excessive free radicals formed will increase oxidative damage, inhibit insulin gene expression, decrease insulin secretion, and ultimately lower protein synthesis (Mulukutla et al 2010, Somwar et al 2000). The intensity of chronic stress in experimental animals in this study was actually a submaximal and light intensity. However, the emotional factor that also involved in the experimental animals added the stress intensity so that the pathological effects of the stress became predominant, indicating by the decrease in GLUT1 expression in brain endothelial and glial cells of he experimental animals.

The decrease in GLUT1 expression in rat brain capillary endothelium, both in high and low intensity swimming exercise treatment as chronic stress induction, indicated that difference in the intensity or degree of stress lowers brain GLUT1 expression. Chronic stress given continuously or without time lag causes the body does not have time for adaptation and recovery process as a form of compensation to overcome the damage up to cellular level that occurs as a result of stress. Long-term and continuous stress treatment induces cell malfunctions.

# **CONCLUSION**

High intensity swimming exercise as chronic stress induction causes the highest decrease in capillary endothelial GLUT1 expression of white rat's brain. Low intensity swimming exercise as chronic stress induction also causes a decrease in capillary endothelial GLUT1 expression of white rat's brain with a statistically significant reduction compared to that in untreated control group. Further research is needed to determine the levels of biological stress markers in experimental animals to find the proper dosage of stress for experimental animals.

# ACKNOWLEDGEMENTS

We thank the research grant program of Airlangga University that had provided fund for the implementation of this study.

# REFERENCES

- Barros LF, Young M, Saklatvala J, Baldwin SA (1997). Evidence of two mechanisms for the activation of the glucose transporter GLUT1 by anisomycin: p38 (MAP kinase) activation and protein synthesis inhibition in mammalian cells. J Physiol 504, 517-525
- Benton D, Parker PY, Donohoe RT (1996). The supply of glucose to the brain and cognitive functioning. J Biosoc Sci 28, 463-479
- Bhatia N, Maiti PP, Choudhary A, Tuli A, Masih D, Khan MMU, Ara T, Jaggi AS (2011). Animal models

- in the study of stress: a review. NSHM Journal of Pharmacy and Healthcare Management 2, 42-50
- Carruthers A, DeZutter J, Ganguly A, Devaskar SU (2009). Will the original glucose transporter isoform please stand up! Am J Physiol Endocrinol Metab 297, E836–E848
- Duelli R and Kuschinsky W (2001). Brain glucose transporters: relationship to local energy demand. News Physiol Sci 16, 71-76
- Liu Y, Lormes W, Baur C, Opitz-Gress A, Altenburg D, Lehmann M, Steinacker JM (2000). Human skeletal muscle HSP70 response to physical training depends on exercise intensity. Int J Sports Med 21, 351-355
- Maher F, Vannucci SJ, Simpson IA (1994). Glucose transporter proteins in brain. FASEB J 8, 1003-1011
- Mulukutla BC, Khan S, Lange A, Hu WS (2010). Glucose metabolism in mammalian cell culture: new insights for tweaking vintage pathways. Trends Biotechnol 28, 476-484
- Munhoz CD, García-Bueno B, Madrigal JLM, Lepsch LB, Scavone C, Leza JC (2008). Stress-induced neuroinflammation: mechanisms and new pharmacological targets. Braz J Med Biol Res 41, 1037-1046
- Nayanatara AK, Nagaraja HS, Anupama BK (2005). The effect of repeated swimming stress on organ weights and lipid peroxidation in rats. Thai Journal of Physiological Sciences 18, 3-9
- Savtchouk I and Liu SJ (2011). Remodeling of synaptic AMPA receptor subtype alters the probability and pattern of action potential firing. Journal of Neuroscience 31, 501
- Selye H (1950). Stress and the general adaptation syndrome. Br Med J 1, 1383–1392
- Somwar R, Perreault M, Kapur S, Taha C, Sweeney G, Ramlal T, Kim DY, Keen J, Côté CH, Klip A, Marette A (2000) Activation of p38 mitogen-activated protein kinase a and β by insulin and contraction in rat skeletal muscle. Diabetes 49, 1794-1800
- Somwar R, Sumitani S, Taha C, Sweeney G, Klip A (1998). Temporal activation of p70 S6 kinase and Akt1 by insulin: PI 3-kinase-dependent and independent mechanisms. Am J Physiol 275, E618-E625
- Wertheimer E, Sasson S, Cerasi E, Ben-Neriah Y (1991). The ubiquitous glucose transporter GLUT-1 belongs to the glucose-regulated protein family of stress-inducible proteins. PNAS 88, 2525-2529