

PROCEEDING

International Seminar

THE ROLE OF VETERINARY SCIENCE
TO SUPPORT MILLENNIUM DEVELOPMENT GOALS
and

THE 12th ASIAN ASSOCIATION OF VETERINARY SCHOOLS CONGRESS




FACULTY OF VETERINARY MEDICINE
UNIVERSITAS AIRLANGGA

PROCEEDING

International Conference

**THE ROLE OF VETERINARY SCIENCE
TO SUPPORT MILLENNIUM DEVELOPMENT GOALS
*and***

THE 12th ASIAN ASSOCIATION OF VETERINARY SCHOOLS CONGRESS



**FACULTY OF VETERINARY MEDICINE
UNIVERSITAS AIRLANGGA**

ISBN 978-602-70438-0-0



9786027043800

290a

PROCEEDING

International Seminar

**THE ROLE OF VETERINARY SCIENCE TO SUPPORT
MILLENNIUM DEVELOPMENT GOALS
AND THE 12th ASIAN ASSOCIATION OF
VETERINARY SCHOOLS CONGRESS
JW MARRIOTT HOTEL, SURABAYA-INDONESIA
4th - 6th SEPTEMBER 2013**

Editors :

Rahaju Ernawati

Fedik Abdul Rantam

Bambang Sektiari

Didik Handijatno

Mustofa Helmi Effendi

Dadik Raharjo

Epy Muhammad Luqman

Lita Rakhma Yustinasari

Widya Paramita Lokapirnasari

**FACULTY OF VETERINARY MEDICINE
UNIVERSITAS AIRLANGGA**

List of Content

	Page
Arts in Pig, Sheep and Goat: Experiences in Thailand M Techakumphu, C. Tretipskul, N, Anakkul, S. Panyaboriban ¹ , J. Suwimon- teerabutr and T. Tharasanit	1
Food Security - The Role of Veterinarians in an Emerging World Problem ID Robertson	6
Past, present and Future of Asian Society off Zoo and Wildlife Medicine (aszwm) Junpei Kimura	11
Standardized Veterinary Competence to Support Global Food Security and Infectious Animal Disease Control Wiwiek Bagja	12
How to Implement Policies and Regulations to Control Antimicrobial Resistance in Animal Husbandry? Yong Ho Park	13
Warning for Animal Parasites on HIV/AIDS Patients Nasronudin	16
Vector Vaccines: a New Generation of Veterinary Vaccines M. Hair-Bejo	21
One Health System for Controlling Zoonotic Diseases Srihadi Agungpriyono, Denny Widaya Lukman	25
Development of a Novel Oral Contraceptive for the Population Control of Wild Mammals Based on the Neuroendocrine Mechanism Generating Gonadotropin Releasing Hormone (GNRH) Pulse Generation Kei-ichiro Maeda, TanuPinyopummintr, Hiroko Tsukamura	27
Reflection of Animal Welfare Principles as a Part of Professionalism Romziah, S.	31
Non Typhoidal Salmonellosis as Food Borne Disease Dadik Raharjo	37
Current Issue on Feed Additives Utilization in Indonesia Budi Tangendjaja	42
Mitochondrial Genetic Defect And Disease in Human Agung Pranoto	51

Expression of Cytochrome C as Apoptotic Indicator and it Relation With Sper Viability and Motility of Domba Ekor Gemuk Frozen Semen In Differe Thawing Duration	
Rahmalia Dwi Suindarti, Imam Mustofa, Suherni Susilowati	213
Effects of the Timing of Insemination by The Use of a Heat Detector on the Incidence of Metestrous Bleeding and Non-Return Rate at Day 21	
Ismudiono, Pudji Srianto and Trilas Sardjito	217
Rapid Detection and Phylogenetic Analysis of West Nile Virus as Zoonosis New Emerging Disease in Patients with Fever of Unknown Origin (FUO) in Surabaya	
E. Bimo Aksono H, Nasronudin, Maria Inge Lusida, Aldise Marieta N, M. Qushai, N. Fajar, Lilis Mundri Jannah, Brian Eka Rachman, Musofa Rusli	221
Comparative Study of Pathogenicity Of H5n1 Virus Between in Tree Sparrow, Scaly Breasted Munia, And Backyard Chickens As Natural Source of Infection in East Java - Indonesia	
Emmanuel Djoko Poetranto, Djoko Legowo, Suwarno, Fedik A Rantam.....	226
Toxoplasmosis :Changes in Trophoblast Apoptosis index Mice (<i>Mus musculus</i>) Given Anti-<i>Toxoplasma gondii</i> ESA (Excretory Secretory Antigen) Immunoglobulin Y	
Lucia Tri Suwanti, Hani Plumeriastuti, Dessy Fajarwati	233
Egg Yolk Derived Anti-Rabies Antibody Production as Immunotherapy Agents	
Suryo Kuncorojakti	240
Relations of Weight And Age to The Front Feet Sole Area of Merino Ram	
Benjamin Christoffel Tehupuring, Dady Soegianto Nazar, Sarmanu.....	244
Molecular Characterization of Nucleoprotein <i>Antigenic Sites</i> of Indonesian Isolate Rabies Virus	
Jola Rahmahani and Suwarno	248
Homology Analysis of G Protein Coding Genes of Rabies Virus Sulawesi Isolate Against Pasteur Strain	
Riski Arya Pradikta and Suwarno	254
The Production of Plastic Progesterone Implants for Estrus Synchronization in Big Tail Sheep From Sapudi Island	
Sunaryo Hadi Warsito, Setyawati Sigit, Herry Agoes Hermadi	258
Production of Frozen Dry Equine Chorionic Gonadotrophin (eCG) from Pregnant Mare Sera	
Herry Agoes Hermadi, Laba Mahaputra	262
Acrosin Half-Breed Etawa Goat (PE) Sperm Characteristic to Increase Spermatozoa Quality	
Budi Utomo	270

RAPID DETECTION AND PHYLOGENETIC ANALYSIS OF WEST NILE VIRUS AS ZONOSIS NEW EMERGING DISEASE IN PATIENTS WITH FEVER OF UNKNOWN ORIGIN (FUO) IN SURABAYA

E. Bimo Aksono H*; Nasronudin*; Maria Inge Lusida*; Aldise Marieta N*; M. Qushai*; N. Fajar*; Lilis Mundri Jannah*; Brian Eka Rachman*; Musofa Rusli**;

*Institute of Tropical Disease, Airlangga University, Surabaya

**Divisi Tropik-Infeksi, Departemen Ilmu Penyakit Dalam, RSUD. Dr. Soetomo/FK-Unair

Surel : baksono@yahoo.com

ABSTRACT

This study aims to perform a rapid detection and phylogenetic analysis of WNV in the serum of patients with clinical isolates with Fever of Unknown Origin. Serum samples in Surabaya came from 2 patients who had been examined by a Specialist Doctors In the June-July, 2013, with symptoms of fever, headache, sore throat, back pain, myalgia, arthralgia, fatigue, conjunctivitis, pain, anorexia, nausea, abdominal pain, diarrhea, and respiratory symptoms. Running RNA extraction procedure and nested PCR PCR is run in accordance with protocols west nile virus detection kit from Intron. Similarly, primers used are designed specifically for the west nile virus. Interpretation of PCR products using 1% agarose gel (306 bp), and then performed a homology analysis using Genetix Mac Ver. 8.0 with reference WNV isolates from the gene bank (JF920741.1.gnu). This research demonstrated the effective PCR methods for the rapid detection of WNV and diagnosis of WNV infection from clinical isolates represent a breakthrough in the clinical world to many cases of infectious disease that has not been diagnosed (Fever Unknown Origin = FUO). Based on homology analysis, This sample showed 95% with WNV were reported in the gene bank (JF920741.1.gnu).

Keywords : PCR, West Nile Virus, FUO, Patients, Mosquitos

INTRODUCTION

West Nile virus (WNV) is an infectious disease that is new emerging diseases, mosquito-borne and has been reported to be found throughout Europe, Russia, Africa, Southwest Asia, India and allegedly including Indonesia. West nile virus is transmitted by mosquitoes of the genus *Culex*, especially that has been infected with WNV. WNV is a member of the Japanese encephalitis antigenic complex of the genus *Flavivirus*, family *Flaviviridae* (Gubler, 2007).

According Smithburn *et al* cited by Gubler (2007) that WNV was first isolated from blood specimens obtained from patients in the province of West Nile of Uganda in 1937. From that time until the fall of 1999, WNV is considered relatively unimportant as human and animal pathogens. The virus was enzootic throughout Africa, West and Central Asia, Middle East, and the Mediterranean, with occasional extension into Europe (Hayes, 1989). A

subtype of WNV (Kunjin) is also found in Australia. Epidemiological characteristics for the period 1937-1999, WNV infection only occurs occasionally in humans, horses, and birds and is generally asymptomatic or mild, while the associated neurological disease and death are rarely reported (Gubler, 2007; Murgue *et al*, 2001; Murgue *et al*, 2002). One exception was a small outbreak of WNV infection in the Camargue, area of France in 1960 at the expected associated with severe and fatal neurological disease in horses and humans. Long intervals between epidemics, and the perception that WNV is not an important issue related to public health, and the lack of reporting of epidemics and / or epizootics in the Mediterranean region opened the realization that the emerging epidemic of WNV infection is strongly associated with severe neurological disease and can be fatal, especially in the case of 1990.

West Nile Virus (WNV) can survive in the environment because of the arthropod vectors that transmit the virus between vertebrate hosts. The main vector for WNV in the United States is the *Culex pipiens* mosquitoes usually breed in urban areas and prefer to bite birds. The main line of West Nile virus infection in humans is through the bite of an infected mosquito. Mosquitoes are infected when they bite infected birds, then WNV mosquito can circulate in the blood for several days. WNV finally entered the salivary glands of mosquitoes. Furthermore, the virus can be transmitted to humans and animals (Peterson *et al*, 2003). WNV has been detected in many species of wild birds, including the American crow.

Imad *et al* (2005) quoting from various sources report that during the year 2002 reported that a WNV outbreak of 2002, can be transmitted through organ transplantation and blood transfusion, one case was reported through organ transplants and one suspected case of transmission through breastfeeding. Reported two cases of WNV infection in laboratory workers.

WNV incubation period is estimated to range between 3 to 14 days. WNV infection is generally mild and often clinically unapparent (Campbell *et al*, 2002; Gubler, 2007). It is estimated that 20% of those infected will show mild symptoms, including fever, headache, and general hospital for 3 to 6 days (Sejvar *et al*, 2003; Peterson *et al*, 2003; Gubler, 2007). An estimated one in 150 patients infected with WNV show symptoms of meningoencephalitis. Patients who have a risk of serious infections and deaths including immunocompromised, elderly, and very young, with symptoms including severe headache, high fever, neck stiffness, stupor, disorientation, coma, tremors, convulsions, muscle weakness, and abnormal movements (Sejvar *et al*, 2003; Peterson *et al*, 2003; Gubler, 2007; Klein *et al*, 2002)

According Guebler (2007) quotes from several sources reported that the vitritis and chorioretinitis have been found in patients with WNV infection. Some patients with WNV infection present with acute flaccid paralysis similar to polio. A minority of patients with severe disease developed into a maculopapular or morbilliform rash involving the neck, trunk, arms, or legs. According to data from the CDC 2002, the mortality rate is estimated at 7%, with most deaths related to complications of meningoencephalitis. Most patients who survive this disease will have a complete recovery, but some patients with meningoencephalitis with acute flaccid paralysis will require long-term rehabilitation.

Diagnosis of WNV infection is based on clinical symptoms and laboratory tests specific. WNV, or other arboviral diseases such as St. Louis encephalitis, should be considered in people suffering from encephalitis or meningitis in summer or early fall. Total number of leukocytes in peripheral blood is generally reported normal or slightly elevated with lymphocytopenia. Sometimes found hyponatremia, especially in patients with encephalitis. Examination of cerebrospinal fluid (CSF) may show pleocytosis, usually with a predominance of lymphocytes. Protein is universally increased. Glucose is usually normal (Sejvar *et al*, 2003; Peterson *et al*, 2003; Guebler, 2007; Klein *et al*, 2002). The most efficient diagnostic method is detection of IgM antibodies to WNV in CSF within 8 days of onset using the IgM antibody capture enzyme-linked immunosorbent assay (MAC-ELISA). Detection of IgM antibodies in the CSF are more specific to central nervous system infection because IgM antibodies do not cross the blood-brain barrier. The finding that WNV IgM antibody may persist in the blood for up to 500 days to make a single blood sample testing problem. Thus, acute and recovered blood samples needed for confirmation.

However in Indonesia have not been many reports of cases of WNV which has infected humans and in mosquitoes. On the other hand there are many reported cases of the infection have no clear diagnosis or Fever of Unknown Origin. This study aims to perform a rapid detection and phylogenetic analysis of WNV in the serum of patients with clinical isolates with Fever of Unknown Origin.

MATERIAL AND METHODS

The research was conducted at the Institute of Tropical Disease, Airlangga University. Serum samples in Surabaya came from 2 patients who had been examined by a Specialist Doctors In the June-July, 2013, with symptoms of fever, headache, sore throat, back pain, myalgia, arthralgia, fatigue, conjunctivitis, pain, anorexia, nausea, abdominal pain, diarrhea, and respiratory symptoms. Running RNA extraction procedure and nested PCR PCR is run in accordance with protocols west Nile virus detection kit from Intron. Similarly, primers used are designed specifically for the west Nile virus. Interpretation of PCR products using 1% agarose gel (306 bp), and then performed a homology analysis using Genetix Mac Ver. 8.0. The study was carried out in accordance with ethical clearance by RS. Unair.

RESULTS AND DISCUSSION

The results showed that serum PCR positive patients contain the infection with WNV. DNA PCR product nucleotide with length 104 bp and the results of homology analysis showed 95% with WNV were reported in the gene bank (JF920741.1.gnu) (figure 3)




Figure 1. PCR products of west Nile virus from clinical isolates in Surabaya in the 1% gel electrophoresis

CONCLUSION

This research demonstrated the effective PCR methods for the rapid detection of WNV and diagnosis of WNV infection from clinical isolates represent a breakthrough in the clinical world to many cases of infectious disease that has not been diagnosed (Fever Unknown Origin = FUO). Based on homology analysis, This sample showed 95% with WNV were reported in the gene bank (JF920741.1.gnu).




Figure 2. Sequencing results of west nile virus from clinical isolates in Surabaya

[GENETYX : Homology Data]

Date : 2013.08.15

Query Sequence

File Name :

Sequence Name : WNV-F6-3-13-7-16

Sequence Size : 111

Target Sequence

File Name : JF920741.1.gnu

Sequence Name : JF920741.1.gnu

Sequence Size : 80

Query Range: 5 - 85

Sbjct Range: 1 - 80

81 bp, INT.Score: 194, OPT.Score: 290

Identity: 77 / 81 (95%)

Similarity: 77 / 81 (95%)

Strand: Plus / Plus

Query 5 GGAAAACAAACAGCGATGAAACACCTTCTGGAGTTTTAAGAAGGGACTAGGGACCTTGA 64

Sbjct 1 GAACAAACAAACAGCGATGAAACACCTTCT - GAGTTTTAAGAAGGAAGGACTAGGGACCTTGA 59

Query 65 CCAGTGCTATCAATCGGCGGG 85

Sbjct 60 CCAGTGCTATCAATCGGCGGG 80

Figure 3. Homology analysis of west nile virus from clinical isolates in Surabaya

REFERENCES

- Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. 2002. West Nile virus. *Lancet Infect Dis* 2: 519-529
- Gubler, D.J. 2007. The Continuing Spread of West Nile Virus in the Western Hemisphere. *Clin Infect Dis*. 45 (8): 1039-1046.
- Hayes, C. 1989. West Nile Fever. In: Monath TP, editor. *The arboviruses: epidemiology and ecology*. Boca Raton, FL: CRC Press. 59-88.
- Imad T.Z; Deniz A; Joseph R; Merline; Subhash C; Karl K.k. 2005. West Nile virus infection. *Am. J. Rontgenology*. Vol 184 (3).
- Klein C; Kimiagar I; Pollak L; et al. 2002. Neurological features of West Nile virus infection during the 2000 outbreak in a regional hospital in Israel. *J Neurol Sci*. 200:63-66.
- Murgue, B; Murri, S; Triki, H, et al. 2001. West Nile in the Mediterranean basin: 1950-2000. In: White DJ, Morse DL, editors. *West Nile virus, detection, surveillance and control*. New York: New York Academy of Sciences. 117-26.
- Murgue, B; Zeller, H; Deubel, V. 2002. The ecology and epidemiology of West Nile virus in Africa, Europe, and Asia. In: MacKenzie JS, Barrett ADT, Deubel V, editors. *Japanese encephalitis and West Nile viruses*. Berlin: Springer-Verlag. 195-221.
- Peterson, LR; Marphin AA; Gubler DJ. 2003. West Nile virus. *JAMA* 290:524-528.
- Sejvar JJ; Haddad MB; Tierney BC; et al. 2003. Neurologic manifestations and outcome of West Nile virus infection. *JAMA* 2003; 290:511-515



CERTIFICATE OF ATTENDANCE

This is to certify that

E. Bimo Aksono

AS SPEAKER

in INTERNATIONAL SEMINAR
THE ROLE OF VETERINARY SCIENCE
TO SUPPORT MILLENNIUM DEVELOPMENT GOALS

was held in JW Marriott Hotel Surabaya - Indonesia

September 5th - 6th, 2013

President of AAVS
Dean of Faculty Veterinary Medicine
Universitas Airlangga - Indonesia

Prof. Hj. Romziah Sidik., DVM., Ph.D.





**KOMITE ETIKA DAN HUKUM
RUMAH SAKIT UNIVERSITAS AIRLANGGA**

**KETERANGAN LOLOS KAJI ETIK
No :141A/KEH/2013**

Komite Etika dan Hukum Rumah Sakit Universitas Airlangga dalam upaya melindungi hak asasi dan kesejahteraan subyek penelitian, telah mengkaji dengan teliti protokol penelitian yang berjudul:

“Rapid Detection and Phylogenetic Analysis of West Nile Virus
as Zoonosis New Emerging Disease in Patients with Fever of Unknown Origin (FUO) in Surabaya”

Peneliti Utama : E. Bimo Aksono H

Anggota Peneliti : 1) Nasronudin
2) Maria Inge Lusida
3) Aldise Marieta N
4) M. Qushai
5) N. Fajar
6) Lilis Mundri Jannah
7) Brian Eka Rachman
8) Musofa Rusli

Institusi Penelitian : Rumah Sakit Universitas Airlangga Surabaya

DINYATAKAN LAIK ETIK.

Surabaya, 17 April 2013



Ketua

Prof.Dr.Med. H. M. Soekry Erfan Kusuma, dr.Sp.F (K), DFM