

DAFTAR PUSTAKA

- Agofure, O. 2017. Prevalence of obesity among adults in Issele-Uku, Delta State Nigeria. *Alexandria Journal of Medicine*, 54(4). pp. 463–468. <https://doi.org/10.1016/j.ajme.2017.10.005>.
- Akpınar, E., Bashan, I., Bozdemir, N. and Saatci, E. 2007. Which is the best anthropometric technique to identify obesity: body mass index, waist circumference or waist-hip ratio?. *Collegium antropologicum*, 31(2). pp. 387–93.
- Aktar, N., Qureshi, N. K. and Ferdous, H. S. 2017. Obesity: A Review of Pathogenesis and Management Strategies in Adult. *Delta Medical College Journal*, 5(1). pp. 35–48. <https://doi.org/10.3329/dmcj.v5i1.31436>.
- Akter, S., Rahman, M.M., Abe, S.K. and Sultana, P. 2014. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bulletin of the World Health Organization*, 92(3). pp. 204-213A. <https://doi.org/10.2471/BLT.13.128371>.
- Ammann, B.C., Knols, R.H., Baschung, P., de Bie, R.A. and de Bruin, E.D. 2014. Application of principles of exercise training in sub-acute and chronic stroke survivors: A systematic review. *BMC Neurology*, 14(1). pp. 1–11. <https://doi.org/10.1186/s12883-014-0167-2>.
- Avloniti, A.A., Douda, H.T., Tokmakidis, S.P., Kortsaris, A.H., Papadopoulou, E.G. and Spanoudakis, E.G. 2007. Acute effects of soccer training on white blood cell count in elite female players. *International journal of sports physiology and performance*, 2(3). pp. 239–249. <https://doi.org/10.1123/ijspp.2.3.239>.
- Bacopoulos, F., Efthymiou, V., Landis, G., Rentoumis, A. and Chrousos, G.P. 2015. Waist circumference, waist-to-hip ratio and waist-to-height ratio reference percentiles for abdominal obesity among Greek adolescents. *BMC Pediatrics*, 15(1). pp. 1–9. <https://doi.org/10.1186/s12887-015-0366-z>.
- Bækkerud, F.H., Solberg, F., Leinan, I.M., Wisløff, U., Karlsen, T. and Rognmo, Ø. 2016. Comparison of Three Popular Exercise Modalities on V' O₂max in Overweight and Obese. *Medicine & Science in Sports & Exercise*, 48(3). pp. 491–498. <https://doi.org/10.1249/MSS.0000000000000777>.
- Bales, C.W. and Buhr, G. 2008. Is Obesity Bad for Older Persons? A Systematic Review of the Pros and Cons of Weight Reduction in Later Life. *Journal of the American Medical Directors Association*, 9(5). pp. 302–312. <https://doi.org/10.1016/j.jamda.2008.01.006>.

- Baltpopoulos, P. 2009. Exercise induced modulation of immune system functional capacity. *Journal Biology of Exercise*, 5(1). pp. 40–50. <https://doi.org/10.4127/jbe.2009.0024>.
- Bautista, R.J.H., Mahmoud, A.M., Königsberg, M. and Guerrero, N.E.L.D. 2019. Obesity: Pathophysiology, monosodium glutamate-induced model and anti-obesity medicinal plants. *Biomedicine & Pharmacotherapy*, 111. pp. 503–516. <https://doi.org/10.1016/j.biopha.2018.12.108>.
- Beechy, L. Galpern, J., Petrone, A. and Das, S.K. 2012. Assessment tools in obesity - Psychological measures, diet, activity, and body composition. *Physiology and Behavior*, 107(1). pp. 154–171. <https://doi.org/10.1016/j.physbeh.2012.04.013>.
- Benedini, S., Dozio, E., Invernizzi, P.L., Vianello, E., Banfi, G., Terruzzi, I., Luzi, L. and Romanelli, M. M. 2017. Irisin: A Potential Link between Physical Exercise and Metabolism - An Observational Study in Differently Trained Subjects, from Elite Athletes to Sedentary People. *Journal of Diabetes Research*, 2017. pp. 1–7. <https://doi.org/10.1155/2017/1039161>.
- Blaak, E. 2001. Gender differences in fat metabolism. *Current Opinion in Clinical Nutrition and Metabolic Care*, 4(6). pp. 499–502. <https://doi.org/10.1097/00075197-200111000-00006>.
- Bloom, M.D. and Kupfer, D.J. 1995. *Psychopharmacology; the fourth generation of progress*. An official publication of the American College of Neuropsychopharmacology. New York: Raven Press. pp. 719-729.
- Bompa, T.O. and Haff, G.G. 1994. *Periodization Theory and Methodology of Training Fourth Edition*. United State of America: Human Kinetics Phublisher.
- Bompa, T.O. and Haff, G.G. 2009. *Periodization Theory and Methology of Training Fifth Edition*. United State of America: Human Kinetic Phublisher.
- Boström, P., Wu, J., Jedrychowski, M.P., Korde, A., Ye, L., Lo, J.C., Rasbach, K.A., Boström, E.A., Choi, J.H., Long, J.Z., Kajimura, S., Zingaretti, M.C., Vind, B.F., Tu, H., Cinti, S., Hojlund, K., Gygi, S.P. and Spiegelman, B.M. 2012. A PGC1 α -dependent myokine that drives browning of white fat and thermogenesis. *Nature*, 481(7382). pp. 463–468. <https://doi.org/10.1038/nature10777.A>.
- Cannon, B. and Nedergaard, J. 2004. Brown Adipose Tissue: Function and Physiological Significance. *Physiological Reviews*, 84(1). pp. 277–359. <https://doi.org/10.1152/physrev.00015.2003>.
- Carlson NR, 1994. *Physiology of behavior*. Boston: Allyn and Bacon. pp. 235-250.

- Cedikova, M., Kripnerová, M., Dvorakova, J., Pitule, P., Grundmanova, M., Babuska, V., Mullerova, D. and Kuncova, J. 2016. Mitochondria in White, Brown, and Beige Adipocytes. *Stem Cells International*, 2016. pp. 1–11. <https://doi.org/10.1155/2016/6067349>.
- Chooi, Y.C., Ding, C. and Magkos, F. 2019. The epidemiology of obesity. *Metabolism: Clinical and Experimental*, 92. pp. 6–10. <https://doi.org/10.1016/j.metabol.2018.09.005>.
- Cocks, M., Shaw, C.S., Shepherd, S.O., Fisher, J.P., Ranasinghe, A., Barker, T.A. and Wagenmakers, A.J.M. 2016. Sprint interval and moderate-intensity continuous training have equal benefits on aerobic capacity, insulin sensitivity, muscle capillarisation and endothelial eNOS/NAD(P) Hoxidase protein ratio in obese men. *Journal of Physiology*, 594(8). pp. 2307–2321. <https://doi.org/10.1113/jphysiol.2014.285254>.
- Cooper, C.E., Vollaard, N.B.J., Choueiri, T. and Wilson, M. 2002. Exercise, free radicals and oxidative stress. *Biochemical Society Transactions*, 30(2). pp. 280–285.
- Coppini, L.Z., Waitzberg, D.L. and Campos, A.C.L. 2005. Limitations and validation of bioelectrical impedance analysis in morbidly obese patients. *Current Opinion in Clinical Nutrition and Metabolic Care*, 8(3). pp. 329–332. <https://doi.org/10.1097/01.mco.0000165013.54696.64>.
- Das, S.K. 2005. Body composition measurement in severe obesity. *Current Opinion in Clinical Nutrition and Metabolic Care*, 8(6). pp. 602–606. <https://doi.org/10.1097/01.mco.0000171122.60665.5f>.
- Daskalopoulou, S.S., Daskalopoulou, S.S., Cooke, A.B., Gomez, Y.H., Mutter, A.F., Filippaios, A., Mesfum, E.T. and Mantzoros, C.S. 2014. Plasma irisin levels progressively increase in response to increasing exercise workloads in young, healthy, active subjects. *European Journal of Endocrinology*, 171(3). pp. 343–352. <https://doi.org/10.1530/EJE-14-0204>.
- Demura, S. and Sato, S. 2007. Prediction of visceral fat area at the umbilicus level using fat mass of the trunk: The validity of bioelectrical impedance analysis. *Journal of Sports Sciences*, 25(7). pp. 823–833. <https://doi.org/10.1080/02640410600875010>.
- Dias, K.A., Ingul, C.B., Tjonna, A.E., Keating, S.E., Gomersall, S.R., Follestad, T., Hosseini, M.S., Hollekim-Strand, S.M., Ro, T.B., Haram, M., Huuse, E.M., Davies, P.S.W., Cain, P.A., Leong, G.M. and Coombes, J.S. 2017. Effect of High-Intensity Interval Training on Fitness, Fat Mass and Cardiometabolic Biomarkers in Children with Obesity: A Randomised Controlled Trial. *Sports Medicine*, 48(3). pp. 733–746. <https://doi.org/10.1007/s40279-017-0777-0>.

- Duclos, M. and Tabarin, A. 2011. Exercise, Training, and the Hypothalamo–Pituitary–Adrenal Axis. *Medicine & Science in Sports & Exercise*, 43(6). pp. 9–15. <https://doi.org/10.1249/01.mss.0000395470.11085.87>.
- Erickson, K. I., Voss, M.W., Prakash, R.S., Basak, C., Szabo, A., Chaddock, L., Kim, J.S., Heo, S., Alves, H., White, S.M., Wojcicki, T.R., Mailey, E., Vieira, V.J., Martin, S.A., Pence, B.D., Woods, J.A., McAuley, E., & Kramer, A.F. 2011. Exercise training increases size of hippocampus and improves memory. *Proceedings of the National Academy of Sciences*, 108(7). pp. 3017–3022. <https://doi.org/10.1073/pnas.1015950108>.
- Esfahani, M., Baranchi, M. and Goodarzi, M.T. 2016. Irisin and Metabolic Disorders. *Avicenna Journal of Medical Biochemistry*, 4(1). pp. 2–5. <https://doi.org/10.17795/ajmb-33230>.
- Eston, R., Hawes, M., Martin, A. and Reilly, T. 2009. *Kinanthropometry and Exercise Physiology Laboratory Manual: Tests, Procedures and Data. Third Edition. "Human body composition"*. Edited by Roger Eston and Thomas Reilly. New York: Routledge Taylor & Francis e-Library. <https://doi.org/10.1136/bjsm.30.2.186>.
- Fatouros, I. G. 2018. Is irisin the new player in exercise-induced adaptations or not? A 2017 update. *Clinical Chemistry and Laboratory Medicine*, 56(4). pp. 525–548.
- Fukushima, Y., Kurose, S., Shinno, H., Thu, H.C.T., Takao, N., Tsutsumi, H., Hasegawa, T., Nakajima, T. and Kimura, Y. 2016. Effects of Body Weight Reduction on Serum Irisin and Metabolic Parameters in Obese Subjects. *Diabetes Metabolism Journal*, 40. pp. 386–395. <http://dx.doi.org/10.4093/dmj.2016.40.5.386>.
- Gadde, K. M., Martin, C.K., Berthoud, H.R. and Heymsfield, S.B. 2018. Obesity: Pathophysiology and Management. *Journal of the American College of Cardiology*, 71(1). pp. 69–84. <https://doi.org/10.1016/j.jacc.2017.11.011>.
- Gallagher, D., Heymsfield, S.B., Heo, M., Jebb, S.A., Murgatroyd, P.R. and Sakamoto, Y. 2000. Healthy percentage body fat ranges: An approach for developing guidelines based on body mass index. *American Journal of Clinical Nutrition*, 72(3). pp. 694–701. <https://doi.org/10.1093/ajcn/72.3.694>.
- Garber, C.E., Blissmer, B., Deschenes, M.R., Franklin, B.A., Lamonte, M.J., Lee, I.M., Nieman, D.C. and Swain, D.P. 2011. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. *Medicine and Science in Sports and Exercise*, 43(7). pp. 1334–1359. <https://doi.org/10.1249/MSS.0b013e318213fefb>.

- Gibala, M.J., Gillen, J.B. and Percival, M.E. 2014. Physiological and Health-Related Adaptations to Low-Volume Interval Training: Influences of Nutrition and Sex. *Sports Medicine*, 44(2). pp. S127–S137. <https://doi.org/10.1007/s40279-014-0259-6>.
- Giriwijoyo, H.Y.S.S. dan Sidik, D.Z. 2012. *Ilmu Kesehatan Olahraga*. Bandung: PT Remaja Rosdakarya.
- Gist, N.H., Fedewa, M.V., Dishman, R.K. and Cureton, K.J. 2014. Sprint interval training effects on aerobic capacity: a systematic review and meta-analysis. *Sports Medicine*, 44(2). pp. 269–279. <https://doi.org/10.1007/s40279-013-0115-0>.
- Gizaw, M., Anandakumar, P. and Debela, T. 2017. A review on the role of irisin in insulin resistance and type 2 diabetes mellitus. *Journal of Pharmacopuncture*, 20(4). pp. 235–242. <https://doi.org/10.3831/KPI.2017.20.029>.
- Gledhill, N. and Jamnik, V.K. 1992. Development and validation of a fitness screening protocol for firefighter applicants. *Can J Sport Sci*, 17. pp. 199–206.
- Goody, D. and Pfeifer, A. 2019. MicroRNAs in brown and beige fat. *BBA-Molecular and Cell Biology of Lipids*, 1864(1). pp. 29–36. <https://doi.org/10.1016/j.bbalip.2018.05.003>.
- Gulyaeva, O., Dempersmier, J. and Sul, H.S. 2019. Genetic and epigenetic control of adipose development. *BBA-Molecular and Cell Biology of Lipids*, 1864(1). pp. 3–12. <https://doi.org/10.1016/j.bbalip.2018.04.016>.
- Hannan, A. L., Hing, W., Simas, V., Climstein, M., Coombes, J.S., Jayasinghe, R., Byrnes, J. and Furness, J. 2018. High-intensity interval training versus moderateintensity continuous training within cardiac rehabilitation: a systematic review and meta-analysis. *Journal of Sports Medicine*, 9. pp. 1–17. <https://doi.org/10.2147/OAJSM.S150596>.
- Hansen, D., Dendale, P., Jonkers, R.A., Beelen, M., Manders, R.J., Corluy, L., Mullens, A., Berger, J., Meeusen, R. and van Loon, L.J. 2009. Continuous low- to moderate-intensity exercise training is as effective as moderate- to high-intensity exercise training at lowering blood HbA (1c) in obese type 2 diabetes patients. *Diabetologia*, 52(9). pp. 1789–1797. <https://doi.org/10.1007/s00125-009-1354-3>.
- Harsono. 1988. *Coaching dan Aspek-aspek Psikologi dalam Coaching*. Jakarta: Dirjen Dikti Proyek Pengembangan Lembaga Pendidikan Tenaga Kependidikan.
- Hastuti, P. 2018. *Genetika Obesitas*. Yogyakarta: Gadjah Mada University Press.

- He, X., Li, Z., Tang, X., Zhang, L., Wang, L., He, Y., Jin, T. and Yuan, D. 2018. Age-and sex-related differences in body composition in healthy subjects aged 18 to 82 years. *Medicine (United States)*, 97(25). pp. 12–17. <https://doi.org/10.1097/MD.00000000000011152>.
- Heymsfield, S. B. 2008. Development of imaging methods to assess adiposity and metabolism. *International Journal of Obesity*, 32. pp. S76–S82. <https://doi.org/10.1038/ijo.2008.242>.
- Higgins, J.K. and Kleimbaun, A.P. 1985. *Design Methodology for randomized clinical trials*. USA: Family Health International. pp. 24-25.
- Hiroshima, Y., Yamamoto, T., Watanabe, M., Baba, Y. and Shinohara, Y. 2018. Effects of cold exposure on metabolites in brown adipose tissue of rats. *Molecular Genetics and Metabolism Reports*, 15. pp. 36–42. <https://doi.org/10.1016/j.ymgmr.2018.01.005>.
- Huh, J.Y., Panagiotou, G., Mougios, V., Brinkoetter, M., Vamvini, M.T., Schneider, B.E., Mantzoros, C.S. 2012. FNDC5 and irisin in humans: I. Predictors of circulating concentrations in serum and plasma and II. mRNA expression and circulating concentrations in response to weight loss and exercise. *Metabolism: Clinical and Experimental*, 61(12). pp. 1725–1738. <https://doi.org/10.1016/j.metabol.2012.09.002>.
- Huh, J. Y., Mougios, V., Kabasakalis, A., Fatourus, I., Siopi, A., Douroudos, I.I., Filippaios, A., Panagiotou, G., Park, K.H. and Mantzoros, C.S. 2014. Exercise-induced irisin secretion is independent of age or fitness level and increased irisin may directly modulate muscle metabolism through AMPK activation. *Journal of Clinical Endocrinology and Metabolism*, 99(11). pp. E2154–E2161. <https://doi.org/10.1210/jc.2014-1437>.
- Huh, J.Y., Siopi, A., Mougios, V., Park, K.H. and Mantzoros, C.S. 2015. Irisin in response to exercise in humans with and without metabolic syndrome. *Journal of Clinical Endocrinology and Metabolism*, 100(3). pp. E453–E457. <https://doi.org/10.1210/jc.2014-2416>.
- Hunn, H. M., Lapuma, P.T. and Holt, D.T. 2002. The Influence of Pre-Test Anxiety, Personality and Exercise on $\text{VO}_{2\text{max}}$ Estimation. *JEPonline Journal of Exercise Physiologyonline*, 5(1). pp. 5–14.
- Jackson, A.S., Stanforth, P.R., Gagnon, J., Rankinen, T., Leon, A.S., Rao, D.C., Skinner, J.S., Bouchard, C. and Wilmore, J.H. 2002. The effect of sex, age and race on estimating percentage body fat from body mass index: The Heritage Family Study. *International Journal of Obesity*, 26(6). pp. 789–796. <https://doi.org/10.1038/sj.ijo.0802006>.
- Kajimura, S., Spiegelman, B.M. and Seale, P. 2015. Brown and beige fat: Physiological roles beyond heat generation. *Cell Metabolism*, 22(4). pp. 546–559. <https://doi.org/10.1016/j.cmet.2015.09.007>.

- Kaminsky, L.A. and Dwyer, G. 2006. Body Composition. In: Kaminsky LA, editor. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*. 5th ed. Philadelphia: Lippincott Williams & Wilkins. pp. 195–205.
- Karlsen, T., Aamot, I.L., Haykowsky, M. And Rognmo, Ø. 2017. High Intensity Interval Training for Maximizing Health Outcomes. *Progress in Cardiovascular Diseases*, 60(1). pp. 67–77. <https://doi.org/10.1016/j.pcad.2017.03.006>.
- Kaupužs, A. 2013. The Relationship Between Physical Activity and Exercise Motivation of the First Year. *Regulation*, pp. 3–15.
- Keil, R. 2004. Coping and stress: a conceptual analysis. *Journal of Advanced Nursing*, 45(6). pp. 659-665. <https://doi.org/10.1046/j.1365-2648.2003.02955.x>.
- Kelly, T., Yang, W., Chen, C.S., Reynolds, K. and He, J. 2008. Global burden of obesity in 2005 and projections to 2030. *International Journal of Obesity*, 32(9). pp. 1431–1437. <https://doi.org/10.1038/ijo.2008.102>.
- Ketel, I.J.G., Volman, M.N.M., Seidell, J.C., Stehouwer, C.D.A., Twisk, J.W. and Lambalk, C.B. 2007. Superiority of skinfold measurements and waist over waist-to-hip ratio for determination of body fat distribution in a population-based cohort of Caucasian Dutch adults. *European Journal of Endocrinology*, 156(6). pp. 655–661. <https://doi.org/10.1530/EJE-06-0730>.
- Khan, S. and Shah, A. J. 2017. Causes and Complication of Obesity among the Children International Journal of Nutrition and Health Sciences Causes and Complication of Obesity among the Children. *Acta Scientific Nutritional Health*, 1(1). pp. 47–50.
- Kibria, G. M. 2019. Prevalence and factors affecting underweight, overweight and obesity using Asian and World Health Organization cutoffs among adults in Nepal: Analysis of the Demographic and Health Survey 2016. *Obesity Research and Clinical Practice*, 13(2). pp. 129–136. <https://doi.org/10.1016/j.orcp.2019.01.006>.
- Kilgore, L. 2010. Adaptation for Fitness. *Cross Fit Journal*, pp. 1–6.
- Koch, A. J. 2010. Immune Response to Exercise. *Journal of Biomotricity*, 4(2). pp. 92–103.
- Koutedakis, Y., Metsios, G.S. and Stavropoulos-Kalinoglou, A. 2006. *The Physiology of Training “Chapter 1. Periodization of Exercise Training in Sport”*. Philadelphia: Chuchill Livingstone Elsevier. pp. 1-21.

- Kraemer, R.R., Shockett, P., Webb, N.D., Shah, U. and Castracane, V.D. 2014. A transient elevated irisin blood concentration in response to prolonged, moderate aerobic exercise in young men and women. *Hormone and Metabolic Research*, 46(2). pp. 150–154. <https://doi.org/10.1055/s-0033-1355381>.
- Kuczmarski, R. J., Fanelli, M. T. and Koch, G. G. 1987. Ultrasonic assessment of body composition in obese adults: Overcoming the limitations of the skinfold caliper. *American Journal of Clinical Nutrition*, 45(4). pp. 717–724. <https://doi.org/10.1093/ajcn/45.4.717>.
- Kuo, T., Chen, T.C., Lee, R.A., Nguyen, N.H.T., Broughton, A.E., Zhang, D. and Wang, J.C. 2017. PIK3R1 is required for glucocorticoid-induced perilipin 1 phosphorylation in lipid droplet for adipocyte lipolysis. *Diabetes*, 66(6). pp. 1601–1610. <https://doi.org/10.2337/db16-0831>.
- Kurdiova, T., Balaz, M., Vician, M., Maderova, D., Vlcek, M., Valkovic, L., Srbecky, M., Imrich, R., Kyselovicova, O., Belan, V., Jelok, I., Wolfrum, C., Klimes, I., Krssak, M., Zemkova, E., Gasperikova, D., Ukropec, J. and Ukropcova, B. 2014. Effects of obesity, diabetes and exercise on *Fndc5* gene expression and irisin release in human skeletal muscle and adipose tissue: *in vivo* and *in vitro* studies. *The Journal of Physiology*, 592(5). pp. 1091–1107. <https://doi.org/10.1113/jphysiol.2013.264655>.
- Kyle, U. G., Bosaeus, I., De Lorenzo, A.D., Deurenberg, P., Elia, M., Giomez, J.M., Heitmann, B.L., Kent-Smith, L., Melchior, J.C., Pirllich, M., Scharfetter, H., Schols, A.M.W.J. and Pichard, C. 2004. Bioelectrical impedance analysis - Part I: Review of principles and methods. *Clinical Nutrition*, 23(5). pp. 1226–1243. <https://doi.org/10.1016/j.clnu.2004.06.004>.
- Lamb, D.R. 1984. *Physiology of Exercise*. New York: Macmillan Pub. pp. 39-63.
- Lim, J. U. Lee, J.H., Kim, J.S., Hwang, Y.I., Kim, T.H., Lim, S.Y., Yoo, K.H., Jung, K.S., Kim, Y.K. and Rhee, C.K. 2017. Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients. *International Journal of COPD*, 12. pp. 2465–2475. <https://doi.org/10.2147/COPD.S141295>.
- Liu, P.Z. and Nusslock, R. 2018. Exercise-Mediated Neurogenesis in the Hippocampus via BDNF. *Frontiers in Neuroscience*, 12. pp. 1–6. <https://doi.org/10.3389/fnins.2018.00052>.
- Lubis, J. 2013. *Panduan Praktis Penyusunan Program latihan*. Jakarta: PT. Rajagrafindo Persada.
- Maalouf, G-E. and Khoury, D.E. 2019. Exercise-Induced Irisin, the Fat Browning Myokine, as a Potential Anticancer Agent. *Journal of Obesity*, Volume 2019, Article ID 6561726, 8 pages. <https://doi.org/10.1155/2019/6561726>.

- Mahajan, R.D. and Patra, S.K. 2013. Irisin, a novel myokine responsible for exercise induced browning of white adipose tissue. *Indian Journal of Clinical Biochemistry*, 28(1). pp. 102–103. <https://doi.org/10.1007/s12291-012-0255-2>.
- Malnick, S.D.H. and Knobler, H. 2006. The medical complications of obesity. *QJM: An International Journal of Medicine*, 99(9). pp. 565–579. <https://doi.org/10.1093/qimed/hcl085>.
- Marie, Ng., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E.C. Biryukov, S., Abbafati, C., Abera, S.F. et al. 2014. Global regional and national prevalence of overweight and obesity in children and adults 1980-2013: A systematic analysis. *Lancet*, 384(9945). pp. 766–781. [https://doi.org/10.1016/S0140-6736\(14\)60460-8.Global](https://doi.org/10.1016/S0140-6736(14)60460-8.Global).
- Mazurek, K., Krawczyk, K., Zmijewski, P., Norkowski, H. and Czajkowska, A. 2014. Effects of aerobic interval training versus continuous moderate exercise programme on aerobic and anaerobic capacity, somatic features and blood lipid profile in collegiate females. *Annals of Agricultural and Environmental Medicine*, 21(4). pp. 844–849. <https://doi.org/10.5604/12321966.1129949>.
- McArdle, W. D., Katch, F.I. and Katch, V.L. 2010. Exercise Physiology. Nutrition, Energy and Puman Performance Seventh Edition. *Lippincott Williams & Wilkins*.
- Moreno-Navarrete, J. M., Ortega, F., Serrano, M., Guerra, E., Pardo, G., Tinahones, F., Ricart, W. and Fernandez-Real, J.M. 2013. Irisin is expressed and produced by human muscle and adipose tissue in association with obesity and insulin resistance. *Journal of Clinical Endocrinology and Metabolism*, 98(4). pp. 769–778. <https://doi.org/10.1210/jc.2012-2749>.
- Moreno-Navarrete, J.M. and Fernández-Real, J.M. 2019. The complement system is dysfunctional in metabolic disease: Evidences in plasma and adipose tissue from obese and insulin resistant subjects. *Seminars in Cell and Developmental Biology*, 85. pp. 164–172. <https://doi.org/10.1016/j.semcdb.2017.10.025>.
- Murawska-Cialowicz, E., Wojna, J. and Zuwala-Jagiello, J. 2015. Crossfit training changes brain-derived neurotrophic factor and irisin levels at rest, after wingate and progressive tests and improves aerobic capacity and body composition of young physically active men and women. *Journal of Physiology and Pharmacology*, 66(6). pp. 811–821.
- Nagai, M., Komiya, H., Mori, Y., Ohta, T., Kasahara, Y. and Ikeda, Y. 2008. Development of a New Method for Estimating Visceral Fat Area with Multi-Frequency Bioelectrical Impedance. *The Tohoku Journal of Experimental Medicine*, 214(2). pp. 105–112. <https://doi.org/10.1620/tjem.214.105>.

- NCD Risk Factor Collaboration. 2016. Trends in adult body-mass index in 200 countries from 1975 to 2014: A pooled analysis of 1698 population-based measurement studies with 19.2 million participants. *The Lancet*, 387(10026). pp. 1377–1396. [https://doi.org/10.1016/S0140-6736\(16\)30054-X](https://doi.org/10.1016/S0140-6736(16)30054-X).
- Nimptsch, K., Konigorski, S. and Pischon, T. 2019. Diagnosis of obesity and use of obesity biomarkers in science and clinical medicine. *Metabolism: Clinical and Experimental*. 92. pp. 61–70. <https://doi.org/10.1016/j.metabol.2018.12.006>.
- Norheim, F., Langleite, T.M., Hjorth, M., Holen, T., Kielland, A., Stadheim, H.K., Gulseth, H.L., Birkeland, K.I., Jensen, J. and Drevon, C.A. 2014. The effects of acute and chronic exercise on PGC-1 α , irisin and browning of subcutaneous adipose tissue in humans. *FEBS Journal*, 281(3). pp. 739–749. <https://doi.org/10.1111/febs.12619>.
- Norton, K., Norton, L. and Sadgrove, D. 2010. Position statement on physical activity and exercise intensity terminology. *Journal of Science and Medicine in Sport*, 13(5). pp. 496–502. <https://doi.org/10.1016/j.jsams.2009.09.008>.
- Novelle, M. G., Contreras, C., Romero-Picó, A., López, M. and Diéguez, C. 2013. Irisin, two years later. *International Journal of Endocrinology*, 2013. <https://doi.org/10.1155/2013/746281>.
- Nygaard, H., Nygaard, H., Slettalokken, G., Vegge, G., Hollan, I., Whist, J.E., Strand, T., Ronnestad, B.R. and Ellefsen, S. 2015. Irisin in blood increases transiently after single sessions of intense endurance exercise and heavy strength training. *PLoS ONE*, 10(3). pp. 1–12. <https://doi.org/10.1371/journal.pone.0121367>.
- Osawa, Y., Azuma, K., Tabata, S., Katsukawa, F., Ishida, H., Oguma, Y., Kawai, T., Itoh, I., Okuda5, S., Matsumoto, I. 2014. Effects of aerobic interval training versus continuous moderate exercise programme on aerobic and anaerobic capacity, somatic features and blood lipid profile in collegiate females. *Open Access Journal of Sports Medicine*, 2014(5). pp. 257–265. <https://doi.org/10.2147/OAJSM.S68932>.
- Palar, C., Wongkar, D. and Ticoalu, S. 2015. Manfaat Latihan Olahraga Aerobik Terhadap Kebugaran Fisik Manusia. *Jurnal e-Biomedik*, 3(1). pp. 316–321.
- Parikh, R., Mathai, A., Parikh, S., Sekhar, G.C., Thomas, R. 2008. Understanding and using sensitivity, specificity and predictive values. *Indian Journal of Ophthalmology*, 56(1). pp. 45–50.

- Parker, L., Parker, L., Shaw, C.S., Banting, L., Levinger, I., Hill, K.M., McAinch, A.J. and Stepto, N.K. 2017. Acute low-volume high-intensity interval exercise and continuous moderate-intensity exercise elicit a similar improvement in 24-h glycemic control in overweight and obese adults. *Frontiers in Physiology*, 7. pp. 1–13. <https://doi.org/10.3389/fphys.2016.00661>.
- Perakakis, N., Triantafyllou, G.A., Fernández-Real, J.M., Huh, J.Y., Park, K.H., Seufert, J. and Mantzoros, C.S. 2017. Physiology and role of irisin in glucose homeostasis. *Nature Reviews Endocrinology*, 13(6). pp. 324–337. <https://doi.org/10.1038/nrendo.2016.221>.
- Peterson, J.M., Mart, R. and Bond, C.E. 2014. Effect of obesity and exercise on the expression of the novel myokines, Myonectin and Fibronectin type III domain containing 5. *PeerJ*, 2. pp. e605. <https://doi.org/10.7717/peerj.605>.
- Pietrobelli, A. and Heymsfield, S.B. 2002. Establishing body composition in obesity. *Journal of Endocrinological Investigation*, 25(10). pp. 884–892. <https://doi.org/10.1007/BF03344052>.
- Powers, S.K. and Howley, E.T. 2015. *Exercise Physiology Theory and Application to Fitness and Performance Tenth Edit.* New York: McGraw-Hill Education.
- Purnomo, M. 2011. Asam Laktat dan Aktivitas SOD Eritrosit pada Fase Pemulihan Setelah Latihan Submaksimal. *Jurnal Media Ilmu Keolahragaan Indonesia*, 1(2). pp. 155–170.
- Qi, G., Zhou, Y., Zhang, X., Yu, J., Li, X., Cao, X., Wu, C., Guo, P. 2019. Cordycepin promotes browning of white adipose tissue through an AMP-activated protein kinase (AMPK)-dependent pathway. *Acta Pharmaceutica Sinica B*, 9(1). pp. 135–143. <https://doi.org/10.1016/j.apsb.2018.10.004>.
- Qiu, S., Bosnyák, E., Treff, G., Steinacker, J.M., Nieß, A.M., Krüger, K., Mooren, F.C., Zügel, M. and Schumann, U. 2018. Acute exercise-induced irisin release in healthy adults: Associations with training status and exercise mode. *European Journal of Sport Science*, 18(9). pp. 1226–1233. <https://doi.org/10.1080/17461391.2018.1478452>.
- Riskesdas. 2018. Laporan Nasional Riset Kesehatan Dasar. Jakarta: Kemenkes RI. Available at: <http://www.kesmas.kemkes.go.id>.
- Robergs, R.A. and Landwehr, R. 2002. The Surprising History of the “HRmax=220-age” Equation. *Journal of Exercise Physiologyonline*, 5(1). pp. 1–10.

- Rodriguez, A.L., Whitehurst, M., Fico, B.G., Dodge, K.M., Ferrandi, P.J., Pena, G., Adelman, A. and Huang, C.J. 2018. Acute high-intensity interval exercise induces greater levels of serum brain-derived neurotrophic factor in obese individuals. *Experimental Biology and Medicine*, 243(14). pp. 1153–1160. <https://doi.org/10.1177/1535370218812191>.
- Rokhmah, F.D., Handayani, D. and Al-Rasyid, H. 2015. Korelasi lingkar pinggang dan rasio lingkar pinggang-panggul terhadap kadar glukosa plasma menggunakan tes toleransi glukosa oral. *Jurnal Gizi Klinik Indonesia*, 12(1). pp. 28–35. <https://doi.org/10.22146/ijcn.22425>.
- Romer, L.M. and Mcconnell, A. 2003. Specificity and Reversibility of Inspiratory Muscle Training. *Medicine and science in sports and exercise*, 35(2). pp. 237–244. <https://doi.org/10.1249/01.MSS.0000048642.58419>.
- Rosella, L.C., Kornas, K., Huang, A., Grant, L., Bornbaum, C. and Henry, D. 2019. Population risk and burden of health behavioral-related all-cause, premature and amenable deaths in Ontario, Canada: Canadian Community Health Survey-linked mortality files. *Annals of Epidemiology*, 32. pp. 49–57.e3. <https://doi.org/10.1016/j.annepidem.2019.01.009>.
- Rosen, E.D. and Spiegelman, B.M. 2014. What we talk about when we talk about fat. *Cell*, 156(1–2). pp. 20–44. <https://doi.org/10.1016/j.cell.2013.12.012>.
- Saleh, O., Majeed, M.J. and Oreaby, G.M. 2014. Descriptive Consideration of Serum Irisin Levels Various Factors: Obesity, Type 2 Diabetes Mellitus, Pre-Diabetic Status, Gender, and Athletics. *Journal of Diabetes & Metabolism*, 5(12). <https://doi.org/10.4172/2155-6156.1000471>.
- Sarifin. 2011. Dampak Faali dari Program Pelatihan (Exercise Program) pada Orang Dewasa. *Jurnal ILARA*, 11(2). pp. 8–14.
- Schumacher, M. A., Chinnam, N., Ohashi, T., Shah, R.S. and Erickson, H.P. 2013. The structure of Irisin reveals a novel intersubunit β -sheet fibronectin type III (FNIII) dimer: Implications for receptor activation. *Journal of Biological Chemistry*, 288(47). pp. 33738–33744. <https://doi.org/10.1074/jbc.M113.516641>.
- Seale, P., Conroe, H.M., Estall, J., Kajimura, S., Frontini, A., Ishibashi, J., Cohen, P., Cinti, S. and Spiegelman, B.M. 2011. Prdm16 determines the thermogenic program of subcutaneous white adipose tissue in mice. *J Clin Invest*, 121(1). pp. 53–56. <https://doi.org/10.1172/JCI44271.96>.
- Selye, H. 1950. Stress and the General Adaptation Syndrome. *Diario Médico Británico*, 1(4667). pp. 1383–1392. <https://doi.org/10.1136/bmj.1.4667.1383>.

- Sergi, G., De Rui, M., Stubbs, B., Veronese, N. and Manzato, E. 2017. Measurement of lean body mass using bioelectrical impedance analysis: a consideration of the pros and cons. *Aging Clinical and Experimental Research*, 29(4). pp. 591–597. <https://doi.org/10.1007/s40520-016-0622-6>.
- Sharp, M.D., Nindl, C., Westphal, K.A. and Friedl, K.E. 1994. The physical performance of female army basic trainees who pass and fail the army body weight and percent fat standards. In: Aghazadeh F, ed. Advances in industrial ergonomics and safety VI. London: Taylor and Francis. pp. 743–50.
- Sillanpää, E., Cheng, S., Hakkinen, K., Finni, T., Walker, S., Pesola, A., Ahtiainen, J., Stenroth, L., Selanne, H. and Sipila, S. 2014. Body composition in 18- to 88-year-old adults - Comparison of multifrequency bioimpedance and dual-energy X-ray absorptiometry. *Obesity*, 22(1). pp. 101–109. <https://doi.org/10.1002/oby.20583>.
- Simanjuntak, R.H., Engka, J.N.A. and Marunduh, S.R. 2016. Pengaruh Latihan Fisik Akut Terhadap Saturasi Oksigen Pada Pemain Basket Mahasiswa Fakultas Kedokteran Unsrat. *Jurnal e-Biomedik*, 4(1). pp. 20–24.
- Stanković, V., Stojanović, S. and Vasiljević, N. 2013. Evaluation of Anthropometric Indices for Metabolic Syndrome and their Association with Metabolic Risk Factors among Healthy Individuals in New Belgrade. *Scientific Journal of the Faculty of Medicine in Niš*, 30(1). pp. 21–30. <https://doi.org/10.2478/v10283-012-0032-4>.
- Sudargo, T., Freitag, H., Rosiyani, F., Kusmayanti, N.A. 2016. *Pola Makan dan Obesitas*. Yogyakarta: Gadjah Mada University Press.
- Sudikno., Syarief, H., Dwiriani, C.M. and Riyadi, H. 2015. Risk Factors of Overweight and Obese in Indonesian Adults (Analysis Data of Basic Health Research 2013). *Journal of The Indonesian Nutrition Association*, 38(2). pp. 91–104.
- Sugiharto. 2000. Pembentukan Radikal Bebas Oksigen Dalam Aktivitas Fisik. *Lab Jurnal Ilmu Keolahragaan dan Pendidikan Jasmani*, 10(1). pp. 22-32.
- Sugiharto. 2003. Adaptasi Fisiologis Tubuh Terhadap Dosis Latihan Fisik. Makalah Disajikan dalam Pelatihan Senam Aerobik. *Jurnal Laboratorium Ilmu Keolahragaan*, Malang: Universitas Negeri Malang.
- Sugiharto and Andiana, O. 2006. The Effects of Exercise Response on Creatine Kinase in White Wistar Rats. *Folia Medica Indonesiana*, 42(4). pp. 228–232.
- Sugiharto. 2012. Fisioneurohormonal pada Stresor Olahraga. *Jurnal Sains Psikologi*, 2(2). pp. 54–66.

- Sugiharto. 2014. *Fisiologi Olahraga: Teori dan Aplikasi Pembinaan Olahraga*. Malang: UM Press.
- Sugiyono, 2017. Metode Penelitian Kuantitatif, Kualitatif dan R&D. Bandung: Alfabeta.
- Sugono, D. 2008. *Kamus Besar Bahasa Indonesia*. Jakarta: Pusat Bahasa Departemen Pendidikan Nasional.
- Sukadiyanto, S. 2010. Stress dan Cara Menguranginya. *Jurnal Cakrawala Pendidikan*, 29(1). pp. 55–66. <https://doi.org/10.21831/cp.v1i1.218>.
- Tew, G.A., Leighton, D., Carpenter, R., Anderson, S., Langmead, L., Ramage, J., Faulkner J., Coleman, E., Fairhurst, C., Seed, M. and and Bottoms, L. 2019. High-intensity interval training and moderate-intensity continuous training in adults with Crohn's disease: A pilot randomised controlled trial. *BMC Gastroenterology*, 19(19). pp. 1–11. <https://dx.doi.org/10.1186/s12876-019-0936-x>.
- The Global Burden of Disease (GBD) 2015 Obesity Collaborators *et al.* 2017. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *New England Journal of Medicine*, 377(1). pp. 13–27. <https://doi.org/10.1056/NEJMoa1614362>.
- Thornton, L.M. and Andersen, B.L. 2006. Psychoneuroimmunology examined: The role of subjective stress. *Cellscience*, 2(4). pp. 66–91.
- Tirtawirya, D. 2012. Intensitas Dan Volume Dalam Latihan Olahraga. *Jurnal Indonesian Sport Scientist Association*, 1(1). pp. 91–99.
- Tjønna, A.R., Lee, S.J., Rognmo, Ø., Stølen, T.O., Bye, A., Haram, P.M., Loennechen, J.P., Al-Sheh, Q.Y., Skogvoll, E., Slørdahl, S.A., Kemi, O.J., Najjar, S.M., and Wisløff, U. 2008. Aerobic Interval Training Versus Continuous ModerateExercise as a Treatment for the Metabolic SyndromeA Pilot Study. *Circulation*, 118(4). pp. 346–354. <https://doi.org/10.1161/CIRCULATIONAHA.108.772822/>.
- Tjønna, A.E., Stølen, T.O., Bye, A., Volden, M., Slørdahl ,S.A., Odegård, R., Skogvoll, E. and Wisløff, U. 2009. Aerobic interval training reduces cardiovascular risk factors more than a multitreatment approach in overweight adolescents. *Clinical Science*, 116(4). pp. 317–326.
- Tohidin, D. 2007. *Adaptasi Fisiologi dalam Olahraga*. Malang: Wineka Media.
- Tschakert, G. and Hofmann, P. 2013. High-Intensity Intermittent Exercise: Methodological and Physiological Aspects. *International Journal of Sports Physiology and Performance*, 8(6). pp. 600–610.

- Tsuchiya, Y., Ando, D., Goto, K., Kiuchi, M., Yamakita, M. and Koyama, K. 2014. High-intensity exercise causes greater irisin response compared with low-intensity exercise under similar energy consumption. *The Tohoku journal of experimental medicine*, 233(2). pp. 135–40. <https://doi.org/10.1620/tjem.233.135>.
- Tsuchiya, Y., Ando, D., Takamatsu, K. and Goto, K. 2015. Resistance exercise induces a greater irisin response than endurance exercise. *Metabolism: Clinical and Experimental*, 64(9). pp. 1042–1050. <https://doi.org/10.1016/j.metabol.2015.05.010>.
- Tsuchiya, Y., Mizuno, S. and Goto, K. 2018. Irisin response to downhill running exercise in humans. *Journal of Exercise Nutrition & Biochemistry*, 22(2). pp. 12–17. <https://doi.org/10.20463/jenb.2018.0011>.
- Vatier, C., Poitou, C. and Clément, K. 2014. *Evaluation of Visceral Fat in Massive Obesity, Nutrition in the Prevention and Treatment of Abdominal Obesity*. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-407869-7.00006-4>.
- Visscher, T.L.S., Snijder, M.B. and Seidell, J.C. 2010. *Epidemiology: Definition and Classification of Obesity, Clinical Obesity in Adults and Children*. Edited by W.H.D. Peter G. Kopelman, Ian D. Caterson. <https://doi.org/10.1002/9781444307627.ch1>.
- Waki, M., Kral, J.G., Mazariegos, M., Wang, J., Pierson, R.N. and Heymsfield, S.B. 1991. Relative expansion of extracellular fluid in obese vs. nonobese women. *American Journal of Physiology-Endocrinology and Metabolism*, 261(2). pp. E199–E203. <https://doi.org/10.1152/ajpendo.1991.261.2.e199>.
- Wardle, J. 1995. The Assessment of Obesity: Theoretical Background and Practical Advice. *Behav. Res. Ther.*, 33(I). pp. 107–117.
- Wen, C. P., Cheng, T.Y.D., Tsai, S.P., Chan, H.T., Hsu, H.L., Hsu, C.C. and Eriksen, M.P. 2008. Are Asians at greater mortality risks for being overweight than Caucasians Redefining obesity for Asians. *Public Health Nutrition*, 12(4). pp. 497–506. <https://doi.org/10.1017/S1368980008002802>.
- Wewege, M., van den Berg, R., Ward, R.E. and Keech, A. 2017. The effects of high-intensity interval training vs. moderate-intensity continuous training on body composition in overweight and obese adults: a systematic review and meta-analysis. *Obesity Reviews*, 18(6). pp. 635–646. <https://doi.org/10.1111/obr.12532>.
- WHO. 2018. *Noncommunicable Diseases Country Profiles 2018*, World Health Organization. <https://doi.org/10.1620/j.ad.2010.09.007>.

- Windarti, N., Hlaing, S.W. and Kakinaka, M. 2019. Obesity Kuznets curve: international evidence. *Public Health*, 169. pp. 26–35. <https://doi.org/10.1016/j.puhe.2019.01.004>.
- Winn, N. C., Grunewald, Z.I., Liu, Y., Heden, T.D., Nyhoff, L.M. and Kanaley, J.A. 2017. Plasma irisin modestly increases during moderate and high-intensity afternoon exercise in obese females. *PLoS ONE*, 12(1). pp. 1–12. <https://doi.org/10.1371/journal.pone.0170690>.
- World Health Organization (WHO). 2016. *Obesity and overweight*. Diakses 04 Januari 2019 dalam <http://www.who.int/mediacentre/factsheets/fs311/en/>.
- Wrann, C.D. 2015. FNDC5/Irisin – Their Role in the Nervous System and as a Mediator for Beneficial Effects of Exercise on the Brain. *Brain Plasticity*, 1. pp. 55–61. <https://doi.org/10.3233/BPL-150019>.
- Xu, B. 2013. BDNF (I) rising from Exercise. *Cell Metabolism*, 18(5). pp. 612–614. <https://doi.org/10.1016/j.cmet.2013.10.008>.
- Yang, M. Wei, D., Mo, C., Zhang, J., Wang, X., Han, X., Wang, Z. and Xiao, H. 2013. Saturated fatty acid palmitate-induced insulin resistance is accompanied with myotube loss and the impaired expression of health benefit myokine genes in C2C12 myotubes. *Lipids in Health and Disease*, 12(1). <https://doi.org/10.1186/1476-511X-12-104>.
- Yu, J. Zhang, S., Cui, L., Wang, W., Na, H., Zhu, X., Li, L., Xu, G., Yang, F., Christian, M. and Liu, P. 2015. Lipid droplet remodeling and interaction with mitochondria in mouse brown adipose tissue during cold treatment. *BBA - Molecular Cell Research*, 1853(5). pp. 918–928. <https://doi.org/10.1016/j.bbamcr.2015.01.020>.
- Zeng, Q., Dong, S.Y., Sun, X.N., Xie, J. and Cui, Y. 2012. Percent body fat is a better predictor of cardiovascular risk factors than body mass index. *Brazilian Journal of Medical and Biological Research*, 45(7). pp. 591–600. <https://doi.org/10.1590/S0100-879X2012007500059>.
- Zhao, J., Su, Z., Qu, C. and Dong, Y. 2017. Effects of 12 weeks resistance training on serum irisin in older male adults. *Frontiers in Physiology*, 8(171). pp. 1–4. <https://doi.org/10.3389/fphys.2017.00171>.
- Zoico, E., Rubele, S., De Caro, A., Nori, N., Mazzali, G., Fantin, F., Rossi, A. and Zamboni, M. 2019. Brown and Beige Adipose Tissue and Aging. *Frontiers in Endocrinology*, 10. pp. 1–10. <https://doi.org/10.3389/fendo.2019.00368>.