

DAFTAR PUSTAKA

- Ampulembang, J. (2004) Hubungan Pajanan Pelarur Organik Metil Etil Keton terhadap Timbulnya Gejala Dini Neurotoksik pada Pekerja di Perusahaan (dengan menggunakan kuesioner Swedish 16). Tesis. Universitas Indonesia, Depok.
- Anselm, B., Gamberale, F. and Iregren, A. (1985) Coexposure to toluene and p-xylene in man: central nervous functions, Br. J. Ind. Med. 42, 117
- Annisa. 2014. Pengaruh Iklim Kerja Terhadap Dehidrasi Pada Karyawan Unit Workhsop PT. Indo Acidatama Karanganyar. *Skripsi*. Surakarta: Universitas Muhammadiyah Surakarta.
- Agency of Toxic Substance and Disease Registry, ATSDR. (2000), Toxicology Profile of Xylene. www.atsdr.cdc.gov/toxprofile/tp.asp?id=234&tid=42.
- Agency for Toxic Substances and Disease Registry (ATSDR). 2002. Toxicological profile for Nitrogen Oxides (nitric oxide, nitrogen dioxide, etc). Atlanta GA: U.S. Department of Public Health and Human Services. Public Health Service.
- Arnold, SM, Angerer, J, Boogaard, PJ, Hughes, MF, O'Lone, RB, Robinson, SH, Schanatter, AR. (2013): The Use of biomonitoring data in exposure and human health risk assessment: benzene case study. Critical Review in Toxicology, 43 (2): 119-153.
- Asilian, H., Nejad, M., Mortazavi, B., Jafari, M.J., Khavanin, A.R., Dehsdasti, A.R. (2008) Assessment of Bagging Operators Exposure to with PVC Airborn Particulates. Iran Journal Environ. Sci. Eng. Vol 5 No. 3. 187-194.
- Astrand, I (1982) Work load and uptake of solvents in tissues of man. In: Mehlman, M A, ed., Advances in Modern Environmental Toxicology. Vol. 2. Princeton Junction, NJ: Senate Press; pp.141-152.
- Astrand, I; Engström, J; Övrum, P (1978) Exposure to xylene and ethylbenzene. I. Uptake, distribution, and elimination in man. Scand J Work Environ Health. 4:185-194.
- ATSDR. Agency for Toxic Substances and Disease Registry. 2005. Toxicological profiles for benzene (Draft for Public Comment). U.S. Department of Health and Human Services, Public Health Service, Atlanta, Georgia, U.S.A.
- ATSDR Agency for Toxic Substance and Disease Registry. 2007. ToxGuide for Benzene C₆H₆. U.S. Department of Health and Human Services, Division of Toxicology and Environmental Medicine Applied Toxicology Branch.

- Agustina, U. (2016). Correlation between Toluene and Urine Hippuric Acid with Central Nervous System Complaints at Workshop Worker. Essay. Surabaya: Faculty of Public Health, Airlangga University.
- Baker, Jr. and Seppäläinen, A.M. (2007) Prevention of neurotoxic illness in working populations. American Journal of Industrial Medicine. Vol. 15, Issue 1, pages 117–118, 2007.
<http://onlinelibrary.wiley.com/doi/10.1002/ajim.v15:1/issuetoc>.
- Badjagbo, K., Loranger, S., Moore, S., Tardif, R., Savre, 2010. BTEX Exposures Among Automobile Mechanics and Painter and Their Associated Health Risks, Human and Ecological Risk Assessment : An International Journal, vol 16, issue 2, p. 301 – 316.
- Bergman K. (1979) Application and results of whole-body autoradiography in distribution studies of organic solvents. CRC Crit Rev Toxicol 12(1):59-118.
- Bloemen, L. J. et al. (2004) ‘Lymphohaematopoietic cancer risk among chemical workers exposed to benzene’, *Occupational and Environmental Medicine*, 61(3), pp. 270–274. doi: 10.1136/oem.2003.007013.
- Bustan, M.N. (2000).Epidemiologi Penyakit Tidak Menular. PT. Rineka Cipta, Jakarta.
- Cassaret and Dousl, (2008) Toxicology : The Basic Science of Poisons 7th edition , Mc Graw Hill Medical Publishing Division, USA.
- Centres for Disease Control and Prevention (CDC). (2012) International Chemical Safety Cards (ICSC) :p-Xylene w.cdc.gov/niosh/ipcsneng/neng0086.htm.
- Clayton G, (1994), *Patty's Industrial Hygiene and Toxicology* vol 1 part B, New York
- Davidson, R.D, Courage, C, Rushton, L, Levy, L. (2001): Benzene In The Environment: An Assessment Of The Potential Risks To The Health Of The Population. Occup Environ Med 2001; 58:2–13.
- Dick, FD, (2006) Solvent neurotoxicity. Occup Environ Med. 2006 Mar; 63(3): 221– 226.<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2078137>.
- Dosing, M., Aelum, JB., Hansen, SH., Lundqvist, GR., and Andersen, N. (1983) Urinary Hippuric Acid and OrthocresolEscretion in Man during Experimental Exposure to Toluene. Bri J Ind Med (40)pp470-473.
- Effendi, Z. 2003. Peranan Leukosit Sebagai Anti Inflamasi Alergik dalam Tubuh. Sumatera Utara: Bagian Histologi Fakultas Kedokteran Universitas

Sumatera Utara

- Eiselen, SC, (2006) Neuropsychological Toxicology: A Theoretical Overview of Neuropsychological Assessment. Mini-Dissertation. Department of Psychology, University of Pretoria. South Africa. <http://repository.up.ac.za/xmlui/bitstream/handle/2263/28742/dissertation.pdf?sequence=1&isAllowed=y>.
- EHS. (2002). Toxicology And Exposure Guidelines. UNL Environmental Health and Safety University of Nebraska Lincoln.
- EPA. 2003. *A Standardized EPA Protocol For Characterization Indoor Air Quality In Large Office Building*.
- Environmental Protection Agency (EPA). (2011) Integrated Risk information System, <http://www.epa.gov/IRIS/subst/0486.htm>.
- Environmental Protection Agency (EPA). (2008) Technology Transfer Network Air Toxic. <http://www.epa.gov/ttn/atw/hithef/n-hexane.htm>.
- Erwinanto, d. (2013). PEDOMAN TATALAKSANA DISLIPIDEMIA. Jurnal Kardiologi Indonesia, 1.
- Faradisha, Jihan. (2018). Analisis Hubungan Paparan Toluena Dengan Risiko Neurotoksik Pada Pekerja Bagian Printing Industri Karung Plastik Di Pt X Sidoarjo. *Tesis*. Surabaya : Universitas Airlangga.
- Gamble, J. (2000). Low-level hydrocarbon solvent exposure and neurobehavioural effects. *Occupational Medicine* 50:81-102.
- Guyton, AC. (2007) Fisiologi Kedokteran, 148 - 168, Edisi ke 12, EGC, Jakarta
- Hamm, R.D. (2013) The Nervous System – Target Organ into the Twenty-First Century. Canadian Board of Occupational Medicine. Canada. <http://spph.ubc.ca/files/2013/10/Occupational-NeuroToxicology-by-Dr.-R.-D.-Hamm.pdf>.
- Harefa, E. 2011. HbA1c Standardization and Recent Updates. Prodia Laboratories. Makassar.
- Health Protection Agency (HPA), (2010) *Xylene: General Information*. CRCE HQ, HPA Version 1. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/318006/hpa_Xylene__General_Information_v1.pdf. diakses pada 10 September 2015.
- Hoffbrand, J.E. Petit, et al. 2005. Kapita Selekta Hematologi. Edisi 4. Penerbit Buku Kedokteran EGC. Jakarta

- Ihrig, A., Triebig, G., Dietz, M. (2001). Evaluation of a Modified German Version of The Q16 Questionnaire for neurotoxic symptoms in workers exposed to International Agency for Research on Cancer , IARC, (2011) Overall Evaluation of Carcinogenicity to Humans List of Agents, Mixtures and Exposures Evaluated. Monographs.iarc.fr/ENG/classification.
- Irawati, Yana. 2010. *Analisis Resiko Kesehatan Pajanan Xylene pada Pekerja Bengkel Sepatu "X" di Kawasan Perkampungan Industri Kecil (PIK) Pulogadung Jakarta Timur*. Jakarta: Universitas Indonesia (Tesis)
- IPCS (2004) *Xylene*. Geneva, World Health Organization, International Programme on Chemical Safety (International Chemical Safety Card 0015;www.inchem.org/documents/icsc/icsc/eics0015.htm.
- Jacobson, GA, McLean S. (2003) Biological Monitoring of Low Level Occupational Xylene Exposure and the Role of Recent Exposure. Oxford Journals Medicine & Health The Annals of Occupational Hygiene Volume 47, Issue 4Pp. 331-336. www.ncbi.nlm.nih.gov/pubmed/12765874.
- Jafari, MJ., Karimi, A., and Azari, MR. (2009) The Challenges of Controlling Organic Solvents in a paint Factory due to Solvent Impurity. Industrial
- Kandayala R., Raghavendra S., Rajasekharan S. (2010). *Xylene*: An Overview of its health hazards and preventive measures. J of Oral and Maxillofacial Pathology. 2010;14(1):1-5.<http://www.jomfp.in/article.asp?issn=0973-029X;year=2010;volume=14;issue=1;spage=1;epage=5;aulast=Kandyala>.
- Kawai T, (1991) Urinary methylhippuric acid isomer levels after occupational exposure to a xylene mixture. Int Arch Occup Environ Health. 1991;63(1):69-75.www.ncbi.nlm.nih.gov/pubmed/1856026.
- Kolluru, R.V.1996. *Risk Assessment and Management Handbook*. New York : McGraw Hill Inc.
- Lamm, A., Savolainen, K., Matikainen, E., (2003) Acute Effects of m-xylene Inhalation on Body Sway, Reaction Times, and Sleep in Male Mice. Int Arch Occup Environ Health 65:179-188.
- Lesitari. 2006. Hubungan Pelarut Organik Dengan Gejala Neurotoksik Pada Pekerja Alas Kaki Di Sektor Informal Ciomas Bogor. (Menggunakan Kuesioner Swedish Q16. *Thesis*. Jakarta: Fakultas Kedokteran Universitas Indonesia.
- Lestari F, (2010) Bahaya Kimia : Sampling & pengukuran kontaminan kimia di udara, Penerbit Buku Kedokteran. Jakarta.
- Louvar,J.F and Louvar, B.D. (1998). *Health and Environment Risk Analysis*:

- Fundamental With Applications.* Voluem 2. New Jersey, Prentice Hall PTR
- Lu, FC. (2006) Toksikologi Dasar: Asas, Organ Sasaran dan Penilaian Risiko. Edisi ke-2. UI Press. Jakarta
- Lundberg I. (1986) Correlation of *xylene* exposure and methyl hippuric acid excretion in urine among paint industry workers. Scand J Work Environ Health. 1:149-53. <http://www.ncbi.nlm.nih.gov/pubmed/3726497>.
- Malathi N, Sharada T, Rajan, N. (2014). Health Hazards of *Xylene*: A Literature Review. J Clin Diagn Res. 2014 Feb;8(2):271-4. doi: 10.7860/JCDR/2014/7544.4079.<http://www.ncbi.nlm.nih.gov/pubmed/24701554>.
- Mirta, DR. (2012) Analisis Gejala Neurotoksi Akibat Pajanan Pelarut Organik *Xylene* pada Pekerja Pembuat Cat PT X Tahun 2012. Tesis. Universitas Indonesia. Jakarta.
- Mulhausen,J and Damiano (2003) Chapter 4th ; Comprehensive Exposure Assessment the Occupational Environment : Its evaluation, control, and management. 2nd edition.
- Murray and Lopez. (2006) Mortality by Cause for 8 region of the world: Global Burden of Disease, 2006, June 9, <http://www.thelancet.com/journal/vol349/iss9062/full/llan.349.9061.original.research.8645.1>. Diakses pada 23 Mei 2015.
- Mohamed, M.Y.A. (2008) Risk Assessment of Hazardous Chemicals A Case Study in Chemistry Research Laboratory of USM. Thesis. University Sains Malaysia.
- Morin M, Chambon P, Chambon R, Bichet N. (1986) Measurement of exposure to *xlyenes* by separate determination of m- and p-methylhippuric acids in urine. J Chromatogr. 1981 Jun 5;210(2):346-9. <http://www.ncbi.nlm.nih.gov/pubmed/7263793>.
- National Research Council, NRC (2010) Keselamatan dan Keamanan laboratorium Kimia : Panduan pengelolaan Bahan Kimia dengan Bijak. The National Academy Press, Washington DC.
- NIOSH. (1996) Volatile Organic Compounds (Screening). www.cdc.gov/niosh/docs/154/pdfs/1005.pdf.
- NIOSH. National Institute for Occupational Safety and Health. 2003. Manual of Analytical Methods (NMAM), Fourth Edition
- Occupational Safety and Health Division, OSHD, (2004) A Semi Quantitative to Assesss Occupational Exposure to Harmful Chemical. The Ministry of

Manpower. Havelock Road Singapore.

- Ogata, M; Tomokuni, K; Takatsuka, Y (1970) Urinary excretion of hippuric acid and m- or pmethylhippuric acid in the urine of persons exposed to vapours of toluene and m- or p-xylene as a test of exposure. *Brit J Indust Med.* 27:43-50.
- Palmer, A. dan Williams, B. 2007. Simple Guides Tekanan Darah Tinggi. EGC. Jakarta
- Pearce, EC. (2006) Anatomi dan Fisiologi untuk Paramedis. Jakarta: Gramedia Pustaka Utama
- Pennline, C. (2005) Evaluating Potential Exposure to Hexavalent Chromium (Cr(VI)) among Maintenance Mechanicwelders. Thesis. University of Cinncinati.<http://etd.ohiolink.edu/view.cgi?acc-num=ucin1182717517>.
- Rahmah, M. D. (2012). Analysis neurotoxic Symptoms caused by Xylene Organic Solvent Exposure at Paint Making Workers. Thesis. Jakarta: Master Program of Occupational Health and Safety, Faculty of Public Health, University of Indonesia. [In Indonesian]
- Rahmatullah Maista, Naiem M. Furqan, Muis Masyitha. (2013). Keamanan Bahan Kimia Berbahaya pada Karyawan Percetakan Kota Makasar. Bagian Kesehatan dan Keselamatan Kerja FKM UNHAS, Makasar.
- Recchia, G; Perbelline, L; Prati, GF; Dean, P; Ancona, G (1985) Coma da prbabile ingestione accidentale di xilene: trattamento mediante emoperfusione con carbone attivato. *Med Lav.* 76: 67-73.
- Riihimäki, V., and Savolainen K. (1980) Human exposure to m-xylene: Kinetics and acute effects on the central nervous system. *Ann. Occup. Hyg.* 23(4):411-422.
- Ridgway, P., Nixon, T., and Leach, J.-P. (2003). Occupational exposure to organic solvents and long-term nervous system damage detectabel by brain imaging, neurophysiology or histopathology. *Food and Chemical Toxicology* 41:153-187.
- Rini. 2007. Kajian Sick Building Syndrome (Studi Kasus : Sick Building Syndrome pada Gedung X di Jakarta). *Jurnal Teknik Sipil* vol.3, No.2, Oktober 2007, hlm.158-173.
- Ruijten, MW., Hooisma, J. Brons, JT., Habets C., Muijser H. (1994) Neurobehavioral effects of long-term exposure to xylene and mixed organic solvents in shipyard spray painter. *Neurotoxicology*. 1994 Fall;15(3):613-20. <http://www.ncbi.nlm.nih.gov/pubmed/7854596>.

- Sato, A; Nakajima, T (1979) Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. Br J Ind Med.36:231-234.
- Salvatore, AIHA-Washington TLV ACGIH (2010) Treshold Limit Value for Chemicals Substances and Physical Agents Biologica Exposure Indicates, Cinncinati.
- Schnatter, A. R. et al. (1996) *The relationship between low-level benzene exposure and leukemia in Canadian petroleum distribution workers, Environmental Health Perspectives*. doi: 10.2307/3433193.
- Seppiilainen, A.M., Leine A., Salmi T, (2009) Changes Induced by Short-Term Xylene Exposure in Human Evoked Potentials. Int Arch Occup Environ Health 61 : 443-449. diakses pada 30 Agustus 2015.
- Sintayati, E., Oginawati, K. (2012) Hubungan Pajanan Toluen dengan Kadar Asam Hipurat Urin Pekerja Pengecatan Mobil (Studi pada Bengkel Mobil Informal di Karasak, Kota Bandung). Tesis. Institut Teknologi Bandung, Bandung.
- Smit, J., Asnong, W. (2009). Industrial Hygiene Guideline : The Assessment of Exposure Vinyl Chloride and 1,2-Dichloroethane. IndustTox Consult, 49, 735-541.
- Swan GE and Lessov-Schlaggar CN1. (2009) Genetic influences on individual differences in nicotine glucuronidation. Twin Res Hum Genet. 2009 Oct;12(5):507-13. <http://www.ncbi.nlm.nih.gov/pubmed/19803778>. diakses pada 14 September 2015.
- Spurgeon, A., Glass, D.C., Calver, I.A., Cunningham-Hill, M., and Harrington, J.M. (1994). Investigation of dose related neurobehavioral effects in paintmakers exposed to low levels of solvents. Occupational and Environmental Medicine, 51, 626-630.
- Sullivan, JB and Krieger, GR. (1991) Alkyl Benzene Solvent and Aromatic Compouns. In Sullivan, JB and Krieger, GR. (ed). Hazardous Materials Toxicology Clinical Principals of Environmental Health. Sydney. William and Wikins pp 1086-1104. Symptoms and signs in workers exposed predominantly to xylene. Int Arch Occup Environ Health 1993;64:597-605. <http://www.ncbi.nlm.nih.gov/pubmed/8314620>.
- Suma'mur. 2009. Hiegiene Perusahaan dan Keselamatan Kerja. Jakarta : CV Sagung Seto.
- Susiana, Nika. (2016) 'Gambaran Profil Darah Pekerja Sektor Informal Industri Sepatu Kulit (Studi Kasus UD. Kurnia dan UD. Malis Jaya)', *Universitas Negeri Semarang*, pp. 1-71.

- Tsai, SY, Chen, JD., Chao, WY., and Wang, JD., (2007) Neurobehavioral Effects of Occupational Exposure to Low-Level Organic Solvents among Taiwanese Workers in Paint Factories, Environmental Research. 2007; 73,146–155.
<http://www.sciencedirect.com/science/article/pii/S1201393519793704>
- Thetkathuek, A., Jaidee, W., Saowakhontha, S., & Ekburanawat, W. (2015). Neuropsychological symptoms among workers exposed to toluene and xylene in two paint manufacturing factories in eastern Thailand. Advances in preventive medicine, 2015. <https://doi.org/10.1155/2015/183728>.
- Unlu, I., and Kesici, G. (2014). A Comparison of The Effects of Solvent and Noise Exposure on Hearing, Together and Separately. Noise Health. 2014 Nov-Dec;16(73):410-5. doi: 10.4103/1463-1741.144422. <http://www.ncbi.nlm.nih.gov/pubmed/25387537>.
- U.S. EPA . (1996) Reproductive Toxicity Risk Assessment Guidelines. Federal Register 61 (212):56274-56322.
- U.S. EPA . (1998a) Guidelines for Neurotoxicity Risk Assessment. Federal Register 63 (93):26926-26954.
- U.S. EPA . (1998b) Science policy council handbook: peer review. Prepared by the Office of Science Policy, Office of Research and Development, Washington, DC. EPA/100/B-98/001.
- U.S. EPA (1998c) Carcinogenic effects of Benzene: An Update. IRIS Cancer Assessment. Prepared by the National Center for Environmental Assessment. Washington, DC.
- U.S. EPA .(1999) Review Draft Guidelines for Carcinogen Risk Assessment. Review Draft. July 1999.
- U.S. EPA (2000) Toxicological Review of benzene in support of summary information on the Integrated Risk Information System (IRIS). National Center for Environmental Assessment, Washington, DC. Available on line from: <http://www.epa.gov/iris>
- US.EPA, 2002. Aquatic Resources Monitoring. Akses internet: <http://www.epa.gov/nheerl/arm/>
- Vaidyanathan A, Foy JWD, Schatz R. Inhibition of rat respiratory-tract cytochrome P-450 isozymes following inhalation of m-xylene: possible role of metabolites. J Toxicol Environ Health. 2003;66:1133–1143.
- Wang, JD., Chen, JD. (1993) Acute and chronic neurological symptoms among paint workers exposed to mixtures of organic solvents. Environ Res. 1993 Apr;61(1):107-16. <http://www.ncbi.nlm.nih.gov/pubmed/8472664>.

- WHO. (2000). *Air Quality Guidelines - Second Edition Chapter 5.14 Toluene.* Europe. WHO Regional office
- Winder, C. (2005). Occupational Toxicology, 2nd Edition. CRC Press, Taylor & Francis e -Library . France.
- Winder, C. and N. Stacey, 2005. Toxicity of organic solvents, in Occupational Toxicology, Second edition, Winder and Stacey (ed). CRC press, 14: 364-388.
- World Health Organization. The world health report 1996: fighting disease-fostering development, Geneva: WHO; 1996. P. 137.
- WHO. (1997) Environmental health criteria 190. *Xylene.* World Health Organization. Geneva, Switzerland.
- WHO. (1986) Early Detection of Occupational Disease. World Health Organization. Geneva, Switzerland.
- Worksafe Western Australia Commision (WWAC).2009. Code of Practice Spray Painting, Worksafe Western Australia.
- Yatim, 2010, Cara ampuh mengontrol kolesterol. Sarana Pustaka Medika: Jakarta