

11. ROLE OF HEPATITIS B AND HEPATITS C VIRUS COINFECTION IN TB DEVELOPMENT _ Lestari _ Folia Medica Indonesiana *by Pudji Lestari*

Submission date: 24-Sep-2019 11:59AM (UTC+0800)

Submission ID: 1178840124

File name: 5204-15192-1-SM.pdf (172.94K)

Word count: 3245

Character count: 16941

1
ROLE OF HEPATITIS B AND HEPATITIS C VIRUS COINFECTION IN TB DEVELOPMENT

Pudji Lestari, Ivan Rahmatullah, Dwi Susanti
Department of Public Health and Preventive Medicine
Faculty of Medicine, Universitas Airlangga

ABSTRAK

Perkembangan penyakit Tuberkulosis (TB) pada tiap individu berbeda. Peran penyakit koinveksi seperti HIV cukup significant namun peran penyakit lain seperti Hepatitis B dan Hepatitis C belum banyak diketahui. Oleh karena itu, penelitian ini bermaksud untuk menggambarkan peran koinveksi virus hepatitis dalam perkembangan TB. Metode penelitian adalah observasional – case control dengan dua kelompok. Kelompok pertama adalah penderita TB dan kelompok kedua adalah kelompok kontrol. Data yang dikumpulkan meliputi kondisi TB, apakah laten atau manifest, dan riwayat infeksi Hepatitis B dan C. Penelitian ini berhasil melibatkan total 32 responden pada masing – masing kelompok. Pada kelompok kasus, jumlah responden laki-laki sebanyak 19 orang dan perempuan sebanyak 13 orang. Sementara pada kelompok kontrol, jumlah responden laki-laki sebanyak 10 orang dan perempuan sebanyak 22 orang. Pada kelompok kasus, didapatkan 20 orang responden positif dengan riwayat hepatitis B. Sedangkan pada kelompok kontrol didapatkan 12 orang responden positif. Hasil perhitungan dengan chi-square menunjukkan adanya perbedaan yang signifikan antara kedua kelompok ($P=0.0455$ dan $\alpha=0.05$) dengan odd ratio $p=0.479$. Sementara itu, penelitian ini mendapati hanya satu orang responden yang positif riwayat koinveksi C dan dua orang pada kelompok kontrol. Uji beda pada dua kelompok menunjukkan hasil yang tidak signifikan ($P=1$ dan $\alpha=0.05$). Penelitian ini menunjukkan bahwa penderita tuberkulosis lebih mungkin memiliki riwayat koinveksi virus hepatitis B 2,7 kali dibandingkan kelompok kontrol yang terinfeksi Hepatitis B namun tidak sakit Tuberkulosis. Sehingga, kemungkinan koinveksi virus hepatitis B berperan pada manifestasi penyakit tuberkulosis namun tidak dengan riwayat koinveksi hepatitis C. (FMI 2016;52:19-23)

Kata kunci: hepatitis B, hepatitis C, koinveksi, tuberkulosis

ABSTRACT

Tuberculosis developed differently in each individual. The role of HIV in the disease development is widely known unlike the role of Hepatitis B and Hepatitis C infection. Therefore, this study is aimed at to describe the role of those co-infections in the TB development. The study employed observational – case control design with two groups, case and control group. Data collection include TB condition, whether latent or manifest, and history of Hepatitis B and Hepatitis C co-infection. The study successfully recruited the total of 32 respondents in each group. At the case group, the number of male and respondent is 19 and 13. Meanwhile, at the control group the number male respondent is 10 and the female respondent is 22. The study found that 20 and 12 respondents were positive Hepatitis B at the case and the control group. Chi-square test resulted significant different between two groups ($P=0.0455$ and $\alpha=0.05$) with odd ratio $p=0.479$. Meanwhile, the study only tested one respondent with positive Hepatitis C at case group and one at control group. Comparison analysis found no significant different between groups ($P=1$ and $\alpha=0.05$). This study reveals that TB patients are more likely to have history of Hepatitis B 2.7 times than those at the control group with the same infection but without TB development. Therefore, the possibility of Hepatitis B co-infection has role in the TB manifestation, but not for Hepatitis C co-infection. (FMI 2016;52:19-23)

Keywords: hepatitis B, hepatitis C, coinfection, tuberculosis

2
Correspondence: Pudji Lestari, Department of Public Health and Preventive Medicine, Faculty of Medicine, Universitas Airlangga

INTRODUCTION

Indonesia is included in high-burden countries for Tuberculosis (TB). The number of new cases in 2012 amounted to 202,319 cases. The threat of such magnitude of TB cases is the drop in productivity, because the ratio of illness between males and females was 1 : 1.5 and nearly 90% of the patients aged over 15 years (WHO 2013). Report of Basic Health Research (Risksdas) 2010, found that the point prevalence of TB

(at least one slide) reached 704/100,000, which many higher than the WHO global report of 244/100,000 (MOH 2011). The development of TB disease in each individual is different, whether becomes latent or manifest. However, how cases of infection becomes latent or manifest, remains unclear. More than 90% of cases remain as latent cases, while the rest are developing into disease. Thus there is a need to find other approaches to the identification of group with a risk of illness. Further efforts for more effective

prevention of tuberculosis can therefore be done (Soemantri et al 2007).

Infection of Human Immunodeficiency Virus (HIV) and the development of Acquired Immune Syndrome Deficiency (AIDS) provide new insights into the course of the tuberculosis disease. The HIV virus is known to allow the bacteria *Mycobacterium tuberculosis*, by creating ideal conditions through the suppression of CD4 cells. With the existence of HIV it is known that there is an interaction between viral infection and latent infection of bacteria. The evidence suggests, someone who recovered from TB, TB does not become free, even having higher risk of illness than people who have latent infection. It shows that TB is a disease of opportunism. Indonesia is endemic TB region, the number of cases of latent estimated doubled from the number of TB patients. It is important to identify the factors that play a role in the development of TB that has become a disease as the health concern of TB is its form as a disease, not a latent TB.

Nagelkerke filed a 'double hit' theory to explain the role of co-infection diseases in the development of TB disease. This theory suspect virus coinfection that would cause someone who is already infected develop the disease, and vice versa when it is infected by this virus and then infected with *Mycobacterium tuberculosis* it will manifest. This virus has characteristics such as sexually transmitted or blood through other body fluids. So people with promiscuity (free sex) are more susceptible to infection with this virus (Nagelkerke 2012).

Hepatitis B and C is a viral infection which is also present in high promiscuity groups (eg sex workers and homosexual) and also more common among inject-ing drug abusers. Hepatitis viruses are also threatening infections in transfusion recipients and users of the dialysis machine. Infection with this virus is also a chronic case of the manifestation that will appear 10-20 years later. Thus these viruses are ideal candidates for co-infected with TB. Therefore, this study aims to identify the description of viral hepatitis coinfection role in the development of TB.

MATERIALS AND METHODS

This study was an observational case control study with two groups, the case group and the control group. The case group was adult patients diagnosed with sputum smear-positive are undergoing treatment at a government hospital in Surabaya, while the control group were health workers or family members of patients. The number of each group was 32 adults. The research was conducted in the period from August to October 2014.

Studied variables included the condition TB, whether latent or manifest, and a history of Hepatitis B and C.

xxx

The study began with a feasibility test of ethics which was done by Bioethics Unit, Faculty of Medicine. After eligibility certificate of conduct was established, the researchers proposed permit to conduct research in the government hospital, Surabaya. Once the permission was obtained, the researchers encountered a group of cases that TB patients who already diagnosis of smear positive TB services at the facility. Willingness to undergo serological examination requested form which preceded availability information for standard procedures consent. Meanwhile, a control group matched by age and sex of the patient, found in TB care (family) or in the homes of people (contacts). They requested a willingness to take blood for examination and serological Mantoux test in the consent form. If the observations found that the household contacts become ill with TB, he/she will be removed from the control group.

Procedures of blood collection from all respondents was that as much as 10 cc of blood to be divided into 4 cc of whole blood + EDTA for examination hematology blood, in control group: Mantoux test was checked, while all respondents checked serum or plasma for ELISA against anti-HBV and anti-HCV. Tests done include: Mantoux test, serology examination of anti-HBV and anti-HCV by ELISA, and Characteristics of respondents: age, their illness (diabetes, HIV, and a history of allergies) using a questionnaire.

The data collected will be described descriptively and presented in the table of distribution. Then, an analysis using univariate test for each of the risk of co-infection. Statistical test is done using fisher, chi-square, t-test, and odds ratios. Statistical significance set at $p \leq 0.05$. Data distribution and analysis using appropriate computer software.

RESULTS

This study was able to recruit a number of each person in the group of 32 adults with diagnosed TB patients' sputum smear-positive and control groups. Of the 32 people in the case group, the number of male respondents as many as 19 people and 13 women among as seen in Table 1. Meanwhile, for the control group the number of male respondents was 10 women and 22 people, as seen in table 1.

In the study also found the average age of the respondents in the case group was 40 years with a standard deviation of 11.5 with the oldest age of

respondents was 62 years, while the youngest was 20 years old. In the control group, the average age is 40.9 and the standard deviation of 13.1 with the oldest aged 65 and the youngest 20 years old. Test performed depending on the age of the two groups with t test showed $P = 0.716$ and $\alpha = 0.05$ which means there is no significant difference between the 2 groups. As seen in Table 2

Table 1. Respondents' gender

	Case Group		Control Group	
	Freq.	%	Freq.	%
Male	19	59.4	10	31.3
Female	13	40.6	22	68.8
Total	32	100	32	100

Comparative test with Chi square ; $P = 0.077794$; $\alpha = 0.05$;
No significant difference between two groups

Table 2. Age of respondents

Groups	Case (32 persons)	Control (32 persons)
Mean and SD	40.00 +/- 11.5	40.9 +/- 13.1
Maximum	62	65
Minimum	20	20

Comparative test with t test ; $P = 0.716$; $\alpha = 0.05$; No significant difference between two groups

Test for history of hepatitis B virus in this study in the case group, obtained 20 respondents 12 respondents positive and negative. While the control group got 12 respondents 20 respondents positive and negative. Different test was also carried out between the two groups. The result of the calculation with chi-square shows the results; $P = 0.0455$ and $\alpha = 0.05$ which means there is a significant difference between the two groups with odd ratio $p = 0.479$. As seen in Table 3.

Table 3. History of co-infection with hepatitis B virus

	Case		Control	
	Freq.	%	Freq.	%
Hepatitis B Antibody positive	20	62.5	12	37.5
Hepatitis B Antibody negative	12	37.5	20	62.5
Total	32	100	32	100

Comparative test with chi-square; $P = 0.0455$; $\alpha = 0.05$; Two groups shows significant difference; Odd ratio $P = 0.479$

Test for history of hepatitis C virus in this study, only one person respondent positive in case group. Meanwhile, 31 other responded negatively. In the control group, there were two people with a positive result and 30 people with a negative result. Different test was also carried out between the two groups also performed. Fisher exact test results show results; $P = 1$ and $p=0.05$ which means there is no significant difference between the 2 groups. As seen in Table 4.

Table 4. History of hepatitis C virus coinfection

	Case		Control	
	Freq.	%	Freq.	%
Hepatitis C Antibody positive	1	96.8	2	93.8
Hepatitis C Antibody negative	31	3.1	30	6.3
Total	32	100	32	100

Comparative test with Fisher exacts test ; $P = 1$; $\alpha = 0.05$; No significant difference between two groups

DISCUSSION

3
The results of this study showed that there was no significant difference between the control group and the case group in terms of age and gender. Thus, these two factors can be balanced, and has been controlled in the hope of another variable results in both groups were compared.

In this study, we used the criterion of control is family tuberculosis with sputum positive stay at home for two consecutive years, according to a study Setiawati et al (2011) and Lestari et al (2011), Mantoux positive and quantitation of Interferon (IGRA) actually showed only partial of the infected group (sensitivity 60%).

This study found quite a lot of people with TB also have been exposed to hepatitis infection (especially hepatitis B), OR in this study of 2.7. This means that patients are 2.7 times more likely to have a history of hepatitis B infection, rather than control. This finding is consistent with research findings in China at risk groups of drug users who find an odd ratio was higher in coinfection Hepatitis virus (Zhang et al 2013). Another study, still in China, also found the risk of clinical manifestations of tuberculosis increased in the group with oncogenic HPV infection (Graubard et al 2011). These facts support the hypothesis that virus infection associated with disease manifestations in tuberculosis.

This study found only one person tuberculosis sufferers who also have been exposed to hepatitis C. Meanwhile, the control group did not obtain similar fact. Although the results showed no significant between the role of hepatitis C coinfection TB circuitry, but this study shows a similar trend to the importance of analyzing the role of co-infection in TB patients. The study in China found that the risk of clinical manifestations of tuberculosis increased in the group with oncogenic HPV infection (Graubard et al 2011). These facts support the hypothesis that virus infection associated with disease manifestations in tuberculosis.

Elucidation of the role of coinfection against tuberculosis disease manifestations may be obtained from meta-analysis of Dextran paper written by Pustynnikov

16

et al. Uptake of hepatitis C virus and *Mycobacterium tuberculosis* share in dendritic cell receptor and mannose receptor same (Pustynnikov 2014). Regulatory T cell helper 1 (Th1) and T helper cell 2 starts here. Can speculate that a history of hepatitis virus uptake interrupt/macrophage response to change in the face of *Mycobacterium tuberculosis*.

Thus, the above facts support the hypothesis that not only HIV but also other viruses are transmitted through body fluids and sex, play a role in the manifestation of tuberculosis. This is in line with the opinion of Nagelkerke (2012) who found similarities distribution age and sex of tuberculosis and sexually transmitted diseases. As an opportunistic disease, tuberculosis appears when there is interference with the balance between Th1 and Th2.

As the implications of the above, then when the results of liver function tests as monitoring treatment showed an increase, then the interpretation should be expanded not only as a result of drug toxicity, but also the possibility of an active viral hepatitis infection should also be explored. Mo et al study found that in the group of multi-infection with HIV/TB/HBV and HIV/TB/HCV incidence of abnormal liver function were significantly higher (OR 5.525). The study also showed that mortality was higher in multi infection (19.7%) (Mo et al 2014). Thus in the treatment of tuberculosis patients with abnormal liver function, requires a serious follow-up.

The limitation in this study is that the study is just a history of never infected with hepatitis B and hepatitis C, by measuring antibodies patient/control of hepatitis B (anti-HBV) and hepatitis C (anti-HCV). It is realized also the limitations of these two antibodies in detection history of hepatitis due to the emergence later than suppose is Ig M anti-HBc, but the period of orbit in the blood longer, although not all anti-HBc. thus, this study did not measure whether there is an antigen hepatitis to be active, was also negativity antibody result does not mean the absence of a history of infection.

CONCLUSION

Tuberculosis sufferers were more likely to have a history of hepatitis B virus 2.7 times than that of infected control but not sick. Coinfection of hepatitis C virus possibly does not play a role in the manifestation of tuberculosis, since Indonesia is an endemic area of hepatitis B, instead of hepatitis C, as individuals with hepatitis C infection history are minority. However, it remains necessary to have deep thought about the handling of tuberculosis. As long as the tuberculosis

bacteria is latent, body response is appropriate, not shifting towards tissue damage, the infection of the bacteria will not pose any problems. When this virus coinfection plays a role in the emergence of disease manifestations, it is necessary to think about this coinfection prevention which would be beneficial for the prevention of tuberculosis. In this case the possible prevention of hepatitis transmission is through blood products or sexual intercourse. In terms of management of the treatment of tuberculosis, screening liver function tests or hepatitis antibody-antigen test may need to be considered in preventing more severe hepatotoxicity and death.

REFERENCES

- Graubard SYH, Chen F, Barrett CJ, Qiao YL, Forman MR (2011). Tuberculosis and oncogenic HPV: potential co-infections in women at high-risk of cervical cancer in rural China. *Asian Pacific Journal of Cancer Prevention* 12, 1409-1415
- Lestari P, Sustini F, Endaryanto A, Asih R (2011). Home humidity increased risk of tuberculosis in children living with adult active tuberculosis cases. *Universa medicina* 30, 138-145
- Ministry of Health Republic of Indonesia (2010). Report on Result of National Basic Health Research (RISKESDAS 2010), Jakarta, Departemen Kesehatan Republik Indonesia
- Mo P, Zhu Q, Teter C, Yang R, Den L, Yan Y, Chen J, Zeng J, ui XE (2014). Prevalence, drug-induced hepatotoxicity, and mortality among patients multi-infected with HIV, tuberculosis, and hepatitis virus. *Int J Infect Dis* 28, 95-100
- Nagelkerke N (2012). Courtesans and consumption, how sexually transmitted infection drive tuberculosis epidemics. Eburon, Delft Netherlands.
- Pustynnikov S, Sagar D, Jain P, Khan Z (2014). Targeting the C-type lectins-mediated host-pathogen interaction with dextran. *J Pharm Pharm Sci* 17, 371-392
- Setiawati L, Endaryanto A, Kusumadewi A, Lestari P (2011). Effect of BCG vaccination and non-tuberculous mycobacterium infection on interferon gamma specific assay and a tuberculin skin test among children with a tuberculosis contact in Surabaya, Indonesia. *Southeast Asian J Trop Med Public Health* 42, 1460-1468
- Soemantri S, Senewe FP, Tjandrarini DH, Day R, Basri C, Manissero D, Mehta F, Dye C (2007). Three-fold reduction in the prevalence of tuberculosis over 25 years in Indonesia. *The International Journal of Tuberculosis and Lung Disease* 11, 398-404
- WHO (2013). TB country profile 2013. Available from <https://extranet.who.int/sree/Reports?op=Replet&nam>

e=%2FWHO_HQ_Reports%2FG2%2FPROD%2FEX
T%2FTBCountryProfile&ISO2=ID&LAN=EN&outty
pe=html. Accessed November 2, 2013
Zhang L, Zhang D, Chen W, Zou X, Ling L (2013).
High prevalence of HIV, HCV and tuberculosis and

associated risk behaviours among new entrants of
methadone maintenance treatment clinics in
Guangdong Province, China. PLoS One 8, e76931

11. ROLE OF HEPATITIS B AND HEPATITS C VIRUS COINFECTION IN TB DEVELOPMENT _ Lestari _ Folia Medica Indonesiana

ORIGINALITY REPORT

9%

SIMILARITY INDEX

7%

INTERNET SOURCES

3%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	repository.unair.ac.id Internet Source	3%
2	garuda.ristekdikti.go.id Internet Source	1%
3	isindexing.com Internet Source	1%
4	digitalcommons.law.ggu.edu Internet Source	<1%
5	bmcpublichealth.biomedcentral.com Internet Source	<1%
6	Rosnah Sutan, Satrinawati Berkat. "Does cultural practice affects neonatal survival- a case control study among low birth weight babies in Aceh Province, Indonesia", BMC Pregnancy and Childbirth, 2014 Publication	<1%
7	ar.scribd.com	

<1%

8

Pingzheng Mo, Qi Zhu, Caroline Teter, Rongrong Yang, Liping Deng, Yajun Yan, Jun Chen, Jie Zeng, Xi-en Gui. "Prevalence, drug-induced hepatotoxicity, and mortality among patients multi-infected with HIV, tuberculosis, and hepatitis virus", International Journal of Infectious Diseases, 2014

Publication

<1%

9

dosen.uta45jakarta.ac.id

Internet Source

<1%

10

repository.unhas.ac.id

Internet Source

<1%

11

es.scribd.com

Internet Source

<1%

12

www.va.gov

Internet Source

<1%

13

www.ncbi.nlm.nih.gov

Internet Source

<1%

14

www.nature.com

Internet Source

<1%

15

jurnal.ugm.ac.id

Internet Source

<1%

16

Zi-Yi Jin, Xing Liu, Ying-Ying Ding, Zuo-Feng Zhang, Na He. "Cancer risk factors among people living with HIV/AIDS in China: a systematic review and meta-analysis", Scientific Reports, 2017

Publication

<1%

17

Zhang, Lei, Di Zhang, Wen Chen, Xia Zou, and Li Ling. "High Prevalence of HIV, HCV and Tuberculosis and Associated Risk Behaviours among New Entrants of Methadone Maintenance Treatment Clinics in Guangdong Province, China", PLoS ONE, 2013.

Publication

<1%

18

Arthur, R.R.. "Hepatitis C antibody prevalence in blood donors in different governorates in Egypt", Transactions of the Royal Society of Tropical Medicine and Hygiene, 199705/06

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

11. ROLE OF HEPATITIS B AND HEPATITS C VIRUS COINFECTION IN TB DEVELOPMENT _ Lestari _ Folia Medica Indonesiana

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
