

Coumarins from Myanmar edible fruit tree (*Casimiroa edulis*)

by Nanik Siti Aminah

Submission date: 16-May-2020 12:50PM (UTC+0800)

Submission ID: 1325573912

File name: C06._Fulltext_-_COUMARIN_JICS_KHUNpdf.pdf (1.1M)

Word count: 2288

Character count: 10937

Coumarins from Myanmar edible fruit tree (*Casimiroa edulis*)

Khun Nay Win Tun^{a,b}, Nanik Siti Aminah^{*a}, Alfinda Novi Kristanti^a, Rico Ramadhan^a, Yoshiaki Takaya^c and HninThanda Aung^d

^aDepartment of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

E-mail: nanik-s-a@fst.unair.ac.id

^bDepartment of Chemistry, Taunggyi University, Shan State (South), Myanmar

^cFaculty of Pharmacy, Meijo University, Tempaku, Nagoya, Japan

^dDepartment of Chemistry, Mandalay University, Mandalay, Myanmar

Manuscript received online 03 April 2019, revised 22 May 2019, accepted 31 May 2019

This research expresses the phytochemical study from the Myanmar edible fruit tree, *Casimiroa edulis* (Rutaceae). The result revealed that the isolation and identification of two furanocoumarins (bergapten **1** and isopimpinellin **2**) from the stem bark of this plant. Their molecular structures were elucidated and identified by using NMR spectroscopy in combination with IR, UV and HRMS spectra data, respectively. Furthermore, these two compounds were investigated for their anti-diabetic activity. According to the result, bergapten **1** and isopimpinellin **2** are not essentially good for anti-diabetic activity. This is the first report of two furanocoumarins from the Myanmar edible fruit tree.

Keywords: *Casimiroa edulis*, furanocoumarins, NMR spectroscopy, anti-diabetic activity.

Introduction

Casimiroa edulis La Llave (Rutaceae) is a genus of comprising only 10 species which is distributed in tropical and subtropical regions, including Myanmar. Among them, *Casimiroa edulis* is the well-known species^{1,2}. The traditional used and the pharmacological studies of the leaves and seeds of *Casimiroa edulis* displayed various biological activities. In Myanmar, this plant is named Thar-kyar-thee.

Phytochemical studied from genus *Casimiroa* assist to the separation of various chemical constituents such as 18 flavonoids (isolated from leaves, fruits, and seeds), 24 alkaloids (isolated from leaves, fruits, bark, and seeds), and 16 coumarins (isolated from leaves, fruits, and seeds). Pharmacological investigations of isolated flavonoids are antioxidant, anti-mutagenic, solid tumor selective cytotoxicity, vasodilation and radical-scavenging activities, the investigations of isolated alkaloids are anti-mutagenic, solid tumor selective cytotoxicity, anti-hypertensive, cardiovascular activities, and the investigations of isolated coumarins are anti-coagulant, vasodilation and radical-scavenging, adipogenesis, solid tumor selective cytotoxicity activities, respectively³⁻¹⁴. This

paper deals with the structure elucidation of two furanocoumarins from the stem bark of *Casimiroa edulis*.

Experimental

General:

The infrared spectra (IR) were obtained on FTIR-8400S (Shimadzu) using KBr. UV spectra were examined in MeOH by using UV-Vis Shimadzu spectrometer. Column chromatography (CC) was achieved on silica gel (BW-820H). Analytical thin-layer-chromatography (TLC) was completed at room temperature on pre-coated Kieselgel silica gel 60 F₂₅₄ (Merck) aluminium plates. Melting points were determined by fisher john melting point apparatus. All NMR spectra were checked in CDCl₃ by using a Bruker Avance 3 (¹H: 600 MHz and ¹³C: 151 MHz) with TMS as an internal standard. Chemical shifts are described in part per million (δ , ppm) based on their solvent signal (chloroform-*d*: δ_H 7.26, δ_C 77.16) coupling constants *J* value in Hertz (Hz). HR-FAB-MS was attained with a JEOL JMS HX-110 mass spectrometer.

Plant material:

Casimiroa edulis La Llave was collected from Namp-see


Village, Taunggyi (Shan State), Myanmar, in August 2016, and was identified by the Professor of Botany Department, University of Taunggyi.

Extraction and isolation:

The dry sample of *Casimiroa edulis* (1 kg) was exhaustively extracted with MeOH (3 L) at room temperature for four weeks. The methanol extract was concentrated under pressure using rotator evaporator for dryness. Then, the methanolic extract was partition between *n*-hexane and MeOH (300 ml×3, v/v). The concentrated MeOH extract (50 g) was subjected to vacuum liquid chromatography (VLC) and eluted with *n*-hexane-EtOAc in a stepwise gradient solvent system (10:0, 6:4, 0:10, v/v) to afford several fractions. Column eluate was collected in 150 ml glass bottles to yield 28 fractions and examined under UV. Based on analytical TLC, the fractions were grouped and evaporated on a rotary evaporator to afford six fractions (J-1 to J-6). Fractions J-5 (2 g) were additionally purified by silica gel column chromatography with a gradient solvent system of *n*-hexane in EtOAc (10:0, 0:10) to give colourless needles of compound **1**, and pale-yellow crystalline solids of compound **2**.

Results and discussion

Compound **1** was obtained as colorless needle crystals. Its melting point was 186–206°C. Based on the molecular ion peak at m/z 217.0501 $[M + H]^+$ (Calcd. 216.0423) in the HR-FAB-MS, the molecular formula was determined as $C_{12}H_8O_4$. IR spectrum informed the characteristic of sharp bands at 3143, 3115, 3088, 3010, 2924, 2852, 1737, 1720, 1625 and 1581 cm^{-1} which could be assigned to $-OCH_3$, sp^2-CH , sp^3-CH , lactone $C=O$, $C=C$ ring skeletal stretching vibration, respectively. The UV spectrum showed the bands



Comps.	R ₁	R ₂
1	OCH ₃	H
2	OCH ₃	OCH ₃

Fig. 1. Chemical structure of compound **1** and **2**.

at λ_{max} 221, 249, 259, 268, and 311 nm, which were typical of a furanocoumarin skeleton. The 1H NMR spectrum was exhibited as an easily distinguishable pair of doublets at δ_H 6.28 (d, 1H, 9.8 Hz, H-3) and δ_H 8.16 (d, 1H, 9.8 Hz, H-4). It also displayed two furan doublets at δ_H 7.60 and 7.02 (each, d, 1H, 2.4 Hz, H-2' and H-3'), and an aromatic singlet at δ_H 7.14 (s, 1H, H-8). Combining analysis of ^{13}C NMR and DEPT spectra data, compound **1** displayed 12 carbons resonance, including five methine, one oxygenated and six quaternary carbons, respectively. In the 1H - 1H COSY experiment, the cross-peak correlations of H-3 with H-4, and the cross-peak correlations of H-2' through H-3' were observed. The long-range HMBC attachment from H-3 to C-2 and C-10, H-4 to C-2 and C-9, H-8 to C-6, C-7, C-9 and C-10, H-2'' to C-3'', C-6 and C-7, H-3'' to C-2', C-6 and C-7 were also observed. The remaining one singlet proton at δ 4.27 (integrating for three protons) showed the presence of methoxy group on the aromatic ring. This confirmation was deduced by HMBC spectrum, which confirms that methoxy proton showed the long-range correlation with the C-5 carbon at δ_C 149.6. Analysis of UV, IR, 1D, 2D NMR and HRMS spectra as well as on comparison with Literature, were allowed to determine the structure of compound **1** as bergapten^{15–17}.

Compound **2** was obtained as pale-yellow crystalline solids with a melting point at 178–180°C. Its molecular formula was assigned as $C_{13}H_{10}O_5$ based on HR-FAB-MS (m/z 247.0606 $[M + H]^+$, Calcd. 246.0528). The UV spectrum ex-




Fig. 2. 1H - 1H COSY correlations (a) and major 1H - ^{13}C long-range HMBC (b) correlation in compound **1**.




Fig. 3. 1H - 1H COSY correlations (a) and major 1H - ^{13}C long-range HMBC (b) correlation in compound **2**.

hibited the absorption maxima at 241, 248, 268 and 311 nm, which is consistent with the presence of a furocoumarin. The IR spectrum revealed the presence of an α,β -unsaturated lactone (ν_{\max} 1751 and 1718 cm^{-1}), methoxy group (3132 and 3159 cm^{-1}), C-H stretching vibration of sp^2 hydrocarbon (3084 and 3005 cm^{-1}), asymmetrical and symmetrical C-H stretching vibration of sp^3 hydrocarbon (2843 and 2953 cm^{-1}), and C=C ring skeleton stretching vibration (ν_{\max} 1604 and 1593 cm^{-1}) groups. ^{13}C NMR of compound **2** included 13 carbon signals, comprising 2 methoxy carbon signals, 4 methine carbon signals, and 7 quaternary carbon signals, which were classified by DEPT and HSQC experiment. Its ^1H NMR spectrum was exhibited a distinct pair of doublets at δ_{H} 6.27 and 8.11 (each, 1H, 9.8 Hz) assignable to C-3, C-4 protons, a furan doublet at δ_{H} 7.62 and 6.99 (each, 1H, 2.2 Hz) assignable to C-2', C-3' protons. The ^1H - ^1H COSY spectrum showed a correlation between H-3 and H-4 protons while H-2' proton had a correlation to the H-4' proton. In HMBC spectrum, the following correlations were observed: the proton signal at δ_{H} 6.27 was connected with C-2 and C-10, the proton signal at δ_{H} 8.11 was connected with C-2 and C-9, the proton signal at δ_{H} 7.62 was correlated with C-2', C-6 and C-7, the proton signal at δ_{H} 6.99 was correlated with C-3', C-6 and C-7. The remaining methoxy protons at δ 4.16 (integrating for 6 protons) were placed at C-5 and C-8 on the

Table 1. ^1H NMR and ^{13}C NMR data of compound **1** in CDCl_3 and bergapten ^1H in CDCl_3

Position	Compd. 1		Lit. [Ref. 16]	
	^1H NMR ^a	^{13}C NMR ^b	^1H NMR ^c	^{13}C NMR ^d
2	—	161.2	—	161.4
3	6.28 (d, 9.8 Hz)	112.6	6.27 (d, 9.8 Hz)	112.6
4	8.16 (d, 9.8 Hz)	139.2	8.15 (d, 9.8 Hz)	139.4
5	—	149.6	—	149.7
6	—	112.7	—	112.9
7	—	158.4	—	158.5
8	7.14 (s)	93.9	7.13 (s)	94.0
9	—	152.7	—	151.9
10	—	106.5	—	106.6
2'	7.60 (d, 2.4 Hz)	144.8	7.59 (d, 2.4 Hz)	144.9
3'	7.02 (d, 2.4 Hz)	105.0	7.02 (d, 2.4 Hz)	105.2
5-OMe	4.27 (s)	60.1	4.27 (s)	60.3
8-OMe	—	—	—	—

^aSpectra recorded in 600 MHz, ^bspectra recorded in 151 MHz, ^cspectra recorded in 400 MHz and ^dspectra recorded in 100 MHz.

Table 2. ^1H NMR and ^{13}C NMR data of compound **2** in CDCl_3 and isopimpinellin in CDCl_3

Position	Compd. 2		Lit. [Ref. 16]	
	^1H NMR ^a	^{13}C NMR ^b	^1H NMR ^c	^{13}C NMR ^d
2	—	160.4	—	160.4
3	6.27 (d, 9.8 Hz)	112.8	6.26 (d, 9.8 Hz)	112.8
4	8.11 (d, 9.8 Hz)	139.4	8.10 (d, 9.8 Hz)	139.3
5	—	144.3	—	144.3
6	—	114.8	—	114.8
7	—	150.0	—	150.0
8	—	128.2	—	128.2
9	—	143.7	—	143.7
10	—	107.6	—	107.6
2'	7.62 (d, 2.2 Hz)	145.1	7.61 (d, 2.3 Hz)	145.1
3'	6.99 (d, 2.2 Hz)	105.1	6.99 (d, 2.3 Hz)	105.1
5-OMe	4.16 (s)	61.7	4.16 (s)	61.7
8-OMe	4.16 (s)	60.8	4.14 (s)	60.8

^aSpectra recorded in 600 MHz, ^bspectra recorded in 151 MHz, ^cspectra recorded in 400 MHz and ^dspectra recorded in 100 MHz.

basic of long range HMBC correlations. With the aid of above data and previous report, compound **2** was identified as isopimpinellin^{16–18}.

Conclusions

In this study, two known compounds were obtained from the stem bark of *Casimiroa edulis*. The structure of isolated compounds identified as bergapten **1** and isopimpinellin **2** based on their spectroscopic data, such as UV-Vis, FTIR, 1D NMR, 2D NMR, and HRMS. The isolated compounds are not important for their anti-diabetic activity. A lot of researchers were reported these two compounds from many sources of natural product. This is the first report on the occurrence of bergapten and isopimpinellin from Myanmar edible fruit tree *Casimiroa edulis*.

Acknowledgements

This work was supported in part by Ailangga Development Scholarship (ADS) and "RISET MANDAT GRANT" of Universitas Airlangga, Surabaya, Indonesia. KNWT gracefully acknowledges Reactors, Vice-Rectors, and Professors from Universitas Airlangga.

Supporting Information

1D and 2D NMR spectrum are described.

References

1. M. Yamamoto, T. Tomita, M. Onjo, K. Ishihata, T. Kubo and S. Tominaga, *HortScience*, 2007, **42(6)**, 1329.
2. N. Satheesh, *American-Eurasian J. Agric. & Environ. Sci.*, 2015, **15(8)**, 1574.
3. A. Ito, L. A. Shamon, B. Yu, E. Mata-Greenwood, S. K. Lee, R. B. Breemen, R. G. Mehta, N. R. Farnsworth, H. H.S. Fong, J. M. Pezzuto and A. Douglas, *J. Agric. Food Chem.*, 1998, **46**, 3509.
4. A. S. Awaad, N. H. El-Sayed, D. J. Maitland and T. J. Mabry, *Pharm. Biol.*, 2006, **44(4)**, 258.
5. A. S. Awaad, N. A. Al-Jaber, G. A. Soliman, M. R. Al-Outhman, M. E. Zain, J. E. Moses and R. M. El-Meligy, *Phytother. Res.*, 2012, **26**, 452.
6. A. N. García-Argáeza, N. M. González-Lugo, H. Parra-Delgado, and M. Martínez-Vázquez, *Biochem. Syst. Ecol.*, 2005, **33**, 441.
7. A. S. Awaad, D. J. Maitland and S. M. Moneir, *Chem. Nat. Compd.*, 2007, **43(5)**, 576.
8. A. E. M. Khaleel, *Monatshefte Fur Chemie*, 2002, **133**, 183.
9. B. Heneka, H. Rimpler, A. Ankli, O. Sticher, S. Gibbons and M. Heinrich, *Phytochemistry*, 2005, **66**, 649.
10. D. L. Dreyer, *J. Org. Chem.*, 1968, **33(9)**, 3577.
11. H. Nagai, T. Tanaka, T. Goto, T. Kusudo, N. Takahashi, and T. Kawada, *Biosci. Biotechnol. Biochem.*, 2014, **78(2)**, 296.
12. A. N. García-Argáeza, N. M. González-Lugob, H. Parra-Delgadob and M. Marínez-Vázquezb, *Z. Naturforsch.*, 2004, **59(b)**, 245.
13. R. Bertin, Z. Chen, M. Martínez-vázquez, A. García-argaéz and G. Froidi, *Phytomedicine*, 2014, **21**, 586.
14. X. U. Ya-ming, M. C. Ramirez-ahumada, F. A. Valeriete and A. A. L. Gunatilaka, *Chin. J. Nat. Med.*, 2011, **9(5)**, 334.
15. C. Chunyan, S. Bo, L. Ping and L. Jingmei, *J. Liq. Chrom. Relat. Tech.*, 2009, **32**, 136.
16. T. O'Neill, J. A. Johnson, D. Webster and C. A. Gray, *J. Ethnopharmacol.*, 2013, **147**, 232.
17. J. N. Ngwendson, E. Bedir, S. M. N. Efange, C. O. Okunji, M. M. Iwu, B. G. Schuster and I. A. Khan, *Pharmazie*, 2003, **58**, 587.
18. G. Wang, Z. Zhou, C. Cheng, J. Yao and Z. Yang, *Vet. Parasitol.*, 2008, **158**, 144.

Coumarins from Myanmar edible fruit tree (Casimiroa edulis)

ORIGINALITY REPORT

22%

SIMILARITY INDEX

9%

INTERNET SOURCES

19%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Chakapong Intaraudom, Nantiya Bunbamrung, Aibrohim Dramaee, Nattawut Boonyuen et al. "Acremonidins F–H and acremoxanthones F–G, antimicrobial substances from the insect fungus *Verticillium* sp. BCC33181", *Tetrahedron*, 2016
Publication 2%
- 2 repository.unair.ac.id
Internet Source 2%
- 3 Girennavar, B.. "Furocoumarins from grapefruit juice and their effect on human CYP 3A4 and CYP 1B1 isoenzymes", *Bioorganic & Medicinal Chemistry*, 20060415
Publication 1%
- 4 Yu Zhao, Yuan-Ling Ku, Xiao Jiang Hao, Shoei-Sheng Lee. "Preparation of Analogues of Territrem B, a Potent AChE Inhibitor", *Tetrahedron*, 2000
Publication 1%
- 5 iopscience.iop.org
Internet Source 1%

6

prp.hec.gov.pk

Internet Source

1%

7

Zheng, Zong-Ping, Yan Yan, Ji Xia, Shuang Zhang, Mingfu Wang, Jie Chen, and Yang Xu. "A phenylacetaldehyde–flavonoid adduct, 8-C-(E-phenylethenyl)-norartocarpetin, exhibits intrinsic apoptosis and MAPK pathways-related anticancer potential on HepG2, SMMC-7721 and QGY-7703", *Food Chemistry*, 2016.

Publication

1%

8

Wei Wang. "Two new pterocarpenes from *Hedysarum multijugum*", *Journal of Asian Natural Products Research*, 1/1/2003

Publication

1%

9

Alessandra Braca, Annahil Armenise, Ivano Morelli, Jeannette Mendez et al. " Structure of Kaurane-Type Diterpenes from and their Potential Anticancer Activity ", *Planta Medica*, 2004

Publication

1%

10

unair.ac.id

Internet Source

1%

11

Mino, T.. "Palladium-catalyzed asymmetric allylic alkylation using chiral hydrazone ligands with ferrocene skeleton", *Journal of Organometallic Chemistry*, 20040901

Publication

1%

12

Wang, L.Q.. "Three new C²1 steroidal glycosides from the roots of *Cynanchum inamoenum*", *Chinese Chemical Letters*, 200710

Publication

1%

13

www.j3.jstage.jst.go.jp

Internet Source

1%

14

Dudutienė, Virginija, Asta Zubrienė, Alexey Smirnov, David D. Timm, Joana Smirnovienė, Justina Kazokaitė, Vilma Michailovienė, Audrius Zakšauskas, Elena Manakova, Saulius Gražulis, and Daumantas Matulis. "Functionalization of Fluorinated Benzenesulfonamides and Their Inhibitory Properties toward Carbonic Anhydrases", *ChemMedChem*, 2015.

Publication

1%

15

P.E.S. de Oliveira, L.M. Conserva, A.C. Brito, R.P.L. Lemos. " Coumarin Derivatives from . and its Larvicidal Activity Against . ", *Pharmaceutical Biology*, 2008

Publication

1%

16

Ligang Zhou, Duan Li, Weibo Jiang, Zhizhong Qin, Shuang Zhao, Minghua Qiu, Jianyong Wu. " Two ellagic acid glycosides from *Lam.* with antifungal activity on ", *Natural Product Research*, 2007

Publication

1%

17

Weihong Lin, Weilin Sun, Jun Yang, Qihang Sun, Zhiquan Shen. "Fabrication of Magnetic Thin Films by a Layer-by-Layer Self-Assembly Approach", The Journal of Physical Chemistry C, 2009

Publication

<1%

18

www.degruyter.com

Internet Source

<1%

19

aip.scitation.org

Internet Source

<1%

20

Hiroyuki Nakagawa, Yoshihisa Takaishi, Yoshinori Fujimoto, Carmenza Duque et al. "Chemical Constituents from the Colombian Medicinal Plant ", Journal of Natural Products, 2004

Publication

<1%

21

Jaime Raul O. Janairo, Gerardo C. Janairo, Consolacion Y. Ragasa, Bruce F. Bowden. "New Xenicanes from ", Natural Product Research, 2007

Publication

<1%

22

Consolacion Y. Ragasa, Angel Lyn Kristin C. Co, John A. Rideout. " Antifungal metabolites from ", Natural Product Research, 2005

Publication

<1%

23

Sang Zin Choi. "Pytochemical constituents of

the aerial parts from *Solidago virga-aurea* var. *gigantea*", *Archives of Pharmacal Research*, 02/2004

Publication

<1%

24

kb.psu.ac.th

Internet Source

<1%

25

Hong-Xiang Lou. "A cytotoxic diterpenoid and antifungal phenolic compounds from *Frullania muscicola* steph", *Journal of Asian Natural Products Research*, 1/1/2002

Publication

<1%

26

Chen, B.. "Sesquiterpenoids from *Ferula kuhistanica*", *Phytochemistry*, 200008

Publication

<1%

27

Aranya Jutiviboonsuk, Hong-Jie Zhang, Tamara P. Kondratyuk, Angkana Herunsalee et al. "Isolation and Characterization of Cancer Chemopreventive Compounds from *Barringtonia maunwongyathiae*", *Pharmaceutical Biology*, 2008

Publication

<1%

28

Junko Ito, Yoshiaki Takaya, Yoshiteru Oshima, Masatake Niwa. "New oligostilbenes having a benzofuran from *Vitis vinifera* 'Kyohou'", *Tetrahedron*, 1999

Publication

<1%

Nakagawa, Hiroyuki, Yoshihisa Takaishi,

29 Yoshinori Fujimoto, Carmenza Duque, Cristina Garzon, Mitsunobu Sato, Masato Okamoto, Tetsuya Oshikawa, and Sharif Uddin Ahmed. "Chemical Constituents from the Colombian Medicinal Plant *Maytenus laevis*", *Journal of Natural Products*, 2004.

Publication

30 Henrieta Stankovičová, Margita Lácová, Anton Gáplovský, Jarmila Chovancová, Nad'a Prónayová. "Reaction of 3-formylchromones with aromatic amino carboxylic acids", *Tetrahedron*, 2001

Publication

31 Takahiro Senda, Yasuhiro Yamada, Masakazu Morimoto, Noriko Nono, Toshiaki Sogabe, Shingo Kubo, Satoshi Sato. "Analyses of oxidation process for isotropic pitch-based carbon fiber using model compounds", *Carbon*, 2018

Publication

32 Moriyama, M.. "Structure and neurotrophic activity of novel sesqui-neolignans from the pericarps of *Illicium fargesii*", *Tetrahedron*, 20070514

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Coumarins from Myanmar edible fruit tree (*Casimiroa edulis*)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
