

# Phytoconstituents of Genus Micromelum and Their Bioactivity —a Review

*by* Nanik Siti Aminah

---

**Submission date:** 02-Jun-2020 04:54PM (UTC+0800)

**Submission ID:** 1336417441

**File name:** 2.\_Phytoconstituents\_of\_Genus\_Micromelum\_2020.pdf (2.72M)


**Word count:** 5888

**Character count:** 34593

# Phytoconstituents of Genus *Micromelum* and Their Bioactivity—a Review

Natural Product Communications  
 Volume 15(5): 1–15  
 © The Author(s) 2020  
 Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
 DOI: 10.1177/1934578X20927124  
[journals.sagepub.com/home/npx](https://journals.sagepub.com/home/npx)



Tin Myo Thant<sup>1,2</sup>, Nanik Siti Aminah<sup>1</sup> , Alfinda Novi Kristanti<sup>1</sup>, Rico Ramadhan<sup>1</sup>, Hnin Thanda Aung<sup>3</sup>, and Yoshiaki Takaya<sup>4</sup>

## Abstract

The genus *Micromelum* belongs to the Rutaceae family. As its rich bioactive constituents its stems, flowers, leaves, and roots have been used in traditional medicine, for the treatment of various diseases from ancient time. Phytochemically, many bioactive compounds, including coumarins, polyoxygenated flavonoids, phenylpropanoic acid derivatives, quinolone alkaloids, and also carbazole alkaloids, have been reported as secondary metabolites of the *Micromelum* spp. including many new compounds. Therefore, *Micromelum* spp. are considered potential for drug leads. In this article, we present an overview of secondary metabolites isolated from genus *Micromelum* and their bioactivities that have been reported between 1982 and 2019.

## Keywords

*Micromelum*, Rutaceae, coumarins, flavonoids, quinolone and carbazole alkaloids

Received: November 27th, 2019; Accepted: April 22nd, 2020.

## Introduction to Chemical Constituents of Genus *Micromelum* and Their Bioactivity

*Micromelum* spp. (Rutaceae) consist of 9 species found mostly in Samoa, Fiji, and Tonga islands, Australia, Southeast Asia, southern China, Ceylon, northeastern India, and West Pakistan.<sup>1,2</sup> *Micromelum* spp. are known for being rich in coumarins such as the 6- and 8-prenylated coumarins. In addition, the carbazole alkaloids, dihydrocinnamic acid derivatives, and flavonoids have been isolated from it.<sup>3,4</sup> The leaves and stems of *Micromelum* species have been found to contain coumarins, phenylpropanoic acid derivatives, polyoxygenated flavonoids, and also alkaloids.<sup>5,6</sup> The following *Micromelum* species have been well investigated due to their pharmaceutical properties.

*M. integerrimum* is a tree that grows up to 8 m high. Its parts have rust-colored pubescent. It usually grows in moist mountain forests, maritime thickets in sandy soil; near sea level to 2000 m and is widely distributed in China, Bhutan, Cambodia, India, Laos, Myanmar, Nepal, Thailand, and Vietnam.<sup>7</sup> *M. integerrimum* is highly diversified for its secondary metabolites, and apart from acridone and carbazole alkaloids, a variety of coumarin derivatives, especially 6,7-di and 7,8-disubstituted coumarin core structures, having at least a prenyl unit have been isolated. Naturally occurring coumarin derivatives from *Micromelum* spp. exhibit a variety of biological activities including anti-corpulence, cytotoxicity, anti-platelet, and antimutagenicity.<sup>3</sup>

*M. minutum* is a small spineless tree that can reach up to 3 m in height. It is widely distributed in Southeast Asia and in Pacific islands. The stems, flowers, leaves, and roots are used pharmaceutically for a variety of indications.<sup>8,9</sup> The plant is known to contain coumarins, some of which are active compounds showing strong cytotoxic activities. The leaves are traditionally used to treat fever and giddiness. The poultice of the boiled roots is used for ague.<sup>10</sup> In Indochina, the roasted and crushed leaves are rubbed to skin irritated by scabies and are considered to be emmenagogues.<sup>11</sup> Previous phytochemical investigations on the different parts of *M. minutum* have showed the presence of coumarins, triterpenes, alkaloids, and phenylpropanoids.<sup>12</sup>

*M. falcatum* is a tree of 1-3 m high and its leaflet blades are alternate, ovate to lanceolate, in equilateral. Flowers are ellipsoid or broad in bud. Petals are white, oblong, outside glabrous, or

<sup>1</sup>Dept of Chemistry, Fac of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

<sup>2</sup>Department of Chemistry, Mandalay Degree College, Mandalay, Myanmar

<sup>3</sup>Department of Chemistry, Mandalay University, Myanmar

<sup>4</sup>Faculty of Pharmacy, Meijo University, Yagotoyama, Tempaku, Nagoya, Japan

## Corresponding Author:

Nanik Siti Aminah, Dept of Chemistry, Fac of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia.

Email: [nanik-s-a@fst.unair.ac.id](mailto:nanik-s-a@fst.unair.ac.id)







Figure 1. Coumarins from *Micromelum minutum*, *M. integerrimum*, *M. falcatum*, and *M. zeylanicum*.

pubescent. Fruits are ellipsoid to ovoid, 1- or 2-seeded. Its flowering period is from January to April. It is usually found in the mountain area at heights below 1200 m in China, Cambodia, Laos, Myanmar, Thailand, and Vietnam.<sup>13</sup> Its leaves and roots are used for treating infected wounds, oedynolysis, rheumatism, muscular atrophy, and insect bites.<sup>4,14</sup> *M. falcatum* has been found to contain several coumarins, such as 7-oxygenated coumarins, dihydrocinnamic acid derivatives, the dimeric indole alkaloid (yuehchukene), the carbazole alkaloid, 5,6-pyranoglycozoline, the

6-prenylated coumarin, micromelin, and the 8-prenylated coumarins, phebalosin and murpanidin.<sup>6,15,16</sup> The dimeric indole alkaloid yuehchukene was found to have potent anti-implantation activity.<sup>17</sup>

*M. compressum* is a tree described as unarmed and glabrous. Its leaves are odd-pinnate, the leaflets are alternate, and the fruits are ovoid. Phytochemical efforts on the  $\text{CH}_2\text{Cl}_2$ -methanol extract of *M. compressum* have resulted in the isolation and identification of 3 polymethoxylated flavones.<sup>1</sup>



**Figure 2.** Coumarins isolated from *Micromelum falcatum*.

*M. hirsutum* is a tree of up to 10 m height with greenish-yellow flowers and aromatic leaves. It is widely distributed in peninsular Malaysia (Melaka and Pahang northward), Andaman Islands, Myanmar, Laos, Thailand, and Vietnam.<sup>18</sup> In 2010, Rodphukdeekul research group investigated antitumor activity of the dichloromethane extract from branches of *M. hirsutum* on human B lymphoma cells, Ramos. The result demonstrated that the extract has antitumor potential against B lymphoma.<sup>19</sup> In Myanmar its leaves are widely used as spices in curries. Phytochemical analysis of this species has reported the presence of many carbazole alkaloids, as well as the  $\gamma$ -lactone compounds.<sup>18</sup>

*M. glanduliferum* is a tropical or subtropical small tree found in Thailand. Coumarins and some carbazole alkaloids compounds, such as osthol, phebalosin, minumicrolin, murrangatin, murralongin, micormarin A, microminutinin, micromelin, and 2,7-dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole and 7-methoxy heptaphylline, were reported from the roots of *M. glanduliferum*.<sup>20</sup>


The chemical analysis of leaves and stems of *M. zeylanicum* has reported an oxazole alkaloid, O-methylhalfordinol, a coumarin, micromelin, and a flavone, 5-hydroxy-3,3',4',7-pentamethoxyflavone. In addition, the stems contained 6-formyl-7-methoxycoumarin, and the leaves contained the carbazole alkaloid, koenigine, and *p*-sitosterol.<sup>21</sup>

In this review the articles were collected from 1982 to 2019. The main purpose of this article was to evaluate the diversity of secondary metabolites isolated from *Micromelum* spp. in order to provide a complete reference of the diverse secondary metabolites from *Micromelum* spp. To the best of our knowledge this is the first review about secondary metabolites isolated from genus *Micromelum*.

## Phytochemical Aspects of *Micromelum* Spp. and Their Tested Bioactivities


### Coumarins

Coumarins are highly active biological substances and major secondary metabolites isolated from *Micromelum* spp. They are widely used in medicine, perfumes, cosmetics, and laser dyes.<sup>6</sup> Coumarins have antioxidant activity and they preserve the



**Figure 3.** Some coumarins isolated from *Micromelum integerrimum* and *M. minutum*.

levels of other antioxidants in human plasma. Coumarins also inhibited lipid peroxidation and increased the activity of antioxidants, superoxide dismutase, and catalase.<sup>22</sup> Many bioactive coumarins have been reported from genus *Micromelum*, and the basic core coumarin compound, umbelliferone (1), was isolated from the roots of *M. minutum*.<sup>8</sup> Scopoletin compound 7-hydroxy-6-methoxy-2H-chromen-2-one (2) was isolated from the fruits of *M. minutum*<sup>23</sup> and 6-hydroxy-7-methoxy-2H-chromen-2-one (3) was isolated from the stems and leaves of *M. integerrimum*.<sup>24</sup> Another coumarin, 6-formyl-7-methoxycoumarin (4), was isolated from the leaves and stems of *M. zeylanicum*.<sup>21</sup>



**Figure 4.** Coumarins from *Micromelum minutum* and *M. falcatum*.





Figure 5. Coumarins from the *Micromelum minutum*.

7-Methoxyprenylated coumarin, osthonon (5), and osthol (6) were isolated from the roots of *M. minutum*.<sup>8</sup> Microfalcrin (7) was isolated from the leaves of *M. falcatum*.<sup>16</sup> This compound (7) was also found in *M. minutum*<sup>8</sup> (Figure 1).

One-prenylated coumarins named micromarin-F (8), -G (9), and -H (10) were isolated from the stems of *M. minutum*.<sup>25</sup> The compound murralongin-1 (11) was isolated from the roots of *M. minutum*.<sup>8</sup> Murralongin-2 (12) and murralonginol (13) were isolated from the fruits of *M. minutum*.<sup>23,26</sup> 7-Oxygenated coumarins, phebalosin (14), murrangatin (15), murrangatin acetate (16), murracarpin (17), and minumicrolin (18) were isolated from the roots of *M. minutum*<sup>8</sup> (Figure 1). Microminutin (19)

was isolated from the acetone extract of the stems of *M. minutum*.<sup>25</sup> Some coumarins, such as 7-methoxy-8-(4'-methyl-3'-furyl)coumarin (20), microminutin B (21), microminutin C (22), and micromeloside A (23), were isolated from the finely powdered fruits of *M. falcatum*.<sup>4</sup> In 2014, Kouloura et al reported microcoumaririn (24) from the leaves of *M. falcatum*. The anti-inflammatory properties of compounds (24, 32) were evaluated by measuring the inhibition of the proinflammatory mediator nuclear factor kappa B induction and nitric oxide (NO) production but these 2 compounds did not show any inhibition.<sup>16</sup>





**Figure 6.** Coumarins from the *Micromelum minutum*, *M. integerrimum*, and *M. falcatum*.

Micromelin or 7-methoxy-6-((1R,2R,5R)-5-methyl-4-oxo-3,6-dioxabicyclo[3.1.0]hexan-2-yl)-2H-chromen-2-one (25) was isolated from the fruits of *M. minutum*.<sup>25</sup> This compound is common to all species of *Micromelum* so far examined and can perhaps be considered as a chemotaxonomic marker for the genus.<sup>27</sup> The compound acetyldihydromicromelin A (26) was isolated from the aerial parts of *M. minutum*.<sup>26</sup> One-prenylated coumarin, namely hydramicromelin D (27), was isolated from the twigs of *M. integerrimum*.<sup>3</sup> Hopeyhopol (28) and marmesin (29) were isolated from the whole plants of *M. minutum*.<sup>28</sup> (-) Decursinol (30) was isolated from the finely powdered fruits of *M. falcatum*.<sup>4</sup> One-prenylated coumarins named micromarin-A (31), -B (32), and -C (33) were isolated from the seeds of *M. minutum*<sup>22</sup> and fruits of *M. falcatum*<sup>16</sup> (Figure 3).

Ester coumarinoids, (3*S*,4*S*)-3,4-dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-ethylenebutyl 3-methylbutanoate (34), (3*R*,4*S*)-3,4-dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl-3-methylbutanoate (35), and 3-hydroxy-4-methoxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl-3-ethylbutanoate (36), were isolated from the finely powdered fruits of *M. falcatum*<sup>4,16</sup> (Figure 4). Another ester coumarin, 7-methoxy-8-(2-hydroxymethyl-1-*O*-isovaleryl-4-butenyl) coumarin (37), was isolated from the stem bark of *M. falcatum*.<sup>29</sup>

Murralonginol isovalerate-1 (38) was isolated from the stems of *M. minutum*.<sup>23</sup> Toxicity of compound (37) isolated

from *M. falcatum* was tested against brine shrimp larvae using a 96-well plate assay and the compound (37) exhibited an LC<sub>50</sub> value of 6.8 μM, indicating that it was a potent toxic natural product and its in vitro antiproliferative activities against lung cancer (HvEvc) cell lines by the MTT method, and (37) displayed moderate activity against the HvEvc cell line with IC<sub>50</sub> value of 35.7.<sup>29</sup> Two-prenylated coumarins, murralonginol isovalerate-2 (39), minutuminolate (40), and 7-demethylmurralonginolisovalerate (41), were isolated from the fruits and roots of *M. minutum*<sup>8,23</sup> (Figure 4). Micromarininate (42) was isolated from the EtOAc extract fruits of *M. falcatum*<sup>4</sup> (Figure 4).

Five other geranylcoumarins, such as 8-hydroxyisocapnolactone-2',3'-diol (43), isolated from *M. minutum* have in vitro antiparasitoid activity against *Plasmodium falciparum* FCR-3 and D-10 strains with IC<sub>50</sub> values of 6.39 μg/mL (16.99 μM) and 24.23 mg/mL (64.45 μM), respectively.<sup>10</sup> 8-methoxycapnolactone (44), 3",4"-dihydrocapnolactone (45), 2',3' epoxyisocapnolactone (46), 8-hydroxy-3",4"-dihydrocapnolactone-2',3'-diol (47), and 8,4-dihydroxy-3,4-dihydrocapnolactone-2,3-diol (48) were isolated from the leaves of *M. minutum*<sup>10,11,30</sup> (Figure 5). These coumarins contained a lactone ring in their structure. In 2009, Ratna et al reported the cytotoxicity of these lactone coumarins on T-lymphoblastic leukemia (CEM-SS), promyelocytic leukemia (HL60), cervical cancer (HeLa), and liver cancer (HepG2) cell lines. Among the tested compounds, (43) showed the most active with IC<sub>50</sub> values of 2.9, 2.5, 6.9, and 5.9 μg/mL, respectively.<sup>11</sup> In 2014, the same group investigated the cytotoxic activity of compound (43) on MCF-7 and T47D cells. The results showed IC<sub>50</sub> values of 8 μg/mL (21.2 μM) and 4 μg/mL (10.6 μM) respectively.<sup>31</sup>

Two monoterpene coumarins, minutin A (49) and minutin B (50), were isolated from the leaves of *M. minutum* (Figure 5) and their cytotoxic activity against *Leishmania major* and cancer cells were already studied. It has been reported that minutin B (50) has strong cytotoxic activity against the cancer cell lines such as SBC3, A549, K562, and K562/ADM with IC<sub>50</sub> values of 8.8, 10.1, 16.9, and 10.1 μM, respectively.<sup>32</sup> The 7-methoxylated coumarin, micromelosidester (51), was isolated from the leaves of *M. falcatum*.<sup>16</sup> Two compounds microminin (52) and 6-methoxymicrominin (53) were isolated from the leaves of *M. falcatum*.<sup>4</sup> These 2 compounds are different from other compounds shown previously, because the prenyl attached on coumarin core forms a bicyclic structure. Other 2-prenylated coumarins named integerrimelin (54), (1*R*,3*R*,4*R*,6*S*)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxabicyclo[3.1.0]hexan-2-yl acetate (55), and scopolin (56) were isolated from the twigs, stems, and leaves of *M. integerrimum*.<sup>3,24,33</sup> A glycoside coumarin 7-methoxy-8-(1-hydroxy-2-*O*-β-glucopyranosyl-3-methyl-4-butene-1-yl) coumarin (57) was isolated from the stem bark of *M. falcatum*.<sup>29</sup> Another coumarin, micromelumoside D (58), was isolated from the aerial portions of *M. minutum*<sup>6</sup> (Figure 6).

In 2018, Kassim et al reported a lignan sesamin (118) (Figure 7) together with 3 coumarins, hydramicromelinin (59), micromelinin (60), and 1-*O*-β-D-glucopyranosylmarmesin (61)




Figure 7. Miscellaneous compounds from the *Micromelum integerrimum*, *M. birsutum*, *M. minutum*, and *M. falcatum*.

(Figure 8) from *M. minutum* methanol bark extract. Of the reported compounds, the ORAC measurements of compounds **59** and **118** showed potent antioxidant activity with the values of 5539 and 4031  $\mu\text{mol TE/g}$ , respectively.<sup>34</sup>

### Alkaloids

Alkaloids have inspired the development of several antibacterial drugs through the synthesis of quinine serendipitously yielding the quinolones, structural alteration of azomycin yielding metronidazole, and working with the quinoline scaffold yielding bedaquiline. In other drugs, alkaloids are present as scaffold substructures, eg linezolid and trimethoprim. Alkaloids remain the focus of much research. Their development as antibacterial drugs pursued within academia, industry, and joint-ventures.<sup>35</sup> A large number of carbazole alkaloids have been

isolated from higher plants of the genera *Murraya*, *Glycosmis*, *Micromelum*, and *Clausena*, all belonging to the family rutaceae.<sup>36,37</sup>

The quinoldione alkaloids 2-(3-hydroxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinolin-3-yl)acetate (**62**) and 3-hydroxy-1-methyl-3-(2-oxopropyl)quinoline-2,4(1H,3H)-dione (**63**) along with 2 quinolinone alkaloids *N*-methylflindersine (**64**), 4-hydroxy-3-methoxy-1-methyl-2(1H)-quinolinone (**65**), and the *N*-methylswietenidine-B (**66**) were reported from the stem of *M. falcatum* (Figure 9). The toxicity of all compounds (**62-66**) was tested in the brine shrimp larvae assay, and their  $\text{LD}_{50}$  values were 143, 355.0, 1.39, 2020.0, and 70.5 mg/mL, respectively, indicating that compound (**64**) was a potent toxic substance.<sup>17</sup>

The alkaloids myosmine (**67**), halfordinol (**68**), and *O*-methylhalfordinol (**69**) were reported from the leaves and




Figure 8. Coumarins from the *Micromelum minutum*.

young stems of *M. zeylanicum*.<sup>21</sup> The methyl carbazole-3-carboxylate (70) was isolated from the leaves of *M. integerrimum*<sup>38</sup> (Figure 9).

Increased death rates due to lung cancer have necessitated the search for potential novel anticancer compounds such as carbazole derivatives. Carbazoles are aromatic heterocyclic




Figure 9. Alkaloids from *Micromelum falcatum*, *M. zeylanicum*, and *M. integerrimum*.

organic compounds, with a tricyclic structure composed of 2 benzene rings each fused onto a 5-membered nitrogen-containing ring with anticancer, antibacterial, and anti-inflammatory activity.

Anti-TB bioassay-directed fractionation led to the isolation of 6 carbazole alkaloids from the  $\text{CH}_2\text{Cl}_2$  extract of the stem bark of *M. birsutum*. The carbazoles including 3-methylcarbazole (71), 3-formylcarbazole (72),<sup>3</sup> lansine (73), 3-formyl-6-methoxycarbazole (74), and methyl carbazole-3-carboxylate (75) were reported from the stem barks of *M. birsutum*.<sup>18</sup> A acridone alkaloid, 1,3-dihydroxy-4-methoxy-10-methylacridone (76), was isolated from the leaves of *M. integerrimum*<sup>38</sup> (Figure 10).

Two carbazole alkaloids, 2,7-dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole (77) and 7-methoxy heptaphylline (78), were isolated from the roots of *M. glanduliferum* by Siridechakorn et al in 1985. They might be a useful chemotaxonomic marker of the species *M. glanduliferum* and could be used to differentiate *M. glanduliferum* from other species of *Micromelum*.<sup>20</sup> Micromeline (79) was isolated from the stem barks of *M. birsutum*.<sup>18</sup> The alkaloid compounds koenimbine (80), koenine (81), koenigine (82), and koenidine (83) were isolated from the leaves and young stems of *M. zeylanicum*<sup>21</sup> (Figures 10 and 11).

In addition, a carbazole alkaloid, mahanine (84), was isolated from the leaves of *M. minutum* (Figure 11). It has been shown to exhibit a wide range of pharmacological effects including antimutagenicity against heterocyclic amines, antimicrobial activity against Gram positive bacteria, an anti-inflammatory effect, and it appeared to be cytotoxic against several cancer cell lines.<sup>35</sup>

Phenethyl cinnamides, micrometam A (85), micrometam D (86), micrometam E (87), and micrometam B (88), and a core of novel marine compound, micrometam C (89), were reported from the whole plant of *M. falcatum*.<sup>15,39</sup> (Figure 11).


### Flavones

A number of phytochemical studies dealing with polymethoxylated flavones from *Micromelum* spp. have been reported and a large number of flavonol-type metabolites were identified with the occurrence of partially methylated derivatives.<sup>1</sup>

Polymethoxylated flavones, 5-hydroxy-3,4',7,8-tetramethoxyflavone (90), 5,7-dihydroxy-3,4',8-trimethoxyflavone (91), 5-hydroxy-3,4',7,8-tetramethoxyflavone (92), and 5-hydroxy-3,4',6,7,8-pentamethoxyflavone (93), were reported from *M. minutum*.<sup>40</sup> Other polymethoxylated flavones such as 3,5,7,4'-tetramethoxyflavone (94), 3,5,7,8,4'-pentamethoxyflavone (95), and 3,5,6,7,4'-pentamethoxyflavone (96) were also reported from the leaves of *M. compressum*<sup>1</sup> (Figure 12).

Three compounds, 5,7-dihydroxy-4',6,8-trimethoxyflavone (97), 7-hydroxy-4',5,6,8-tetramethoxyflavone (98), and 7-hydroxy-4',6,8-trimethoxyflavone-5-yacetate (99), were isolated from the aerial parts of *M. minutum*.<sup>26</sup> 5-hydroxy-3,3',4',7,8-pen-





**Figure 10.** Alkaloids from *Micromelum minutum*, *M. birsutum*, *M. zeylanicum*, *M. glanduliferum*, *M. falcatum*, and *M. integerrimum*.

tamethoxyflavone (**100**) was also isolated from the leaves and stems of *M. zeylanicum*<sup>21</sup> (Figure 12).

#### Coumarin and Flavone Condensed Binary Compounds<sup>16</sup>

The compounds 8-((1*S*,2*R*)-1-hydroxy-2-(5-hydroxy-6, 8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-7-yl)oxy)-3-methylbut-3-enyl)-7-methoxy-2*H*-chromen-2-one (**101**),

8-((1*S*,2*R*)-1-hydroxy-3-methyl-2-(5,6,8-trimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-7-yl)oxy)but-3-enyl)-7-methoxy-2*H*-chromen-2-one (**102**), and 7-((1*S*,2*R*)-1-hydroxy-1-(7-methoxy-2-oxo-2*H*-chromen-8-yl)-3-methylbut-3-en-2-yl)oxy)-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4*H*-chromen-5-yl acetate (**103**) were reported from the above ground parts of *M. minutum*<sup>26</sup> (Figure 13).





Figure 11. Alkaloids from *Micromelum falcatum*.

### Miscellaneous Compounds


The stigmasterol (**104**) was isolated from the leaves of *M. minutum*.<sup>10</sup> Two triterpenes, 5(6)-gluten-3 $\alpha$ -ol (**105**) and 5(6)-gluten-3-on (**106**), were isolated from the leaves' and barks' crude petroleum ether and chloroform extracts of *M. minutum*.<sup>11</sup> The derivative of oleic acid with trivial name micromolide (**107**) was isolated from the stem barks of *M. hirsutum*<sup>18</sup> (Figure 14).

In 2014, Zhi-yao et al have reported 2 phenylpropanoids, microintegerrin A (**108**) and microintegerrin B (**109**), from the air-dried stems and leaves of *M. integerrimum*. These 2 compounds were tested for their cytotoxicity against 3 cancer cell lines (HeLa, A549, and BGC-823) but did not show any activity.<sup>24</sup> Dihydrocinnamic methyl ester derivatives, 3-(5-((2*R*, 3*R*, 4*R*)-3,4-dihydroxy-4-methyl-5-oxotetrahydrofuran-2-yl)-2-hydroxy-4-methoxyphenyl)




**Figure 12.** Flavones from *Micromelum minutum*, *M. zeylanicum*, and *M. compressum*.

propanoic acid (110) and secomicromelin (111), were isolated from the fruits of *M. falcatum*<sup>4</sup> (Figures 7 and 14). Three dihydrocinnamic acid derivatives, 3,4-dihydro-1,2-secomicrominin (112), 3,4-dihydro-1,2-secomicrominin methylester (113), and 3,4-dihydro-1,2-secomicrominin-9-O-glucoside<sup>9</sup> (114), were isolated from the leaves of *M. falcatum*.<sup>14</sup> Other 3



**Figure 14.** Miscellaneous compounds from *Micromelum minutum* and *M. birsutum*.

micromelumosides A (115), micromelumosides B (116), and micromelumosides C (117), and sesamin (118) were reported from the aerial parts of *M. minutum*<sup>6</sup> (Figure 7).



**Figure 13.** Coumarin and flavone condensed compounds from *Micromelum minutum*.

## Conclusion

*Micromelum* spp. possess very rich sources of bioactive secondary metabolites. As a member of rutaceae family it contains mainly coumarins (from simple to prenyl, and geranyl), alkaloids (quinolidine and carbazole), polymethoxyflavones, and a few glycosides and phenylpropanoic acid derivatives (Table 1). The coumarins are the most abundant compounds from *Micromelum* spp., and some compounds, such as 8-hydroxyisocapnolactone-2',3'-diol, have positive effects on the anticancer activity. Currently, at least 118 compounds have been reported from different plant parts of *M. integrimum*, *M. minutum*, *M. falcatum*, *M. compressum*, *M. glanduliferum*, *M. birsutum*, and *M. zeylanicum* (Table 1). Among them, *M. integrimum*, *M. minutum*, and *M. falcatum* were well studied chemically and pharmacologically and others, *M. compressum*, *M. glanduliferum*, *M. birsutum*, and *M. zeylanicum*, have been studied less. So, the review suggests that there is a great potential for researchers to search drug lead secondary metabolites from the genus *Micromelum*.

## Acknowledgments

Appreciation is conveyed to Prof. M. Iqbal Choudhary from H.E.J. Research Institute of Chemistry, International Center of Chemical and Biological Sciences, University of Karachi, Karachi-75270, Pakistan, for his help in improving this review article.

**Table 1.** Isolated Compounds Reported From *Micromelum* Spp.

No. of compounds	Plant parts	Name of species	References
<b>Simple coumarins</b>			
Umbelliferone (1)	52 Roots	<i>M. minutum</i>	8,13
7-Hydroxy-6-methoxy-2H-chromen-2-one (2)	Fruits	<i>M. minutum</i>	23
6-Hydroxy-7-methoxy-2H-chromen-2-one (3)	Leaves and stems	<i>M. integririmum</i>	24
6-Formyl-7-methoxycoumarin (4)	Leaves and stems	<i>M. zeylanicum</i>	21
Osthenon (5)	Roots	<i>M. minutum</i>	8
<b>Prenylated coumarins</b>			
Osthol (6)	Leaves	<i>M. minutum</i>	8
Microfalcrin (7)	Roots	<i>M. falcatum</i>	16
Micromarin-F (8)	Stems	<i>M. minutum</i>	25
Micromarin-G (9)	Stems	<i>M. minutum</i>	25
Micromarin-H (10)	Stems	<i>M. minutum</i>	25
Murralongin-1 (11)	Fruits	<i>M. minutum</i>	8,23
Murralongin-2 (12)	Fruits	<i>M. minutum</i>	8,23
Murralonginol (13)	Fruits	<i>M. minutum</i>	8,23
Phebalosin (14)	Roots	<i>M. minutum</i>	8
Murrangatin (15)	Roots	<i>M. minutum</i>	8
Murrangatin acetate (16)	Roots	<i>M. minutum</i>	8
Murracarpin (17)	Roots	<i>M. minutum</i>	8
Minumicrolin (18)	Roots	<i>M. minutum</i>	8
Microminutin (19)	Stem	<i>M. minutum</i>	25
7-Methoxy-8-(4'-methyl-3'-furyl)coumarin (20)	Fruits	<i>M. falcatum</i>	4
Microminutin B (21)	Fruits	<i>M. falcatum</i>	4
Microminutin C (22)	Fruits	<i>M. falcatum</i>	4
Micromeloside A (23)	Fruits	<i>M. falcatum</i>	4
Microcoumaririn (24)	Leaves	<i>M. falcatum</i>	16
Micromelin (25)	Fruits	<i>M. minutum</i>	23
Acetyldihydromicromelin A (26)	Aerial parts	<i>M. minutum</i>	26
Hydramicromelin D (27)	Twigs	<i>M. integririmum</i>	3
Hopeyhopol (28)	Whole plants	<i>M. minutum</i>	28
Marmesin (29)	Whole plants	<i>M. minutum</i>	28
(-) Decursinol (30)	Fruits	<i>M. falcatum</i>	4
Micromarin-A (31)	Stems	<i>M. falcatum</i>	16,22
Micromarin-B (32)	Stems	<i>M. minutum</i>	16,22
Micromarin-C (33)	Stems	<i>M. minutum</i>	22
(3S,4S)-3,4-Dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-ethylenebutyl 3-methylbutanoate (34)	Fruits	<i>M. falcatum</i>	4
(3R,4S)-3,4-Dihydroxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl 3-methylbutanoate (35)	Fruits	<i>M. falcatum</i>	4
3-Hydroxy-4-methoxy-4-(7-methoxy-2-oxo-2H-chromen-8-yl)-2-methylenebutyl 3-ethylbutanoate (36)	Fruits	<i>M. falcatum</i>	4
7-Methoxy-8-(2-hydroxyethyl-1-O-isovaleryl-4-butenyl) coumarin (37)	Stem bark	<i>M. falcatum</i>	29
Murralonginol isovalerate-1 (38)	Stems	<i>M. minutum</i>	23
Murralonginol isovalerate-2 (39)	Roots	<i>M. minutum</i>	8
Minutaminolate (40)	Roots	<i>M. minutum</i>	8
7-Demethylmurralonginol isovalerate (41)	Roots	<i>M. minutum</i>	23
Micromarinate (42)	Fruits	<i>M. falcatum</i>	4
8-Hydroxyisocapnolactone-2',3'-diol (43)	Leaves	<i>M. minutum</i>	10,11,28,32
8-Methoxycapnolactone (44)	Leaves	<i>M. minutum</i>	10,11,28,32
3",4"-Dihydrocapnolactone (45)	Leaves	<i>M. minutum</i>	10,11,28,32
2',3'-Epoxyisocapnolactone (46)	Leaves	<i>M. minutum</i>	10,11,28,32
8-Hydroxy-3",4"-dihydrocapnolactone-2',3'-diol (47)	Leaves	<i>M. minutum</i>	10,11,32
8,4-Dihydroxy-3,4-dihydro capnolactone-2,3-diol (48)	Leaves	<i>M. minutum</i>	10,11,32
Minutin A (49)	Leaves	<i>M. minutum</i>	10,11,32
Minutin B (50)	Leaves	<i>M. minutum</i>	10,11,32
Micromelosidester (51)	Leaves	<i>M. falcatum</i>	4

(Continued)



Table 1. Continued

No. of compounds	Plant parts	Name of species	References
Microminutinin (52)	Leaves	<i>M. falcatum</i>	4
6-Methoxymicrominutinin (53)	Leaves	<i>M. falcatum</i>	16,25
Integerrimelin (54)	Twigs	<i>M. integririmum</i>	3
(1R,3R,4R,6S)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxabicyclo[3.1.0]hexan-2-yl acetate (55)	Leaves	<i>M. integririmum</i>	33
scopolin (56)	Stems and leaves	<i>M. integririmum</i>	24
7-Methoxy-8-(1-hydroxy-2-O- $\beta$ -glucopyranosyl-3-Methyl-4-butene-1-yl) coumarin (57)	Stem bark	<i>M. falcatum</i>	29
Micromelumside D (58)	Aerial parts	<i>M. minutum</i>	6
Hydramicromelinin (59)	Bark	<i>M. minutum</i>	34
Micromelinin (60)	Bark	<i>M. minutum</i>	34
1-O- $\beta$ -D-Glucopyranosylmarmesin (61)	Bark	<i>M. minutum</i>	34
<b>Alkaloids</b>	60		
2-(3-Hydroxy-1-methyl-2,4-dioxo-1,2,3,4-tetrahydroquinolin-3-yl)acetate (62)	Stems	<i>M. falcatum</i>	17
3-Hydroxy-1-methyl-3-(2-oxopropyl)quinoline-2,4(1H,3H)-dione (63)	Stems	<i>M. falcatum</i>	17
N-Methylflindersine (64)	Stems	<i>M. falcatum</i>	17
4-Hydroxy-3-methoxy-1-methyl-2(1H)-quinolinone (65)	Stems	<i>M. falcatum</i>	17
N-Methylswietenidine-B (66)	Stems	<i>M. falcatum</i>	17
Myosmine (67)	Leaves and stem	<i>M. zeylanicum</i>	21
Halfordinol (68)	Leaves and stem	<i>M. zeylanicum</i>	21
O-Methylhalfordinol (69)	Leaves and stem	<i>M. zeylanicum</i>	21
Methyl carbazole-3-carboxylate (70)	Leaves	<i>M. integririmum</i>	38
3-Methylcarbazole (71)	Stem barks	<i>M. birsutum</i>	18
3-Formylcarbazole (72)	Stem barks	<i>M. birsutum</i>	18
Lansine (73)	Stem barks	<i>M. birsutum</i>	18
3-Formyl-6-methoxycarbazole (74)	Stem barks	<i>M. birsutum</i>	18
Methyl carbazole-3-carboxylate (75)	Leaves	<i>M. integririmum</i>	38
1,3-Dihydroxy-4-methoxy-10-methylacridone (76)	Stem bark	<i>M. birsutum</i>	18
2,7-Dihydroxy-3-formyl-1-(30-methyl-20-butenyl)carbazole (77)	Roots	<i>M. glanduliferlum</i>	20
7-Methoxy heptaphylline (78)	Roots	<i>M. glanduliferlum</i>	20
Micromeline (79)	Stem bark	<i>M. birsutum</i>	18
Koenimbine (80)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenine (81)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenigine (82)	Leaves and stem	<i>M. zeylanicum</i>	21
Koenidine (83)	Leaves and stem	<i>M. zeylanicum</i>	21
Mahanine (84)	Roots	<i>M. minutum</i>	35
<b>Miscellaneous compounds</b>			
Micrometam A (85)	Whole plants	<i>M. falcatum</i>	15
Micrometam D (86)	Whole plants	<i>M. falcatum</i>	15
Micrometam E (87)	Whole plants	<i>M. falcatum</i>	15
Micrometam B (88)	Whole plants	<i>M. falcatum</i>	15
Micrometam C (89)	Whole plants	<i>M. falcatum</i>	15
<b>Polymethoxylated flavones</b>	30		
5-Hydroxy-3,4',7,8-tetramethoxyflavone (90)	Aerial parts	<i>M. minutum</i>	40
5,7-Dihydroxy-3,4',8-trimethoxyflavone (91)	Aerial parts	<i>M. minutum</i>	40
5-Hydroxy-3,4',7,8-tetramethoxyflavone (92)	Aerial parts	<i>M. minutum</i>	40
5-Hydroxy-3,4',6,7,8-pentamethoxyflavone (93)	Leaves	<i>M. minutum</i>	40
3,5,7,4'-Tetramethoxyflavone (94)	Leaves	<i>M. ompresum</i>	1
3,5,7,8,4'-Pentamethoxyflavone (95)	Leaves	<i>M. ompresum</i>	1

(Continued)

**Table 1.** Continued

No. of compounds	Plant parts	Name of species	References
3,5,6,7,4'-Pentamethoxyflavone (96)	Leaves	<i>M. compressum</i>	1
5,7-Dihydroxy-4',6,8-trimethoxyflavone (97)	Aerial parts	<i>M. minutum</i>	26
7-Hydroxy-4',5,6,8-tetramethoxyflavone (98)	Aerial parts	<i>M. minutum</i>	26
7-Hydroxy-4',5,6,8-trimethoxyflavone-5-ylacetate (99)	Aerial parts	<i>M. minutum</i>	26
5-Hydroxy-3,3',4',7,8-pentamethoxyflavone (100)	Leaves and stems	<i>M. zeylanicum</i>	21
<b>Coumarin and flavone condensed compounds</b>			
8-((1 <i>S</i> ,2 <i>R</i> )-1-Hydroxy-2-(5-hydroxy-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4 <i>H</i> -chromen-7-ylloxy)-3-methylbut-3-enyl)-7-methoxy-2 <i>H</i> -chromen-2-one (101)	Aerial parts	<i>M. minutum</i>	26
8-((1 <i>S</i> ,2 <i>R</i> )-1-Hydroxy-3-methyl-2-(5,6,8-trimethoxy-2-(4-methoxyphenyl)-4-oxo-4 <i>H</i> -chromen-7-ylloxy)but-3-enyl)-7-methoxy-2 <i>H</i> -chromen-2-one (102)	Aerial parts	<i>M. minutum</i>	26
7-((1 <i>S</i> ,2 <i>R</i> )-1-Hydroxy-1-(7-methoxy-2-oxo-2 <i>H</i> -chromen-8-yl)-3-methylbut-3-en-2-ylloxy)-6,8-dimethoxy-2-(4-methoxyphenyl)-4-oxo-4 <i>H</i> -chromen-5-yl acetate (103)	Aerial parts	<i>M. minutum</i>	26
<b>Stigmasterol and triterpenes</b>			
Stigmasterol (104)	Leaves	<i>M. minutum</i>	10
5(6)-Gluten-3 $\alpha$ -ol (105)	Leaves and barks	<i>M. minutum</i>	23
5(6)-Gluten-3-one (106)	Leaves and barks	<i>M. minutum</i>	23
<b>Derivative of oleic acid and phenylpropanoic acid</b>			
Micromolide (107)	Stem barks	<i>M. hirsutum</i>	18
Microintegerrin A (108)	stems and leaves	<i>M. integerrimum</i>	24
Microintegerrin B (109)	stems and leaves	<i>M. integerrimum</i>	24
3-(5-((2 <i>R</i> ,3 <i>R</i> ,4 <i>R</i> )-3,4-Dihydroxy-4-methyl-5-oxotetrahydrofuran-2-yl)-2-hydroxy-4-methoxyphenyl)propanoic acid (110)	fruits	<i>M. falcatum</i>	4
Secomicromelin (111)	fruits	<i>M. falcatum</i>	4
3,4-Dihydro-1,2-secomicrominin (112)	leaves	<i>M. falcatum</i>	14
3,4-Dihydro-1,2-secomicrominin methylester (113)	leaves	<i>M. falcatum</i>	14
3,4-Dihydro-1,2-secomicrominin-9- <i>O</i> -glucoside (114)	leaves	<i>M. falcatum</i>	14
Micromelumosides A (115)	aerial parts	<i>M. minutum</i>	6
Micromelumosides B (116)	aerial parts	<i>M. minutum</i>	6
Micromelumosides C (117)	aerial parts	<i>M. minutum</i>	6
Sesamin (118)	bark	<i>M. minutum</i>	34

4


**Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**Funding**

The author(s) received no financial support for the research, authorship, and/or publication of this article.

**ORCID ID**

Nanik Siti Aminah  <https://orcid.org/0000-0002-2767-6006>

**References**

- Vidar WS, Macabeo APG, Knom M, Kohls P, Aguinakdo AM. Polymethoxylated flavones from *Micromelum compressum*. *Biochem Syst Ecol*. 2013;50:48-50. doi:10.1016/j.bse.2013.03.010
- Singh RK. Lectotypification of the name *Micromelum integerrimum* (Rutaceae). *Phytotaxa*. 2016;255(2):172-174. doi:10.11646/phytotaxa.255.2.7
- Phakhodee W, Pattarawarapan M, Pongpam P, Laphookhieo S. Naturally occurring prenylated coumarins from *Micromelum integerrimum* twigs. *Phytochem Lett*. 2014;7:165-168. doi:10.1016/j.phytol.2013.11.009
- Suthiwong J, Sriphana U, Thongsri Y, Promsuwan P, Prariyachattigul C, Yenjai C. Coumarinoids from the fruits of *Micromelum falcatum*. *Fitoterapia*. 2014;94:134-141. doi:10.1016/j.fitote.2014.02.004
- Susidarti RA. Chemical Constituents and Biological Activities of *Micromelum Minutum* (Rutaceae) and Two Eugenia Species (Myrtaceae). [PhD thesis]. Universiti Putra Malaysia; 2003.
- Sahakitpichan P, Tanpatanan W, Chimnoi N, Ruchirawat S, Kanchanapoom T. Glucosides of phenylpropanoic acid derivatives and coumarins from *Micromelum minutum*. *Phytochem Lett*. 2015;14:12-16. doi:10.1016/j.phytol.2015.08.009

7. Zhang D, Hartley TG, Mabberley DJ. *Flora of China, FOC* ; 1998;11: 51.
8. Lekphrom R, Kanokmedhakul K, Sangsopha W, Kanokmedhakul S. A new coumarin from the roots of *Micromelum minutum*. *Nat Prod Res*. 2016;30(21):2383-2388. doi:10.1080/14786419.2016.1185720
9. Tantishaiyakul V, Pummangura S, Chaichantipwath C, McLaughlin JL, Division LS. phebalsin from the stem bark of *Micromelum minutum*. *J Nat Prod*;49(1):49-50.
10. Susidarti R, Rahmani M, Ali A. 8-Methoxycapnolactone and stigmaterol From *Micromelum minutum*. *Maj Farm*. 2007;18(2):105-109.
11. Asmah Susidarti R, Rahmani M, Ismail HBM, et al. Cytotoxic activity of coumarins from *Micromelum minutum*. *Pharm Biol*. 2009;47(2):182-185. doi:10.1080/13880200802436513
12. Hasan CM, Ahsan M. Chemical constituents, cytotoxic activities and traditional uses of *Micromelum minutum* (Rutaceae): a review. *Pharm Pharmacol Int J*. 2019;7(5):229-236.
13. Ramesh B, Pugalendi KV. Antioxidant role of umbelliferone in STZ-diabetic rats. *Life Sci* 2006;79(3):306-310. doi:10.1016/j.lfs.2006.01.005
14. Kamperdick C, Phuong NM, Sung TV, Schmidt J, Adam G. Coumarins and dihydrocinnamic acid derivatives from *Micromelum falcatum*. *Phytochemistry*. 1999;52(8):1671-1676. doi:10.1016/S0031-9422(99)00243-5
15. Luo X, Huang Y, Zhang S, Yin H, Li C, Li Q. Five new phenethyl cinnamides from the mangrove associates *Micromelum falcatum*. *Biochem Syst Ecol*. 2014;56:191-195. doi:10.1016/j.bse.2014.05.018
16. Kouloura E, Danika E, Kim S, et al. Rapid identification of coumarins from *Micromelum falcatum* by UPLC-HRMS/MS and targeted isolation of three new derivatives. *Molecules*. 2014;19(9):15042-15057. doi:10.3390/molecules190915042
17. Luo XM, Qi SH, Yin H, Gao CH, Zhang S. Alkaloids from the stem bark of *Micromelum falcatum*. *Chem Pharm Bull*. 2009;57(6):600-602. doi:10.1248/cpb.57.600
18. Ma C, Case RJ, Wang Y, et al. Anti-Tuberculosis constituents from the stem bark of *Micromelum hirsutum*. *Planta Med*. 2005;71(3):261-267. doi:10.1055/s-2005-837826
19. Rodphukdeekul S, Sueblinvong T, Reungrungsi N, Chansrinoyom C. Anti-tumor activity of *Micromelum hirsutum* extract. *Thai J Pharmacol*. 2010;32(1):106-110.
20. Siridechakorn I, Ritthiwigrom T, Laphookhieo S. Coumarins and carbazole alkaloids from the roots of *Micromelum glanduliferum*. *Biochem Syst Ecol*. 2012;40:69-70. doi:10.1016/j.bse.2011.10.002
21. Bowen IH, Perera KPWC. Alkaloids, coumarins and flavonoids of *Micromelum zeylanicum*. *Phytochemistry*. 1982;21(2):433-437. doi:10.1016/S0031-9422(00)95283-X
22. Koriem KMM, Aminuddin ME, Kader AS. Antihyperglycemic SNR. Antihyperlipidemic and Antiapoptotic Activities of *Micromelum minutum* Seeds in Diabetic Rats. *J Mol Genet Med*. 2013
23. Lekphrom R, Kanokmedhakul S, Kukongviriyapan V, Kanokmedhakul K. C-7 oxygenated coumarins from the fruits of *Micromelum minutum*. *Arch Pharm Res*. 2011;34(4):527-531. doi:10.1007/s12272-011-0402-y
24. Wang Z-Y, He W-J, Zhou W-B, et al. Two new phenylpropanoids from *Micromelum integerrimum*. *Chin J Nat Med*. 2014;12(8):619-622. doi:10.1016/S1875-5364(14)60094-7
25. Ito C, Otsuka T, Ruangrunsi N, Furukawa H. Chemical constituents of *Micromelum minutum*. Isolation and structural elucidation of new coumarins. *Chem Pharm Bull*. 2000;48(3):334-338. doi:10.1248/cpb.48.334
26. Das S, Baruah RH, Sharma RP, Barua JN, Kulanthaivel P, Herz W. 7-methoxycoumarins from *Micromelum minutum*. *Phytochemistry*. 1984;23(10):2317-2321. doi:10.1016/S0031-9422(00)80543-9
27. Yun-Cheung Ket al. *Micromelum*: a key genus in the chemosystematics of the Clausenea. *Biochem Syst Ecol*. 1988;16(5):485-489.
28. Rahman MS, Rahman MZ, Hasan CM, Rashid MA, Kaiser MA. A review on phytochemicals from some medicinal plants of Bangladesh. *J. Phar. Nutri. Sci*. 2011;1(1):87-95. doi:10.6000/1927-5951.2011.01.01.14
29. Luo X, He W, Yin H, et al. Two new coumarins from *Micromelum falcatum* with cytotoxicity and brine shrimp larvae toxicity. *Molecules*. 2012;17(6):6944-6952. doi:10.3390/molecules17066944
30. Rahmani M, Susidarti RA, Ismail HBM, et al. Coumarins from Malaysian *Micromelum minutum*. *Phytochemistry*. 2003;64(4):873-877. doi:10.1016/j.phytochem.2003.08.007
31. Susidarti RA, Jenie RI, Ikawati M, Putri DDP, Meiyanto E. Cytotoxic activity and apoptosis induction of 8-hydroxyisocapnolactone-2',3'-diol and its combination with Doxorubicin on MCF-7 and T47D cells. *J Appl Pharm Sci*. 2014;4(6):089-097.
32. Sakunpak A, Matsunami K, Otsuka H, Panichayupakaranant P. Isolation of new monoterpene coumarins from *Micromelum minutum* leaves and their cytotoxic activity against Leishmania major and cancer cells. *Food Chem*. 2013;139(1-4):458-463. doi:10.1016/j.foodchem.2013.01.031
33. Phakhodee W, Laphookhieo S, Prior TJ, Rujiwatra A. 1R,3R,4R,6S)-4-(7-Methoxy-2-oxo-2H-chromen-6-yl)-1-methyl-3,6-dioxo-bicyclo-[3.1.0]hexan-2-yl acetate. *Acta Crystallogr Sect E Struct Rep Online*. 2012;68(Pt 12):3421-3422. doi:10.1107/S1600536812047617
34. Kassim NK, Lim PC, Ismail A, Awang K. Isolation of antioxidative compounds from *Micromelum minutum* guided by preparative thin layer chromatography-2,2-diphenyl-1-picrylhydrazyl (PTLC-DPPH) bioautography method. *Food Chem*. 2019;272:185-191. doi:10.1016/j.foodchem.2018.08.045
35. Roy MK, Thalang VN, Trakoontivakorn G, Nakahara K. Mechanism of mahanine-induced apoptosis in human leukemia cells (HL-60). *Biochem Pharmacol*. 2004;67(1):41-51. doi:10.1016/j.bcp.2003.07.021
36. Schmidt AW, Reddy KR, Knölker H-J, Occurrence KH-J. Occurrence, biogenesis, and synthesis of biologically active carbazole alkaloids. *Chem Rev*. 2012;112(6):3193-3328. doi:10.1021/cr200447s
37. Chakraborty DP, Roy S. Carbazole alkaloids. III. *Fortschr Chem Org Naturst*. 1991;57:71-152. doi:10.1007/978-3-7091-9119-4\_2
38. Yang X-L, Xie Z-H, Jiang X-J, Huang Y-B, Liu J-K, Yan-Bing Huang J-KL. A new acridone alkaloid from *Micromelum integerrimum*. *Chem Pharm Bull*. 2009;57(7):734-735. doi:10.1248/cpb.57.734

39. Tang H, Ge H, Chen Z-B, et al. Micrometam C protects against oxidative stress in inflammation models in zebrafish and RAW264.7 macrophages. *Mar Drugs*. 2015;13(9):5593-5605. doi: 10.3390/md13095593
40. Sohrab MH, Chowdhury R, Hasan CM, Rashid MA. Chemotaxonomic significance of polyoxygenated flavonoids from the leaves of *Micromelum minutum*. *Biochem Syst Ecol*. 2004;32(9):829-831. doi: 10.1016/j.bse.2003.11.009



# Phytoconstituents of Genus *Micromelum* and Their Bioactivity—A Review

## ORIGINALITY REPORT

25%

SIMILARITY INDEX

15%

INTERNET SOURCES

23%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Hoong-Kun Fun, Surat Laphookhieo, Wisanu Maneerat, Suchada Chantrapromma. " 2,8-Dihydroxy-1-(3-methylbut-2-enyl)-9 -carbazole-3-carbaldehyde ", Acta Crystallographica Section E Structure Reports Online, 2007  
Publication 1%
- 2 [psasir.upm.edu.my](http://psasir.upm.edu.my)  
Internet Source 1%
- 3 Youn, S. W., H. K. Kim, H. J. Lee, and J. Lee. "Quantification of cerebral circulation and shunt volume in a tentorial dural arteriovenous fistula using two-dimensional phase-contrast magnetic resonance imaging", Acta Radiologica Short Reports, 2014.  
Publication 1%
- 4 [epub.wu.ac.at](http://epub.wu.ac.at)  
Internet Source 1%
- 5 [www.medcraveonline.com](http://www.medcraveonline.com)  
Internet Source 1%

6

Tin Myo Thant, Nanik Siti Aminah, Alfinda Novi Kristanti, Rico Ramadhan, Preecha Phuwapraisirisan, Yoshiaki Takaya. " A new pyrano coumarin from roots displaying dual inhibition against  $\alpha$ -glucosidase and free radical ", Natural Product Research, 2019

Publication

1%

7

[thehomeopathicremedies.blogspot.com](http://thehomeopathicremedies.blogspot.com)

Internet Source

1%

8

[www.medscape.com](http://www.medscape.com)

Internet Source

1%

9

Wong Phakhodee, Mookda Pattarawarapan, Preetiya Pongparn, Surat Laphookhieo. "Naturally occurring prenylated coumarins from *Micromelum integerrimum* twigs", Phytochemistry Letters, 2014

Publication

1%

10

P. Valenti, A. Rampa, R. Budriesi, A. Bisi, A. Chiarini. "Coumarin 1,4-Dihydropyridine Derivatives", Bioorganic & Medicinal Chemistry, 1998

Publication

1%

11

Kazuhiko Nakahara, Gassinee Trakoontivakorn, Najeeb S. Alzoreky, Hiroshi Ono, Mayumi Onishi-Kameyama, Mitsuru Yoshida. " Antimutagenicity of Some Edible Thai Plants, and a Bioactive Carbazole Alkaloid, Mahanine,

1%

Isolated from ", Journal of Agricultural and Food Chemistry, 2002

Publication

---

12

A. Sakunpak, K. Matsunami, H. Otsuka, P. Panichayupakaranant. "Isolation of new monoterpene coumarins from *Micromelum minutum* leaves and their cytotoxic activity against *Leishmania major* and cancer cells", Food Chemistry, 2013

Publication

---

13

Mohammad H Sohrab, Rasheduzzaman Chowdhury, Choudhury M Hasan, Mohammad A Rashid. "Chemotaxonomic significance of polyoxygenated flavonoids from the leaves of *Micromelum minutum*", Biochemical Systematics and Ecology, 2004

Publication

---

14

Zhi-Yao WANG, Wen-Jun HE, Wen-Bing ZHOU, Guang-Zhi ZENG, Zhi-Qi YIN, Shou-Xun ZHAO, Ning-Hua TAN. "Two new phenylpropanoids from *Micromelum integerrimum*", Chinese Journal of Natural Medicines, 2014

Publication

---

15

[www.fluorochem.co.uk](http://www.fluorochem.co.uk)

Internet Source

---

16

[www.genome.jp](http://www.genome.jp)

Internet Source

---

1%

1%

1%

1%

<1%

17 Shu-Ming Li. "Prenylated indole derivatives from fungi: structure diversity, biological activities, biosynthesis and chemoenzymatic synthesis", Nat. Prod. Rep., 2010

Publication

<1%

18 Kazuhiko Nakahara. "Mahanine, a carbazole alkaloid from *Micromelum minutum*, inhibits cell growth and induces apoptosis in U937 cells through a mitochondrial dependent pathway", British Journal of Pharmacology, 05/2005

Publication

<1%

19 Refilwe P. Molatlhegi, Alisa Phulukdaree, Krishnan Anand, Robert M. Gengan, Charlette Tiloke, Anil A. Chuturgoon. "Cytotoxic Effect of a Novel Synthesized Carbazole Compound on A549 Lung Cancer Cell Line", PLOS ONE, 2015

Publication

<1%

20 [researchspace.ukzn.ac.za](http://researchspace.ukzn.ac.za)

Internet Source

<1%

21 Yun-Cheung Kong, Paul P-H. But, Kam-hung Ng, Qian Li, Kin-Fai Cheng, Peter G. Waterman. "Micromelum: a key genus in the chemosystematics of the Clauseneae", Biochemical Systematics and Ecology, 1988

Publication

<1%

22 Xiao-Fei Shang, Susan L. Morris-Natschke, Guan-Zhou Yang, Ying-Qian Liu et al.

<1%



"Biologically active quinoline and quinazoline alkaloids part II", Medicinal Research Reviews, 2018

Publication

23

Harald Greger. "Phytocarbazoles: alkaloids with great structural diversity and pronounced biological activities", Phytochemistry Reviews, 2017

Publication

<1%

24

[www.japsonline.com](http://www.japsonline.com)

Internet Source

<1%

25

[citeseerx.ist.psu.edu](http://citeseerx.ist.psu.edu)

Internet Source

<1%

26

[indonesianjpharm.farmasi.ugm.ac.id](http://indonesianjpharm.farmasi.ugm.ac.id)

Internet Source

<1%

27

Belluti, F.. "Design, synthesis and anticancer activities of stilbene-coumarin hybrid compounds: Identification of novel proapoptotic agents", Bioorganic & Medicinal Chemistry, 20100515

Publication

<1%

28

[www.eflora.cn](http://www.eflora.cn)

Internet Source

<1%

29

Poolsak Sahakitpichan, Waraporn Tanpatanan, Nitirat Chimnoi, Somsak Ruchirawat, Tripetch Kanchanapoom. "Glucosides of

<1%

phenylpropanoic acid derivatives and coumarins  
from *Micromelum minutum*", *Phytochemistry  
Letters*, 2015

Publication

30

[www.scribd.com](http://www.scribd.com)

Internet Source

<1%

31

[pt.scribd.com](http://pt.scribd.com)

Internet Source

<1%

32

[spatialcomplexity.blogweb.casa.ucl.ac.uk](http://spatialcomplexity.blogweb.casa.ucl.ac.uk)

Internet Source

<1%

33

[git.embl.de](http://git.embl.de)

Internet Source

<1%

34

Ahmad Alkofahi. "Mutagenic and Toxic Activity  
of some Jordanian Medicinal Plants",  
*Pharmaceutical Biology*, 3/1/1995

Publication

<1%

35

Maria de Fátima Agra, Patrícia França de  
Freitas, José Maria Barbosa-Filho. "Synopsis of  
the plants known as medicinal and poisonous in  
Northeast of Brazil", *Revista Brasileira de  
Farmacognosia*, 2007

Publication

<1%

36

Elodie Faurel-Paul, Kenta Yoshida, Magali  
Sépulcre, Hamid Dhimane, Yves Le Merrer.  
"Synthesis of Three Regioisomeric (7-  
Methoxychromenonyl)methyl Guanosine 5'-

<1%

37

[eprints.um.edu.my](http://eprints.um.edu.my)

Internet Source

<1%

38

Christiane Schultze, Bernd Schmidt.  
"Prenylcoumarins in One or Two Steps by a  
Microwave-Promoted Tandem Claisen  
Rearrangement/Wittig Olefination/Cyclization  
Sequence", The Journal of Organic Chemistry,  
2018

Publication

<1%

39

Xing-Wei Yang, Robert B. Grossman, Gang Xu.  
"Research Progress of Polycyclic  
Polyprenylated Acylphloroglucinols", Chemical  
Reviews, 2018

Publication

<1%

40

[jstagebeta.jst.go.jp](http://jstagebeta.jst.go.jp)

Internet Source

<1%

41

Longhuo Wu, Pei Li, Xiao Wang, Zhenjing  
Zhuang, Farzin Farzaneh, Ruian Xu. "  
Evaluation of anti-inflammatory and  
antinociceptive activities of ", Pharmaceutical  
Biology, 2010

Publication

<1%

42

Kouloura, Eirini, Eirini Danika, Sothea Kim,  
Mélanie Hoerlé, Muriel Cuendet, Maria  
Halabalaki, and Leandros Skaltsounis. "Rapid

<1%

Identification of Coumarins from *Micromelum falcatum* by UPLC-HRMS/MS and Targeted Isolation of Three New Derivatives", *Molecules*, 2014.

Publication

---

43

Nolan, Karen A., Jeremy R. Doncaster, Mark S. Dunstan, Katherine A. Scott, A. David Frenkel, David Siegel, David Ross, John Barnes, Colin Levy, David Leys, Roger C. Whitehead, Ian J. Stratford, and Richard A. Bryce. "Synthesis and Biological Evaluation of Coumarin-Based Inhibitors of NAD(P)H: Quinone Oxidoreductase-1 (NQO1)<sup>†</sup>", *Journal of Medicinal Chemistry*, 2009.

<1%

Publication

---

44

[www.molport.com](http://www.molport.com)

Internet Source

<1%

---

45

Arvind S. Negi. "Antitubercular potential of plants: A brief account of some important molecules", *Medicinal Research Reviews*, 2009

<1%

Publication

---

46

Peter Winterhalter. "Glycoconjugated aroma compounds: Occurrence, role and biotechnological transformation", *Advances in Biochemical Engineering/Biotechnology*, 1997

<1%

Publication

---

Temple. "Alkyl- or Aryl-Trisubstituted 1,2,4-

- |    |                                                                                                                                                                   |     |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| 47 | Triazoles", Chemistry of Heterocyclic Compounds A Series Of Monographs, 01/01/1981<br>Publication                                                                 | <1% |
| 48 | Ratna Asmah Susidarti. "A new coumarin and triterpenes from Malaysian Micromelum minutum", Natural Product Research, 2/8/2006<br>Publication                      | <1% |
| 49 | <a href="http://www.ifrpd.ku.ac.th">www.ifrpd.ku.ac.th</a><br>Internet Source                                                                                     | <1% |
| 50 | <a href="http://flora.huh.harvard.edu">flora.huh.harvard.edu</a><br>Internet Source                                                                               | <1% |
| 51 | <a href="http://www.digitalnaturalhistory.com">www.digitalnaturalhistory.com</a><br>Internet Source                                                               | <1% |
| 52 | Alexandre Ferraz. "Anti-tumour Screening of Brazilian Plants", Pharmaceutical Biology (Formerly International Journal of Pharmacognosy), 12/1/2002<br>Publication | <1% |
| 53 | <a href="http://stuartxchange.com">stuartxchange.com</a><br>Internet Source                                                                                       | <1% |
| 54 | <a href="http://archive.org">archive.org</a><br>Internet Source                                                                                                   | <1% |
| 55 | Ratna Asmah Susidarti, Mawardi Rahmani, Hazar B. M. Ismail, M. Aspollah Sukari et al. " A                                                                         | <1% |



new coumarin and triterpenes from Malaysian ",  
Natural Product Research, 2006

Publication

---

56

Yanfei Ban, Tianshuang Xia, Rui Jing, Yaoli Guo, Yiya Geng, Qi Ye, Chengjian Zheng. "Vitex Diterpenoids: Structural Diversity and Pharmacological Activity", Current Pharmaceutical Design, 2020

Publication

---

57

[www.globinmed.com](http://www.globinmed.com)

Internet Source

---

58

Wei-Lun Huang, Arun Raja, Bor-Cherng Hong, Gene-Hsiang Lee. " Organocatalytic Enantioselective Michael–Acetalization–Reduction–Nef Reaction for a One-Pot Entry to the Functionalized Aflatoxin System. Total Synthesis of (–)- Dihydroaflatoxin D and (–)- and (+)-Microminutinin ", Organic Letters, 2017

Publication

---

59

Taylor A. Choi. "Transition metals in organic synthesis - Part 83#: Synthesis and pharmacological potential of carbazoles", Medicinal Chemistry Research, 06/2008

Publication

---

60

Mariana Meckes. "A microbiological evaluation of medicinal plants used by the Maya people of Southern Mexico", Phytotherapy Research,

<1%

<1%

<1%

<1%

<1%

06/1995

Publication

---

61

Bernd Schmidt, Stefan Krehl, Alexandra Kelling, Uwe Schilde. " Synthesis of 8-Aryl-Substituted Coumarins Based on Ring-Closing Metathesis and Suzuki–Miyaura Coupling: Synthesis of a Furyl Coumarin Natural Product from ", The Journal of Organic Chemistry, 2012

<1%

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

# Phytoconstituents of Genus *Micromelum* and Their Bioactivity—a Review

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---

PAGE 10

---

PAGE 11

---

PAGE 12

---

PAGE 13

---

PAGE 14

---

PAGE 15

---