


VOLUME 49
ISSUE 6, 2014



ISSN 1314-7471 (print)
ISSN 1314-7978 (on line)
www.uclm.edu

JOURNAL

**OF CHEMICAL TECHNOLOGY
AND METALLURGY**



SOFIA 2014

CONFUSARIN AND NUDOL, TWO PHENANTHRENE GROUP COMPOUNDS, FROM *Dioscorea esculenta* L. AND THEIR ANTIOXIDANT ACTIVITIES

Nanik Siti Aminah, Ratih Hidayah, Mulyadi Tanjung

Dept. of Chemistry, Faculty of Science and Technology, Universitas Airlangga
Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, East Java, Indonesia
E-mail: nanik-s-a@fst.unair.ac.id

Received 05 January 2017

Accepted 20 July 2017

ABSTRACT

Two phenanthrene group compounds namely: confusarin and nudol are isolated from *Dioscorea esculenta* L. This species belongs to Dioscoreaceae family. The two compounds extraction is conducted by the maceration method using methanol solvent and partition with *n*-hexane and ethyl acetate. The ethyl acetate extract is purified using various chromatographic techniques yielding pure compounds. The latter structure is determined by spectroscopic methods including UV/Vis, IR, 1D and 2D NMR. Confusarin and nudol are tested for their antioxidant activity against DPPH radical scavenging. They show IC_{50} of $19,63 \pm 0,09$ and $37,91 \pm 0,08$ ppm, correspondingly.

Keywords: phenanthrene, confusarin, nudol, *Dioscorea esculenta* L., antioxidant.

INTRODUCTION

Dioscorea esculenta L. with the local name "gembili" is one of species belonging to *Dioscorea* genus. In Indonesia, this plant is one of species used by the community as a substitute for rice [1]. *Dioscorea*, a genus in Dioscoreaceae family, is comprised of about 600 species. It has been reported that various kinds of phenolic compounds are contained in *Dioscorea* genus, such as: stilbenoid, dihydrostilbene, phenanthrene, dihydrophenanthrene, diarylheptanoid and acetophenone [2 - 6]. Secondary metabolite of compounds from *Dioscorea* genus show [7-11] an interesting biological activity, such as: antiinflammatory, antibacterial, antimicrobial, anticancer, antiallergenic, antifungal and antioxidant ones.

Based on previous studies, the methanol extract of *D. esculenta* shows an antioxidant activity [12]. But a phytochemical study of the secondary metabolite compounds as well as an antioxidant activity of pure compounds from this species has not been so far reported. The present communication presents two phenolic compounds, namely confusarin (1) and nudol (2) isolated from *D. esculenta* and verifies their

antioxidant activity using DPPH reagent.

EXPERIMENTAL

Gravitation column chromatography (GCC) was carried out using Merck Si gel 60 (700-200 mesh), radial chromatography was conducted out using Merck Si gel 60 PF₂₅₄, while pre-coated Si gel plates (Merck Kieselgel 60 GF₂₅₄, 0.25 mm) were used for TLC analysis. UV and IR spectra were recorded with a Shimadzu UV-1800 and FT IR Spectrum One Perkin-Elmer instruments, respectively. ¹H and ¹³C-NMR spectra were obtained with JEOL ECA 400, operating at 400 (¹H) and 100 (¹³C) MHz, using residual and deuterated solvent peaks as reference standards. The determination of the antioxidant activity was based on the inhibition of free radical against DPPH using a spectroscopy method.

Samples of *Dioscorea esculenta* tuber were collected from Pengampon District, Kabuh, Jombang, East Java, Indonesia. The plant was identified by the staff of the Herbarium Bogoriense, Bogor Botanical Garden, Bogor, Indonesia, and a voucher specimen were deposited in the herbarium.

The powdered tuber of *Dioscorea esculenta* L. (10,0 kg) was maserated with methanol for 2 x 24 h and then concentrated under reduced pressure to give a gummy brownish extract. The methanol extract was partitioned into n-hexane and ethyl acetate soluble fractions. A portion (5.2 g) of the ethyl acetate fraction was fractionated using GLC (n-hexane : ethyl acetate, increasing polarity) to give two group fractions A & B by combining fractions with similar TLC profiles. Fraction B (159 mg) was subjected to further partition using radial chromatography (n-hexane:chloroform = 8:2, 7:3, 1:1, 3:7, chloroform and chloroform:ethyl acetate = 9:1). It gave three sub-fractions B1, B2, and B3. The purification of B2 fraction by radial chromatography (n-hexane : chloroform = 7:3, 1:1, 3:7 and chloroform) gave compound **1** (3,3 mg). Fraction B3 yielded compound **2** (19.6 mg) using the same methods (n-hexane : ethyl acetate = 9:1 and 8:2 as eluent), 1,1-diphenyl-2-picryl-hydrazyl (DPPH) scavenging activity assay was used to determine compound's inhibition capacity. Its reaction principle based on the mechanism of free radicals inhibition by hydrogen transfer provided the determination of the sample antioxidant activity expressed in IC₅₀ (Inhibition Concentration 50 %). The

procedure used was as follows: a total of 500 µL of the test solutions of a varying concentration (10 ppm - 500 ppm), 500 µL of 0.2 M acetate buffer of pH 5.5, and 1000 µL of methanol were mixed in a test tube; 500 µL of 5x10⁻⁴ M DPPH were added to this mixture; the latter was homogenized using a vortex in a dark room (resistant to UV light) and incubated for 30 min. Then the mixture UV absorbance was measured by a spectrophotometer at λ_{max} of 517 nm. Ascorbic acid was used as a positive control [13]. To prevent the sample from light disturbance, the test tube wrapped with aluminum foil. The inhibition capacity was evaluated in correspondence with the following equation:

$$\% \text{ inhibition} = \frac{A. \text{blanko} - A. \text{sampel}}{A. \text{blanko}} \times 100\%$$

RESULTS AND DISCUSSION

Confusarin (**1**) – an yellow powder, mp 220.8°C (dec.); UV spectrum (MeOH) λ_{max} nm (log ε) : 264 (4.41), 286 (3.82), 308 (3.64), 350 (2.94) and 367 nm (2.97); spectrum ¹H-NMR (CDCl₃, 400 MHz) and ¹³C-NMR (CDCl₃, 100 MHz) - see Table 1.

Table 1. ¹H-NMR and ¹³C-NMR data (CDCl₃) of compounds **1** and **2**.

No	Compound 1		Compound 2	
	δ _H	δ _C	δ _H	δ _C
1	7.19(s)	108.1	7.18(s)	108.2
2	-	142.3	-	147.5
3	-	136.1	-	140.9
4	-	145.4	-	150.7
4a	-	114.4	-	133.5
4b	-	119.8	-	118.9
5	9.19 (d, J=9.2)	123.9	9.36(d, J=9.1)	128.5
6	7.29(d, J=9.2)	116.1	7.19(dd, J=9.1, 2.8)	116.5
7	-	140.9	-	153.2
8	-	135.1	7.16(d, J=2.8)	111.9
8a	-	121.6	-	129.6
9	7.85(d, J=9.1)	119.3	7.49(d, J=8.8)	126.4
10	7.60(d, J=9.1)	127.3	7.53(d, J=8.8)	127.4
10a	-	122.7	-	124.3
3-OCH ₃	4.10(s)	61.3	4.11(s)	61.4
4-OCH ₃	3.96(s)	61.9	3.98(s)	59.8
8-OCH ₃	3.97(s)	59.7	-	-
2-OH	6.00(s)	-	6.03(s)	-
7-OH	5.80(s)	-	5.34	-

Nudol (**2**) - an amorphous pale yellow solid; UV spectrum (MeOH) (λ_{\max} nm (log ϵ): 211 (3.88), 258 (4.39), 283 (3.73), 347 (2.69) dan 363 nm (2.75); spektrum IR ν_{\max} (cm^{-1}): 3273 (OH), 2935 and 2850 (CH aromatic), 1618-1357 (C=C aromatic) and 1287 - 1159 (C-O-C ether). Spectrum $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) and $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz) - see Table 1.

The $^1\text{H-NMR}$ spectrum of Confusarin (**1**) reveals that the three aromatic protons, that are a pair of doublet signals ($J = 9.2$ Hz) at δ_{H} 9.19 and 7.29 ppm, a singlet signal at δ_{H} 7.19 ppm and a pair of doublet signals ($J = 9.1$ Hz) at δ_{H} 7.85 and 7.60 ppm could be assigned to penta-substituted phenanthrene. The spectrum indicates the presence of five oxyaryl protons through three singlet signals from methoxy groups at δ_{H} 4.10 ppm, 3.97 ppm and 3.96 ppm and two hydroxyl groups at δ_{H} 6.00 ppm and 5.80 ppm. This means that there are three methoxy and two hydroxyl groups attached to the aromatic ring.

The spectrum $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz) shows 17 perfectly separated carbon signals. They refer to nine quaternary carbon atoms (δ_{C} 145.4 ppm, 142.3 ppm, 140.9 ppm, 136.1 ppm, 135.1 ppm, 119.8 ppm, 114.4 ppm, 122.7 ppm and 121.6 ppm), five methine carbons (δ_{C} 127.3 ppm, 123.9 ppm, 119.3 ppm, 116.1 ppm and 108.1 ppm) and three methyl carbons from a methoxy group (δ_{C} 61.9 ppm, 61.3 ppm and 59.7 ppm). Five from the nine quaternary carbon atoms are of an oxyaryl carbons type (δ_{C} 145.4 ppm, 142.3 ppm, 140.9 ppm, 136.1 ppm and 135.1 ppm). The data presented shows that compound **1** belongs to the phenanthrene derivatives group.

Further information of the methoxy and hydroxyl group position in compound **1** phenanthrene is obtained from HMQC and HMBC analyses. The doublet signal ($J = 9.2$ Hz) at δ_{H} 9.19 ppm (H-5) is correlated with two quaternary carbon signals (δ_{C} 121.6 ppm and 114.4 ppm) and one oxyaryl carbon signal (δ_{C} 140.9 ppm). This data confirms the carbon signals (δ_{C} 140.9 ppm, 121.6 ppm and 114.4 ppm) at position C-4a, C-7 and C-8a. The correlation of the signal singlet at δ_{H} 7.19 with two quaternary carbon signals (δ_{C} 114.4 ppm and 122.7 ppm) and two oxyaryl carbon signals (δ_{C} 136.1 ppm and 142.3 ppm) attributes δ_{C} of 122,7 ppm to position C-10a, while δ_{C} of 114,4 ppm - to position C-4a. Meanwhile the two

oxyaryl carbons (δ_{C} 136.1 ppm and 142.3 ppm) occupy positions C-2 and C-3. The absence of a correlation between the signals at δ_{H} 7.19 ppm and δ_{C} 119.8 ppm (C-4b) indicates δ_{H} 7,19 ppm attribution to H-1 position. The UV, $^1\text{H-NMR}$, and $^{13}\text{C-NMR}$ data of compound **1** verify that it is 2,7-dihydroxy-3,4,8-trimethoxyphenantrene (Fig. 1). A compound of this structure named confusarin has been obtained from *Eria confuse* [14] but for the first time it is found in *Dioscorea esculenta*.

The IR spectrum of Nudol (**2**) shows a strong absorption band in the range of 1618 cm^{-1} -1357 cm^{-1} referring to aromatic rings, that of 1287 cm^{-1} -1159 cm^{-1} indicative of C-O-C ether and that at 3273 cm^{-1} attributed to hydroxyl groups. The absence of carbonyl groups is evident. The $^1\text{H-NMR}$ spectrum (CDCl_3 , 400 MHz) whose data is listed in Table 1 shows the presence of three signals of an aromatic type: ABX system signals at δ_{H} 9.36 ppm (d, $J = 9.1$ Hz), 7.19 ppm (dd, $J = 9.1$ and 2.8 Hz) and 7.16 ppm (d, $J = 2.8$ Hz), *ortho* doublet signals at δ_{H} 7.53 ppm and 7.49 ppm ($J = 8.8$ Hz) and a singlet signal of *penta* substituted benzene at δ_{H} 7,18 ppm. Two methoxy aromatic signals referred to two singlet protons at δ_{H} 4,11 ppm and 3,98 ppm are also outlined. In addition, there are two hydroxyl aromatic signals at δ_{H} 5,34 ppm and 6,03 ppm. Based on the molecular formula of compound $\text{C}_{16}\text{H}_{14}\text{O}_4$ and presence of four oxyaryl carbons (δ_{C} 153.2 ppm, 150.7 ppm, 147.5 ppm and 140.8 ppm), it can be suggested that the isolated compound is two hydroxy and two methoxy substituted phenanthrene at C-2, C-3, C-4 and C-7 positions, respectively.

The $^{13}\text{C-NMR}$ spectrum (CDCl_3 , 100 MHz) reported in Table 1 shows 16 perfectly separated carbon signals. The signals refer to eight quaternary carbons (δ_{C} 153.2 ppm, 150.7 ppm, 147.5 ppm, 140.9 ppm, 133.5 ppm, 129.6 ppm, 124.3 ppm and 118.9 ppm), four quaternary




Fig. 1. Chemical structure of confusarin (**1**).

oxyaryl carbons (δ_c 153,2 ppm; 150,7 ppm; 147,5 ppm and 140,8 ppm), six methin carbons (δ_c 128.5 ppm, 127.4 ppm, 126.4 ppm, 116.5 ppm, 111.9 ppm and 108.2 ppm), and two methyl carbons from a methoxy group (δ_c 61,4 ppm and 59,8 ppm). The data of ^{13}C -NMR analysis indicates that compound **2** belongs to the phenanthrene derivatives group.

The position of the methoxy and the hydroxyl group of compound **2** phenanthrene is determined on the ground of HMQC and HMBC analyses. The correlation between the singlet signal referring to an aromatic proton (H-1) at δ_H 7,18 ppm with two oxyaryl carbon signals (δ_c 147.5 ppm and 140.9 ppm) and one methin signal (δ_c 127.4 ppm) indicates that the position of the methin carbon δ_c 127.4 ppm is at C-10. The two oxyaryl carbons (δ_c 147.5 ppm and 140.9 ppm) occupy positions C-2 and C-3. The correlation between the hydroxyl singlet signal at δ_H 6.03 ppm, the two oxyaryl carbon signals (δ_c 147.5 ppm and 140.9 ppm) and the methin carbon signal (δ_c 108.2 ppm) confirms that OH position is at C-2, while that of the methin carbon (δ_c 108.2 ppm) is at C-1.

The position of $-\text{OCH}_3$ at C-3 is indicated by the correlation between the singlet methoxy group signal at δ_H 4.11 ppm and that of the oxyaryl carbon at δ_c 140.9 ppm. This in turn means that the position of the oxyaryl carbons (signals at δ_c 147.5 ppm and 140.9 ppm) is at C-2 and C-3. The correlation between the methoxy group signal (δ_H 3,98 ppm) and the oxyaryl carbon signal (δ_c 150,7 ppm) as well as the absence of a correlation between the ABX proton system signals at δ_H 9.36 ppm (d, $J=9.1$ Hz), 7.19 (dd, $J=9.1$ and 2.8 Hz) and 7.16 ppm (d, $J=2.8$ Hz) indicate that the position of the methoxy group (δ_c 150.7 ppm) is at C-4, while that of the hydroxyl group is at C-7. The UV, IR, ^1H -NMR, and ^{13}C -NMR data referring to compound **2** postulate that it is 2,7-dihydroxy-3,4-dimethoxyphenantrene (Fig. 2). A compound of this structure named nudol has been obtained from *Eulophia nuda* [15] but for the first time it is found in *Dioscorea esculenta*.

The antioxidant activity of compounds **1** and **2** against DPPH radical reagent is estimated to IC_{50} 19,63 \pm 0,09 and IC_{50} 37,91 \pm 0,08 ppm, correspondingly. The values pointed out provide to assume that the methoxy group at C-8 in compound **2** is responsible for the higher




Fig. 2. Chemical structure of nudol (**2**).

antioxidant activity displayed. Ascorbic acid of IC_{50} of 10,95 \pm 0.08 ppm is used as a reference in the antioxidant activity test.

CONCLUSIONS

Confusarin and nudol had been isolated from tuber of *dioscorea esculenta* L. The two isolated compounds showed an antioxidant activity of IC_{50} 19,63 \pm 0,09 and 37,91 \pm 0,08 ppm, respectively.

Acknowledgements

The research was supported by grants of Penelitian Unggulan Perguruan Tinggi (PUPT) from the Ministry of Research and Technology, Republic of Indonesia, 2016.

REFERENCES

1. Purnomo, Daryono, B. S., Rugayah, Sumardi, Issirep, Studi Etnobotani *Dioscorea* spp. (Dioscoreaceae) dan Kearifan Budaya Lokal Masyarakat di Sekitar Hutan Wonosadi Gunung Kidul Yogyakarta, J. Nat. Ind., 14, 2012, 191-198.
2. D. Gupta, J. Singh, p-Hydroxy acetophenone derivatives from *Dioscorea bulbifera*, Phyt., 28, 1989, 947-949.
3. K.H. Kim, M.A. Kim, E. Moon, S.Y. Kim, S.Z. Choi, M.W. Son, K.R. Lee, Furastonol saponins from the rhizomes of *Dioscorea japonica* and their effects of NGF induction, Bio. and Med. Chem., 21, 2011, 2075-2078.
4. M. Sautour, A.C. Mitaine-Offer, T. Miyamoto, H. Wagner, M.A. Lacaille-Dubois, A new phenanthrene glycoside and other constituents from *Dioscorea opposita*, Chem. Pharm. Bull., 52, 2004, 1235-1237.
5. R. Sunder, S. Rangaswami, G.C.S. Reddy, A new dihydrophenanthrene from *Dioscorea decipiens*, Phyt., 17, 1978, 1067.

6. K.W. Woo, E. Moon, O.W. Kwon, S.O. Lee, S.Y. Kim, S.Z. Choi, M.W. Son, K.R. Lee, Anti-neuroinflammatory diarylheptanoids from the rhizomes of *Dioscorea nipponica*, *Bio. and Med. Chem. Lett.*, 23, 2013, 3806-3809.
7. C. Chiu, J. Deng, H. Cheng, Y. Chen, M. Lee, W. Hou, C. Lee, S. Huang, G. Huang, Antioxidant and anti-inflammatory properties of taiwanese yam (*Dioscorea japonica* Thunb. var. *pseudojaponica* (Hayata) Yamam and its reference compounds, *Food Chem.*, 141, 2013, 1087-1096.
8. J. Cho, H. Choi, J. Lee, M. Kim, H. Sohn, D.L. Lee, The antifungal activity and membrane-disruptive action of dioscin extracted from *Dioscorea nipponica*, *Bioc. et Biop. Act.*, 1828, 2013, 1153-1158.
9. V. Kuete, R.B. Teponno, A.T. Mbaveng, L.A. Tapondjou, J.J.M. Meyer, L. Barboni, N. Lall, Antibacterial activities of the extracts, fractions and compounds from *Dioscorea bulbifera*, *BMC Comp. and Alt. Med.*, 12, 2012, 228.
10. D. Luo, Structural investigation of polysaccharide (DMB) purified from *Dioscorea Nipponica Makino*, *Carb. Poly.*, 103, 2014, 261-266.
11. S. Tewtrakul, A. dan Itharat, Anti-allergic substances from the rhizomes of *Dioscorea membranaceae*, *Bio. and Med. Chem.*, 14, 2006, 8707-8711.
12. M. Murugan, V.R. Mohan, In vitro antioxidant studies of *Dioscorea esculenta* (Lour). Burkill, *Asian Pas. J. of Trop. Biom.*, 2012, S1620-S1624.
13. P. Molyneux, The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity, *J. Sci. Tech.*, 26, 2004, 211-219.
14. P.L. Majumder, A. Kar, Confusarin and confusarinidin, two phenanthrene derivatives of the orchid *Eria confusa*, *Phyt.*, 26, 1987, 1127-1129.
15. S.R. Bhandari, A.H. Kapadi, P.L. Mujumder, M. Joardar, J.N. Shoolery, Nudol, a phenanthrene of the orchids *Eulophia nuda*, *Aria carinata* and *Eria stricta*, *Phyt.*, 24, 1985, 801-804.



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Journal of Chemical Technology and Metallurgy

Country [Bulgaria](#) -  [SJR Ranking of Bulgaria](#)

Subject Area and Category

[Chemical Engineering](#)
[Chemical Engineering \(miscellaneous\)](#)

[Engineering](#)
[Industrial and Manufacturing Engineering](#)

15

H Index

Publisher

[University of Chemical Technology and Metallurgy](#)

Publication type

Journals

ISSN

13143859, 13147471

Coverage

2013-ongoing




[How to publish in this journal](#)









[Join the conversation about this journal](#)

Quartiles



SJR 

Citations per document 



Journal of Chemical Technology and Metallurgy

Q2 Industrial and Manufacturing Engineering
best quartile

SJR 2018
0.26

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```

B **Bilal** 1 year ago

Dear Sir/M'am,

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Home](#)

Editorial Board

EDITOR-IN-CHIEF

Prof. Dr. Bogdana Koumanova

University of Chemical Technology and Metallurgy, Bulgaria

S.J. Allen,

Queens University of Belfast, UK

N.Yu. Bashkirceva,

National Research Technological University,
Kazan, Russia

M. Bojinov,

University of Chemical Technology and
Metallurgy, Bulgaria

V. Bojinov, University of Chemical Technology
and Metallurgy, Bulgaria

J. Carda,

University Jaume I, Castellon, Spain

D. Danalev,

University of Chemical Technology and
Metallurgy, Bulgaria

V. Dimitrov,

Bulgarian Academy of Sciences

N. Dishovsky,

University of Chemical Technology and
Metallurgy, Bulgaria

S.J.C. Foyo de Azevedo,

Universidade do Porto, Portugal

M. Jitaru,

I.P. Mazur,

Lipetsk State Technical University, Russia

D. Mehandjiev,

Bulgarian Academy of Sciences

E. Mihailov,

University of Chemical Technology and
Metallurgy, Bulgaria

L. Mörl,

University "Otto-von-Guericke", Magdeburg,
Germany

B. Nath,

European Centre for Pollution Research,
London, UK

A.B. Nayzabekov

Rudny Industrial Institute, Rudny,
Kazakhstan

L. Petrov,

Bulgarian Academy of Sciences

S. Piskin,

Yildiz Technical University, Istanbul, Turkey

A. K. Pogodaev,

University "Babeş –Bolyai", Cluj–Napoca,
Romania

S. Kalcheva,

University of Chemical Technology and
Metallurgy, Bulgaria

F. Keil,

Hamburg University of Technology, Germany

T. Koinov,

University of Chemical Technology and
Metallurgy, Bulgaria

T. Komatsu,

Nagaoka University of Technology, Japan

J.M. LeLann,

Institut National Polytechnique, École nationale
supérieure des ingénieurs en arts chimiques
et technologiques, France

S.N. Lezhnev,

Rudny Industrial Institute, Rudny, Kazakhstan

A. Mavrova,

University of Chemical Technology and
Metallurgy, Bulgaria

Lipetsk State Technical University, Russia

G. Radeva,

University of Chemical Technology and
Metallurgy, Bulgaria

A. Di Schino,

University of Perugia, Italy

M. Simeonova,

University of Chemical Technology and
Metallurgy, Bulgaria

V. Stefanova,

University of Chemical Technology and
Metallurgy, Bulgaria

D. Stoilova,

Bulgarian Academy of Sciences

N. Tsarevsky,

Southern Methodist University, Dallas,
Texas, USA

I. Turunen,

Lappeenranta University of Technology,
Finland

L. Vezenkov,

University of Chemical Technology and
Metallurgy, Bulgaria

[Log in](#) or [register](#) to post comments

uctm.edu/journal

[Home](#)

Volume 52, Iss. 6, 2017

Submitted by stanislav on Thu, 09/28/2017 - 20:40

Journal of Chemical Technology and Metallurgy

52, Iss. 6, 2017
ISSN 1314-7471 (print)
ISSN 1314-7978 (on line)

EDITOR-IN- CHIEF

Prof. Dr. Bogdana Koumanova

Tel: (+ 359 2) 81 63 302

University of Chemical Technology and Metallurgy

8 Kl. Ohridski, 1756 Sofia, Bulgaria

E-mail: journal@uctm.edu

SELECTED ARTICLES

International Conference

**“Collaboration Seminar of Chemistry and
Industry (CoSCI-2016)”**

5-6 October, 2016

Surabaya, Indonesia

Invited Editor

Dr. Purkan Purkan

[Primary study of cellulose acetate hollow fiber as a green membrane applied to hemodialysis](#)

Yanuardi Raharjo, Siti Wafiroh, Mahdya Nayla, Vita Yuliana,
Mochamad Zakki Fahmi

[Composite beads of chitosan/bentonite as a matrix for phosphate fertilizer controlled-release](#)

Bambang Piluharto, Veinardi Suendo, Ida Maulida, Asnawati

[Serum acetaldehyde as a potential biomarker for the detection of pathogenic biofilm formation by *Candida albicans*](#)

Masfufatun, Sumayyah Luqman Bayasud, Mei Shirli Yasinta, Ni'matuzahro, Afaf Baktir

[The influence of ascorbic acid, creatine, and creatinine on the uric acid analysis by potentiometry using a carbon paste modified imprinting zeolite electrode](#)

Miratul Khasanah, Muji Harsini, Alfa Akustia Widati, Prihantari Mukti Ibrani

[A novel spectrophotometric method for determination of histamine based on its complex reaction with Cu\(II\) and alizarin red S](#)

Miftakhul Jannatin, Ayu Nabila I.L, Ganden Supriyanto, Pratiwi Pudjiastuti

[Application of ionic liquid dispersive liquid-liquid microextraction for analysis of n-nitrosodipropylamine in salted fish](#)

Aning Purwaningsih, Yanuardi Raharjo, Hendarta Agasi

[Determination of chlorpyrifos pesticide by effervescence liquid phase microextraction HPLC UV-Vis](#)

Usreg Sri Handajani, Yanuardi Raharjo, Bagas Wantoro

[Effect of aliphatic and aromatic hydrocarbons on the oxygenase production from hydrocarbonoclastic bacteria](#)

Sri Sumarsih, Ni'matuzahroh, Fatimah, Miranti Puspitasari, Meilisa Rusdiana

[Identification of *Candida* species by assimilation and Multiplex-PCR methods](#)

Hermansyah, Nurmalina Adhiyanti, Julinar, Kemas Yakub
Rahadiyanto, Susilawati

[Xylanase enzyme from a local strain of *Pseudomonas stutzeri*](#)

Purkan Purkan, Emma Huruniawati, Sri Sumarsih

[Study of a catalyst of citric acid crosslinking on locust bean gum](#)

Wuryanto Hadinugroho, Suwaldi Martodihardjo, Achmad Fudholi,
Sugeng Riyanto

[Production and characterization of sulfonated chitosan-calcium oxide composite membrane as a proton exchange fuel cell membrane](#)

Siti Wafiroh, Abdulloh, Winda Kusuma Wardani

[An excellent way to prepare conductive glass using a simple glass plate aiming a promising solar cell](#)

Harsasi Setyawati, Handoko Darmokoesoemo, Hamami, Faidur
Rochman, Ahmadi Jaya Permana

[Partial oxidative synthesis of fluorescent carbon derived from local bamboo leaves](#)

Ahmadi Jaya Permana, Abdul Haris, Harsasi Setyawati, Mochamad
Zakki Fahmi

[Stability of coordination compounds obtained by reduction of copper\(II\) halide and 1,3-bis\(diphenylphosphino\)propane \(DPPP\)](#)

Nike Prilil Puspita Sari, Lis Siaturohmah, Effendy, Fariati

[Phenolic compounds from *Aquilaria microcarpa* stem bark](#)

Alfinda Novi Kristanti, Mulyadi Tanjung, Okky P. Rahayu, Erika Herdiana

[Electrochemical degradation of naphthol AS-BO batik dyes](#)

Muji Harsini, Suyanto, Yhosep Gita Y. Y., Lilik Rhodifasari, Handoko Darmokoesomo

[Silica-methyltrimethoxysilane based hydrophobic coatings on a glass substrate](#)

Alfa A. Widati, Nuryono Nuryono, Indriana Kartini, Noah D. Martino

[Chitosan-based neem seed extract nanocapsules: a new approach on enhancing its effectiveness as an insecticide delivery agent](#)

Mochamad Zakki Fahmi, Hery Suwito, Achmadi Susilo, Elika Joeniarti, Anninda Mughniy Rahayu Jaswadi, Nindayu Indrasari

[Confusarin and nudol, two phenanthrene group compounds, from *Dioscorea esculenta* L. and their antioxidant activities](#)

Nanik Siti Aminah, Ratih Hidayah, Mulyadi Tanjung

[Drug delivery hard shell capsules from seaweed extracts](#)

Pratiwi Pudjiastuti, Muhammad Al Rizqi Dharma Fauzi, Handoko Darmokoesoemo

[Organic template free hierarchical ZSM-5 prepared by desilication](#)

Hartati, Alfa Akustia Widati, Aning Purwaningsih, Alfinda Novi Kristanti, Anggarani Nur Oktavia

[Modification of gresik's dolomite to CaO•MgO nanocomposite as a catalyst for synthesis of biodiesel from tamanu oil](#)

Abdulloh Abdulloh, Alfa Akustia Widati, Oditio Arizal

[Guide for Authors](#)