



Search by any title, abstract, authors...



Research.Publish.Connect.



Proceedings

Proceedings of the 1st International Conference on Chemical Science and Technology Innovation

July 14-15, 2019, in Medan, Indonesia



Editors: Basuki Wirjosentono ¹; Seri Bima Sembiring ¹; Zul Alfian ¹; Rizalman Mamat ²; Irwana Nainggolan ³; Saharman Gea ⁴ and Markus Fischer ⁵

Affiliations: ¹ Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sumatera Utara, Indonesia ; ² Department of Mechanical Engineering, Universiti Malaysia Pahang, Pahang, Malaysia, Malaysia ; ³ Universitas Sumatera Utara, Indonesia ; ⁴ Department of Chemistry, Faculty of Mathematics and Natural Science, Universitas Sumatera Utara, Jalan Biotechnologi No.1, Medan 20155, Sumatera Utara, Indonesia, Indonesia ; ⁵ Institut für Lebensmittelchemie, Fachbereich Chemie, Universität Hamburg, Germany

ISBN: 978-989-758-415-2

Conference Link: <https://ocs.usu.ac.id/ICOCTI/ICOCTI2019>

Foreword: On behalf of the organizing committee of the International Conference and Department of Chemistry, Universitas Sumatera Utara, we would like to express our warmest welcome for all our participants and speakers to the 1st International

Conference on Chemical Science and Technology Innovation (ICOCTI) 2019. The conference itself was held in Medan, July 18-19th 2019, with the theme " A Chemical Breakthrough for Science Technology Innovation". With over 60 selected manuscripts presented in the conference from authors across the globe, this proceeding aimed to serve as a platform to exchange information of the research methods, results, and experiences from various field of chemical science and engineering all over the world, including organic chemistry, inorganic chemistry, natural product chemistry, material science, polymer chemistry, engineering chemistry, and environmental sciences. In addition to that, we hope that this proceeding could promote research opportunities for the **(More)**

Volumes:

Vol. 1 - 978-989-758-415-2

Papers	Authors	
Show All  papers		
Synthesis, Antioxidant and Toxicity Activity of Compounds (E)-1-(3-bromophenyl)-3-p tolylprop-2-en-1-on	Eti Meirina Brahmana , Jamaran Kaban , Ginda Haro , Juliati Br Tarigan , Basuki Wirjosentono , Tamrin and Mimpin Ginting	P. 5 - 9 DOI: 10.5220/00088364
Examination of Micro-structural, Mechanical Properties and Investigation of Optimum Conditions of Diffusion Bonding between Grey Cast Iron and Low Carbon Steel	Fauzuddin Ayob , Patthi Hussain , Mokhtar Awang , Aiziz Abdullah , Bakhtiar Arif Baharuddin , Aziz Abdul Rahim and Asmawi Ismail	P. 10 - 17 DOI: 10.5220/00088370
Mangan Coated N-Graphene for Good Performance Electrode in Primary Battery Anode	Rikson Siburian , Sabarmin Perangin-angin , Helmina Sembiring , Crystina Simanjuntak and Yopi Sihombing	P. 18 - 20 DOI: 10.5220/00088383
Furfural Synthesis from Mile-a-Minute Weed (<i>Mikania micrantha</i>) using Roselle Petal Extract as Catalyst	Taslim , Iriany , Okta Bani and Apri Wardiana Sinaga	P. 21 - 25 DOI: 10.5220/00088384
Nanometers Formation Model of Iron (Fe) and Magnesium (Mg) on Graphene Nano Sheets	Rikson Siburian , Herlince Sihotang , Saur Lumban Raja , Minto Supeno , Crystina Simanjuntak and Hana Manurung	P. 26 - 29 DOI: 10.5220/00088386
Analysis of Total Phenolics and Flavonoid from the Root Bark of Flacourtie rukam	Muharni , Elfita , Julinar and Heni Yohandini	P. 30 - 35 DOI: 10.5220/00088392
Formation Process of Graphene Nano Sheets		P. 36 - 38

The Synthesis of Graphene from Coconut Shell Charcoal

P. 39 -
44

Minto Supeno , Rikson Siburian and Desi Natalia

DOI:10.5220/00088396

Study of Environmental Approaching on Lake Toba

P. 45 -
51

Parlindungan Purba , Sirozuliam , Agus Purwoko and Hidayati

DOI:10.5220/00088487

A Tensile Test of Bio-Board Made from Bamboo - Pineapple Fiber - Coconut Fiber as Nonvolatile Material

P. 52 -
56

Suryani Salim , Teuku Rihayat , Shafira Riskina , Nurhanifa Aidy , Mawaddah Razali , Ana Siti Nurmasyitah , Murni , Fitria and Yuhanis Yunus

DOI:10.5220/00088528

Distillation of Agarwood Oil (*Aquillaria sp*) using Photovoltaic Methods

P. 57 -
62

Nelly Safitri , Teuku Rihayat , Suryani , Shafira Riskina , Rudi Saputra , Aida Safitri , Tjut Yayang Risqatia Hasnah , Kiki Yolanda Putri and Yuhanis Yunus

DOI:10.5220/00088533

Synthesis of Carboxymethyl Polysaccharide from Arenga Pinnata Polysaccharide and Monochloroasetic

P. 63 -
68

Juliaty Br. Tarigan , Diana A. Barus and Nico Hot Asi Naibaho

DOI:10.5220/00088541

Synthesis Succinic Galactomannan from Galactomannan Arenga pinnata Merr. and Succinic Anhydride using Microwave Method

P. 69 -
73

Juliaty Br. Tarigan and Mutiara H. Siregar

DOI:10.5220/00088542

Sustainable Supply Chain Management in Manufacture

P. 74 -
78

Nurhayati Sembiring , Mangara M. Tambunan , Elisabeth Ginting and Intan Sari Dewi

DOI:10.5220/00088543

Pillarization of Bentonite using Fe3+/Ti4+ and Its Application for Congo Red and Direct Violet Removal

P. 79 -
88

Muhammad Said , Riza Antini , Tarmizi Taher and Aldes Lesbani

DOI:10.5220/00088544

Bioactivity and Phytochemical Constituents of Extract Etanol from *Musa paradisiaca* Linn

P. 89 -
95

Mayang Sari , Tamrin , Jamaran Kaban and Zul Alfian

DOI:10.5220/00088551

***Illicium verum* Essential Oil as Antibacterial Agent**

P. 96 -
99

Rizki Damayanti , Tamrin , Eddyanto and Zul Alfian

DOI:10.5220/00088552

Phosphate Adsorption using KOH Activated Coal Bottom Ash

P. 100 -
104

Fatimah Agussalim , Alfikri Ramadhan and Budi Pratama Tarigan

DOI:10.5220/00088555

**Kinetics of Petroleum Oil Recovery in Bio Surfactant of
Brevundimonas diminuta and Pseudomonas citronellolis Bacteria
Solutions** P. 105 -
112

Bambang Yudono , Sri Pertiwi Estuningsih , Muhammad Tri
Handoko and Muhammad Said

DOI:10.5220/00088564

Phase Formation of M-Type BaFe₁₂O₁₉/ZnO Magnetic Material P. 113 -
115

Syahrul Humaidi , Tua Raja Simbolon , Diana A Barus ,
Veryyon Hrp , Perdamean S , Chandra K , Eko Arif S and
Silviana S **DOI:**10.5220/00088567

**Synthesis and Characterization of Superabsorbent Polymer based
on Carboxymethyl Cellulose, Breadfruit Starch and Aluminum
Sulfate** P. 116 -
120

Darwin Yunus Nasution , Marpongahtun , Dede
Ibrahim Muthawali , A. D. Budiman and Zulfikar **DOI:**10.5220/00088571

Cocoa Pod (*Theobroma cacao L.*) Utilization as Material for CMC P. 121 -
123

Yeni Afriani , Zanira Urfa Harahap , Era Fazira Matondang and Cut
Fatimah Zuhra **DOI:**10.5220/00088572

**Production of Bioethanol Gel from Sugar Cane Waste with
Carbopol as Alternative Fuel** P. 124 -
129

Wilza Fithri Az-zahra , N. Nurlina Harahap , S. Haidar Putra and
M. Zulham Efendi Sinaga **DOI:**10.5220/00088575

**Isolation and Identification of Secondary Metabolite Compound
Extract Etil Acetate from the Leaves of Durian *Durio zibethinus L.*** P. 130 -
132

Dyna Grace Romatua Aruan , Tonel Barus , Ginda Haro and
Partomuan Simanjuntak **DOI:**10.5220/00088578

**Lead (II) Nitrate Catalyzed Esterification Palmitic Acid with
Alcohols** P. 133 -
135

Nimpan Bangun , Justaman Karo-Karo , Hamongan Nainggolan
, Slamet Silaban , Nia Erisa Tarigan and Rahmad Ramadan **DOI:**10.5220/00088579

**Chemical Compounds from Fungus *Syncephalastrum racemosum*
Isolated as Endophytic from *Ageratum conyzoides*** P. 136 -
140

Elfita , Muharni , Mardiyanto , Fitrya and
Rismawati Simangunsong **DOI:**10.5220/00088580

**Cytotoxic Carbazole Alkaloid from the Root of *Clausena excavata*
on Hela Cell Line** P. 141 -
144

Tin Myo Thant , Nanik Siti Aminah , Alfinda Novi Kristanti ,
Rico Ramadhan , Hnin Thanda Aung and Yoshiaki Takaya **DOI:**10.5220/00088581

Modification and Characterization Natural Cycle Rubber (Resipren-35) with Oleat Acid using Dicumyl Peroxide and Divinilbenzena as Compatibility P. 145 - 151

Barita Aritonang , Tamrin , Basuki Wirjosentono and Eddiyanto

DOI:10.5220/00088597

Preparation and Characterization of Biochar from Palm Kernel Shells as an Activated Carbon Precursors with the Pyrolysis Method P. 152 - 155

Ratni Dewi , Harry Agusnar , Zul Alfian and Tamrin

DOI:10.5220/00088631

Cytotoxicity of Heterophyllene A, the Derivative of Arylbenzofuran from Stem Bark of *Artocarpus calophylla* P. 156 - 159

Rumzil Laili , Nanik Siti Aminah , Tin Myo Thant , Yoshiaki Takaya , Ainun Sa'adah and Alfinda Novi Kristanti

DOI:10.5220/00088632

Synthesis and Identification of Furfural from Cocoa Pod Husk (CPH) with Pretreatment Process before Hydrolysis Process P. 160 - 164

Lisa Aulia Lubis , Amir Husin and Maulida

DOI:10.5220/00088638

Making Composites from Mixing Limestone with Addition of Latex P. 165 - 173

Amir Hamzah Siregar , Saharman Gea and Nora Indriani

DOI:10.5220/00088640

Characterization of α -Cellulose from Bagasse Cane Bz 132 (Saccharum officinarum) P. 174 - 177

Tengku Rachmi Hidayani , Basuki Wirjosentono , Darwin Yunus Nasution and Tamrin

DOI:10.5220/00088646

The Effectiveness of Chitosan as an Antimicrobial on Bacterial Cellulose-based Scaffold Skin Tissue Engineering P. 178 - 181

Khatarina Meldawati Pasaribu , Saharman Gea , Safruddin Ilyas and Tamrin

DOI:10.5220/00088647

Effect of Comparative Flour of Purple Sweet Potato (*Ipomea batatas*) and Wheat Flour of Nutrition Value on Brownies P. 182 - 186

Emma Zaidar , Rumondang Bulan , Firman Sebayang , M. Zulham Efendi Sinaga and Rina Latipah

DOI:10.5220/00088657

Identifying Trends of Sustainable Supply Chain Management P. 187 - 190

Nurhayati Sembiring , Mangara M Tambunan and Elisabeth Ginting

DOI:10.5220/00088681

Modification and Characterization Starch Nanoparticles of Mangrove Fruit using Chemical-mechanical Method and Application as Basic Materials Making Hydrogel P. 191 - 196

Gimelliya Saragih , Tamrin , Marpongahtun and Darwin Yunus Nasution

DOI:10.5220/00088685

Antioxidants Activity of the Kecombrang Flower (*Etlingera elatior*) Extract by using 1,1-diphenyl-2-picrilhidrazyl (DPPH) Method P. 197 - 203

- Making Hydrogel with Crosslinked Reactions between Chitosan and Dialdehyde Cellulose from Coconut Fiber as Wound Healers** P. 204 -
Firman Sebayang , Rumondang Bulan , Emma Zaidar Nasution , 210
M. Zulham Efendi Sinaga and Windi Anggara Putri DOI:10.5220/00088694
- Effect Croslinking on Characteristics of Silica Chitosan Composite from Vulcanic Ash of Sinabung Mount and Shrimp Husk by Sol Gel Method** P. 211 -
Lisnawaty Simatupang , Manihar Situmorang , Rikson Siburian 214
and Basuki Wirjosentono DOI:10.5220/00088759
- Potential Antibacterial and Antioxidant Activiy of Methanolic Extract of Vigna unguiculata (L.) Walp Leaves** P. 215 -
Sovia Lenny and Dinda Widya Rizky 217
DOI:10.5220/00088786
- Synthesis of Cross-linked Breadfruit Starch (*Artocarpus communis*) using Epichlorohydrin** P. 218 -
Cut Fatimah Zuhra , Mimpin Ginting and Desi Sonya Siregar 222
DOI:10.5220/00088789
- Synthesis of Quatenary Ammonium Compounds from Eugenol through Mannich and Methylation Reactions and Its Antibacterial Activity** P. 223 -
Sabarmin Perangin-angin and Sri Wahyuni Barus 228
DOI:10.5220/00088792
- Synthesis 2-(3-Phenylallylidenamino) Pentanedioic Acid by Condensation of Cinnamaldehyde with Glutamic Acid and the Activity Test as Antibact** P. 229 -
Mimpin Ginting , Deni Aprilina Lumban Gaul , Indra Masmur and 233
Andy Candra DOI:10.5220/00089197
- Isolation and Characterization of an Antioxidant Compound from Kayu Hitam Leaves (*Diospyros celebica* Bakh.F.)** P. 234 -
Helmina Br. Sembiring and Yuni Romasni Purba 238
DOI:10.5220/00089198
- Development of Maceration Methods in the Investigation and Analysis of Lard in Industrial Food Products to Improve Halal Product in Indonesia** P. 239 -
Muhammad Taufik , Zul Alfian , Desi Ardilla , Mariany Razali , 244
Irfan Kurniawan , Nur Waridah Angriani Nasution , Rika Astuti
Pulungan and Sri Dewi Sitohang DOI:10.5220/00089199
- Identification and Analysis of Potential Antioxidants from Leaves of *Eucalyptus robusta* PT. Toba Pulp Lestari, Tbk.** P. 245 -
Chintya Cahaya , Muhammad Taufik , Zul Alfian , Sovia Lenny and 248
Rizky Hidayati DOI:10.5220/00089200
- The AAS Method for Arsen Analysis in Cabbage in the Area of Sinabung Post Eruption** P. 249 -
252

Identification of Potential Antioxidants from Leaves of Eucalyptus grandis PT Toba Pulp Lestari, Tbk. P. 253 - 256

Rizky Hidayati , Muhammad Taufik , Zul Alfian , Sopia Lenny ,
Chintya Cahaya , Simon Sidabuke and E. Manullang
DOI:10.5220/00089202

Analysis of Physical Properties and Chemical Contents in Eucalyptus citriodora PT. Toba Pulp Lestari, Tbk P. 257 - 261

Zul Alfian , Muhammad Taufik , Sopia Lenny , Chintya Cahaya ,
Rizky Hidayati , Simon Sidabuke , E. Manullang and
Jeremi Sitohang
DOI:10.5220/00089203

Quality Improvement of Liquid Smoke of Coconut Shell by Tar Scrubber P. 262 - 265

Reka M. Sari , Saharman Gea , Basuki Wirjosentono and
Sunit Hendrana
DOI:10.5220/00089208

Thermal Properties of Chitosan-roselle Films P. 266 - 273

Irwana Nainggolan , Tulus Ikhsan Nasution and Khairel Rafezi
Ahmad
DOI:10.5220/00089214

Magnesium Impregnated Silica Mesoporous Prepared using Ester Ricinoleic as Template for the Esterification P. 274 - 279

Andriayani , Marpongahtun and Yugia Muis
DOI:10.5220/00089217

Preparation and Characterization of Chitosan with Activated Carbon as Adsorbent to Reduce Level Metal Cadmium (Cd) and Nickel (Ni) P. 280 - 286

Fitri Purnama Sari , Harry Agusnar and Muhammad Taufik
DOI:10.5220/00089218

Mechanical Properties and Morphology Biocomposites of Polycaprolactone (PCL)/Modified using Trisodium Trimetaphosphate P. 287 - 292

Marpongahtun , Cut Fatimah Zuhra , Sopia Lenny , Darwin Yunus
Nasution , Aulya M. , Suci A. Amaturrahim , Mahmud ,
Fitria Puspa Dewi and Irmayani
DOI:10.5220/00089258

Effect of Particle Size from Oil Palm Wood Powder as Pore Forming Agent on Porosity and Hardness of Porous Ceramics based on Clay and Polyvinyl Alcohol P. 293 - 296

Marpongahtun , Darwin Yunus Nasution and Taufik Hidayat
DOI:10.5220/00089259

In Vitro Investigation of Bacterial Cellulose/Turmeric Extract (BC-TE) Nanocomposite for Burn Wound Dressing P. 297 - 300

Rio Cahyono Siahaan , Melati Sitorus , Rayhan Mulia and
Saharman Gea
DOI:10.5220/00089279

Preparation and Characteristics of Polyvinyl Alcohol-based Hydrogel Containing Natural Microbentonite

P. 301 -

304

Mawarani Manullang , Diana Adnanda Nasution and Basuki Wirjosentono

DOI:10.5220/00089281

Modification of Pulp Cellulose of Belangke Bamboo (*Gigantochloa pruriens*) using [2-(Acryloyloxy)Ethyl] Trimethyl Ammonium Chloride and Maleic Anhydride

P. 305 -

311

Rina Ridara , Diana Adnanda Nasution and Basuki Wirjosentono

DOI:10.5220/00089323

Thermal and Morphological Properties of Polyvinyl Alcohol-based Hydrogel Containing Microcrystal Cellulose

P. 312 -

318

Riski Aulia Hasibuan , Diana Adnanda Nasution and Basuki Wirjosentono

DOI:10.5220/00089324

Antimicrobial Properties of [2-(Acryloyloxy)Ethyl]-Trimethyl Ammonium Chloride and Maleic Anhydride Surface Grafted-Cotton Fibers

P. 319 -

323

Hazna Sartiva , Diana Adnanda Nasution and Basuki Wirjosentono

DOI:10.5220/00089341

X-Ray Diffraction and Morphology Studies of Sulfonated Polystyrene and Maleated Natural Rubber Blend with PE-g-MA as Compatibilized

P. 324 -

328

Ahmad Nasir Pulungan , Basuki Wirjosentono , Eddiyanto and Sunit Hendrana

DOI:10.5220/00089350

Optimization of Polymer Networks using Variations in Moleculer Weight of Polyetilen Glycol in the Manufacture of Semi – IPN Hidrogels from Coconut Water Cellulose Bacteria

P. 329 -

334

Tamrin and Emma Zaidar

DOI:10.5220/00090032

Addition of Taro, Breadfruit, Seaweeds and Soymilk Waste Powders into Cassava Starch and Corn Powder-based Rice Analogues

P. 335 -

338

Basuki Wirjosentono , Tamrin , Amir Hamzah Siregar , Diana Adnanda Nasution , Isna Shara Siregar , Siti Sarah Fazira , Hanifah Siregar and Nurainun Mardhiah

DOI:10.5220/00090051

Proximate and Physical Characteristics of Cookies Made of Cassava Starch and Wheat Flour Blends Containing Soymilk Waste Powder

P. 339 -

342

Basuki Wirjosentono , Tamrin , Amir Hamzah Siregar , Diana Adnanda Nasution and Paula Netti Sihombing

DOI:10.5220/00090052

RESOURCES

Proceedings

Papers

Authors

Ontology

CONTACTS

Science and Technology
Publications, Lda
Avenida de S. Francisco
Xavier, Lote 7 Cv. C,
2900-616 Setúbal,
Portugal.

Phone: +351 265 520 185
Fax: +351 265 520 186
Email: info@scitepress.org

EXTERNAL LINKS

PRIMORIS
INSTICC
SCITEVENTS
CROSSREF

PROCEEDINGS SUBMITTED FOR INDEXATION BY:

Thomson Reuters
dblp
Ei Compendex
SCOPUS
Semantic Scholar

© 2020 SciTePress, Science and Technology Publications, Lda - All rights reserved.

Cytotoxic Carbazole Alkaloid from the Root of *Clausena excavata* on HeLa Cell Line

Tin Myo Thant^{1,2}, Nanik Siti Aminah^{3*}, Alfinda Novi Kristanti³, Rico Ramadhan³, Hnin Thanda Aung⁴ and Yoshiaki Takaya⁵

¹Ph.D. Student of Mathematics and Natural Sciences, Faculty of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

²Dept. of Chemistry, Mandalay Degree College, Mandalay, Myanmar

³Dept. of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Komplek Kampus C UNAIR, Jl. Mulyorejo, Surabaya, Indonesia

⁴Dept. of Chemistry, Mandalay University, Mandalay, Myanmar

⁵Fac. Of Pharmacy, Meijo University, 150 Yagotoyama, Tempaku, Nagoya, 468-8503 Japan

Keywords: *Clausena excavata*, Carbazole Alkaloid, 7-hydroxyheptaphylline, MTT assay, HeLa.

Abstract: In a search for bioactive constituents from Myanmar medicinal plants, a carbazole alkaloid, named 7-hydroxy heptaphylline (1) was isolated from the root of *Clausena excavata*. The structure of isolated compound was elucidated based on spectrophotometric data such as UV-vis, FT-IR, NMR and HRMS data. The cytotoxicity of the isolated compound (1) was evaluated by MTT assay against on HeLa cancer cells. The compound (1) exhibited moderate inhibition activity with IC₅₀ 41.4 µg/ml.

1 INTRODUCTION

Clausena excavata Burm. f. is a wild shrub, a member of Rutaceae family predominantly distributed in India, China and Southeast Asia. The leaves, twigs, and root barks of *C. excavata* have long been used in Asian folk medicine for the treatment of colic, cough, rhinitis, sore, wounds, malaria, abdominal pain, snake-bite, preliminary stage of AIDS and dermatopathy, dysentery, enteritis, and urethra infection (Waziri et al., 2016a) (Kumar et al., 2012) (Peh et al., 2013).

The constituents of *C. excavata* have been frequently studied. Phytochemical analyses in the past have revealed that *C. excavata* is a rich source of coumarins, carbazole alkaloid along with a small group of flavonoids, limonoids and triterpenoids (Cheng et al., 2009) (Mohan, 2012) (Sunthitikawinsakul et al., 2003) (Kumar et al., 2012) (Peh et al., 2013) (Peng et al., 2013) (Thant et al., 2019). Many compounds reported from *C. excavata* showed diverse therapeutic activities which are antibacterial, antifungal, antiplatelet, antiplasmodial, antitumor, antinociceptive,

antimycobacterial, and anti-HIV-1 activities (Kongkathip and Kongkathip, 2009).

The coumarins isolated from this plant have attracted attention due to its bioactive properties such as the furanone-coumarins named clauslactones A–J isolated from leaves exhibited tumor promotion inhibitory effects, nordentatin showed antibacterial properties and a pyranocoumarin clausenidin isolated from roots displayed anti-HIV-1 activity (Kumar et al., 2012). Nevertheless, the other potent bioactivities of the constituents from *C. excavata* are still unknown and worthy of exploration (Cheng et al., 2009). Moreover, four isolated pyranocoumarins from *C. excavata* and screened their cytotoxic potentials in cancer cells. The study revealed that the pyranocoumarins are good modulators of tumor cell death (Waziri et al., 2016b) (N. W. Muhsin Sharif, 2011).

Cancer is the second leading cause of death worldwide. Cervical cancer is one of the most dead list diseases among women and it is occurred when the abnormal cells are undergoing to the rapid and uncontrolled growth on the cervix. Current treatments for cervical cancer may include surgery, drugs (hormonal therapy and chemotherapy),

radiation and/or immunotherapy. Conventional cancer treatments such as chemotherapy and radiotherapy have shown some effectiveness for reducing or eradicating cancers; however, they can produce unpleasant side effects, e.g. nausea, vomiting, changes in bowel habits, fatigue and hair loss. Complementary and alternative medicine (CAM), herbals and multivitamin supplements, or herbal medicine is increasingly used as an adjunctive treatment for cancer patients to reduce or manage side effects of conventional cancer treatments. Several studies have confirmed the anti-proliferative and cell cycle regulatory effects of some plants which behave as cancer prevention (Amin et al., 2009) (Azarifar et al., 2015). More than 25% of drugs used during the last 20 years are directly derived from plants, while the other 25% are chemically altered natural products. Still, only 5-15% of the approximately 250,000 higher plants have ever been investigated for bioactive compounds. The advantage of using such compounds for cancer treatment is their relatively non-toxic nature and availability in an ingestive form (Amin et al., 2009).

In continuing our research, isolation of bioactive constituents from natural product, isolated compound (**1**) was tested on his cytotoxicity activity on HeLa cancer cell by MTT assay.

2 MATERIALS AND METHODS

2.1 Plant Material

The plant sample of *C. excavata* was collected from Pyin Ma Nar Township, Mandalay Division, Myanmar in October 2016. It was identified by Prof. Soe Myint Aye, botanist from Department of Botany, Mandalay University, Myanmar.

2.2 MTT Assay

Anticancer activity tests on isolated compound was carried out using the MTT assay method (3-[4,5-dimethylthiazol-2-il]-2,5 diphenyl tetrazolium bromide) following the protocol of Suwito et al (Suwito et al., 2018). The cancer cells were seeded in a 96-well plate at a density of 1×10^4 cells/well with a phenol red-free RPMI (Roswell Park Memorial Institute medium) 1640 medium (containing 10% FBS (fetal bovine serum)) and maintained for 24 h. Subsequently, the tested compound (various concentrations) was applied for

24 h. After addition of 0.5% MTT solution, the incubation was continued for a further 4 h at 37 °C/5% CO₂. The stop solution (0.04 N HCl in isopropanol) was added to the culture medium to each well. Then, the spectroscopic measurement was carried out at 570 nm (peak) and 630 nm (bottom) using an ELISA (Enzyme-Linked Immunosorbent Assay) reader. It was conducted in triplicate. Doxorubicin was used as a positive control. The value of IC₅₀ was determined using a probit analysis (SPSS 17, IBM Analytics, New York, NY, USA).

2.3 Extraction and Isolation

The dried roots (3.6 kg) were extracted successively with 95% EtOH (12.0 L) over a period of two weeks at ambient temperature. After removing the solvent 156 g of ethanolic extract was obtained. Among them the extract (100g) was partitioned by using solvents; *n*-hexane: methanol (1:1, v/v). Then methanol portion (80.4 g) was subjected to vacuum liquid chromatography over silica gel (150g) eluted with different mixtures of *n*-hexane: ethylacetate by stepwise increasing gradient polarity gave a total of 7 combined fractions (MF-1 to -7) were obtained. MF-6.2.1.1 was purified by using *n*-hexane: ethylacetate, ethylacetate (0-30%) with gradient polarity and silica gel column, gave 44 fractions. Among them 7-21 showed one spot on tlc with three different solvent elution system (*n*-hex: EtOAc; *n*-hexane: CHCl₃; *n*-hexane : Acetone) afforded as pure compound (**1**).

2.4 Statistical Analysis

Results were presented as mean \pm SD in triplicate experiment. Differences were determined using SPSS 17, IBM Analytics, New York, NY, USA at significant difference of 0.05.

3 RESULTS AND DISCUSSIONS

Spectroscopic Data of Compound-1 (7-hydroxyheptaphylline)

Compound-1: greenish yellow solid. UV(MeOH), λ_{max} (log_ε) 342 (0.629), 303 (2.519), 237 (1.408) nm. IR (KBr) ν_{max} cm⁻¹: 3250, 2958, 2922, 2852, 1743, 1614, 1454, 1327, 1186, 9966 ¹H NMR (600 MHz, DMSO-d6) δ 11.53, 11.27, 9.88, 9.49, 8.14, 7.80, 7.78, 6.86, 6.86, 6.66, 6.65, 6.64, 6.64, 5.29, 5.28, 5.28, 5.27, 5.27, 5.27, 5.26, 5.26, 5.26, 3.51, 3.50, 1.78, 1.63. ¹³C NMR (151 MHz, DMSO-d6) δ

196.45, 156.98, 156.59, 145.16, 143.00, 132.15, 124.25, 122.11, 120.96, 117.83, 115.91, 115.01, 109.84, 109.32, 97.80, 25.92, 23.08, 18.39.




Figure 1: Structure of compound-1 isolated from the root of *C. excavata*.

Table 1: Cytotoxic activity of isolated compound (**1**) against on *HeLa* cell lines.

Compounds	IC ₅₀ (ug/mL)
7-hydroxy heptaphylline (1)	41.4
Doxorubicin	1.19

Table 2: ¹H (600 MHz), ¹³C (151 MHz) NMR and HMBC spectral data of compound-1 (7-hydroxyheptaphylline).

Position	δ_{H} (mult, J Hz)	δ_{C} (ppm)	HMBC
1a	-	145.2	
1	-	109.1	
2	-	157.9	
3	-	123.7	
4	8.03 (s, 1H)	125.3	C-1a,C-2,C-3,C-4a, CHO
4a	-	115.5	
5a	-	140.2	
5	7.97 (dd, J = 7.7 Hz, 1H)	119.8	C-5a,C-8,C-8a
6	7.27 (m, 1H)	120.5	C-5a,C-7,C-8
7	7.40 (dt, J = 7.7, 3.8 Hz, 1H)	125.9	C-6
8	7.41 (dd, J = 7.7,3.8 Hz, 1H)	110.9	C-5a
8a	-	117.4	
1'	3.64 (d, J = 6.9 Hz, 2H,H-1')	22.9	C-1,C-1a,C-2,C-10, C-11
2'	5.33 (d, J = 6.9 Hz, 1H,H-2')	121.3	C-1,C-9,C-12,C-13
3'	-	134.2	
4'	1.91 (s, 3H)	18.1	C-10,C-11,C-13
5'	1.78 (s, 3H)	25.7	C-10, C-11, C-12
-CHO	9.91	195.4	C-1,C-2,C-4
-NH	8.21		
-OH	11.66		

Compound-**1** (7-hydroxyheptaphylline) was afforded as yellow green crystal UV (MeOH), λ_{max} (log ϵ) 342 (0.629), 303(2.519), 237(1.408), 249(1.64) revealed as basic carbazole alkaloid. The IR spectrum of compound-1 showed the presence of -OH and -NH

functional groups at (3250cm⁻¹) and also carbonyl and aromatic benzene groups at (1743, 1612 and 1588 cm⁻¹). In the ¹H NMR spectrum the compound-**1** revealed a total of 11 signals representing 13 protons as characteristic of carbazole alkaloid. In the down field region three singlets; intramolecular bonding OH-(δ 11.66) with -CHO (δ 9.91,C-3) and one broad signal due to -NH (δ 8.21) and another singlet proton H-4 at δ 8.03 (s,1H).In the aromatic region four signals with δ values 7.26 (td, J=3.8, 7.7Hz, 1H), 7.40 (td, J=3.8, 7.7Hz, 1H) showed the absence of substituent in ring A. the presence of prenyl group was revealed by one methylene protons at δ 3.64 (d, J= 6.9Hz,2H), δ 5.33 (t, J= 6.9Hz,1H) and one dimethyl group (δ 1.89(s,3H) , δ 1.72(s, 3H) (Table 2, Figure 1).

According to ¹³C NMR and DEPT (90, 135) spectra it was revealed that the presence of 18 signals and 18 carbons. Furthermore, the DEPT spectrum showed the presence of one aldehyde carbon, 8 quaternary carbons, 6 methine carbons, one methylene carbon and two methyl carbon as shown in table 2. The 2D NMR spectra such as DQF-COSY revealed the correlation of two adjacent protons. The HSQC spectrum gave the direct connection between protons and carbon. The HMBC data showed the position of H-4 singlet proton to C-1a, C-2, C-3, C-4a, -CHO. The presence and attachment of prenyl group to core carbazole alkaloid was revealed by the proton H-1' to C-1,C-1a C-2, C-10, C-11 (Table 2). Finally, the combination of fragments and attachments of substituents were confirmed by HMBC spectrum and compound (**1**) was elucidated as 7-hydroxyheptaphylline. It was also agreement with literature values.

The isolated compound was tested for their cytotoxicity on HeLa cell lines (Table 1). Both of them showed toxicity against to cervix cancer cell (HeLa) with IC₅₀ values 41.4 and μg/ml, where doxorubicin was used as positive control.

4 CONCLUSIONS

Nowadays the demand for anticancer drugs which are effective and less side effects are increasing. To fulfill this aim we conducted the isolation of bioactive compounds from root of *C. excavata*. The isolated compound was evaluated for their cytotoxicity. The compound (**1**), 7-hydroxyheptaphylline with IC₅₀ 41.4 μg/ml. So the result revealed that the compound-**1** should be further study for a potential natural anticancer candidate for HeLa cell in future.

ACKNOWLEDGEMENTS

Authors would like to thank to Universitas Airlangga and Ministry of Research, Technology and Higher Education for research grant. We also would like to acknowledge Ms. Helda Dwi Hardiyanti and Universitas Gadjah Mada for cytotoxicity measurements.

REFERENCES

- Amin, A., Gali-muhtasib, H., Ocker, M., Schneider-stock, R., 2009. Overview of Major Classes of Plant-Derived Anticancer Drugs 5, 1–11.
- Azarifar, Z., Mortazavi, M. M., Farhadian, R., Parvari, S., Mohammadi roushandeh, A., 2015. Cytotoxicity Effects of Aqueous Extract of *Purtulaca oleracea* on HeLa cell Line. Pharm. Sci. 21, 41–45.
- Cheng, S. S., Chang, H. T., Lin, C. Y., Chen, P. S., Huang, C. G., Chen, W. J., Chang, S. T., 2009. Insecticidal activities of leaf and twig essential oils from *Clausena excavata* against *Aedes aegypti* and *Aedes albopictus* larvae. Pest Manag. Sci. 65, 339–343.
- Kongkathip, N., Kongkathip, B., 2009. Constituents and bioactivities of *Clausena excavata*. Heterocycles 79, 121–144.
- Kumar, R., Saha, A., Saha, D., 2012. A new antifungal coumarin from *Clausena excavata*. Fitoterapia 83, 230–233.
- Mohan, D., 2012. *Clausena excavata* Burm. f. (Rutaceae): A review of its traditional uses, pharmacological and phytochemical properties. J. Med. Plants Res.
- N. W. Muhd Sharif, 2011. Cytotoxic constituents of *Clausena excavata*. African J. Biotechnol. 10, 16337–16341.
- Peh, T. H., Lim, G. K., Taufiq-yap, Y. H., Cheng, G., Ee, L., 2013. A New Cytotoxic Carbazole Alkaloid Isolated from the Stem Bark of Malaysian *Clausena excavata*. Can. Chem. Trans. 1, 165–172.
- Peng, W. W., Zheng, Y. Q., Chen, Y. S., Zhao, S. M., Ji, C. J., Tan, N. H., 2013. Coumarins from roots of *Clausena excavata*. J. Asian Nat. Prod. Res. 15, 215–220.
- Sunthitikawinsakul, A., Kongkathip, N., Kongkathip, B., Phonnakhu, S., Daly, J.W., Spande, T. F., Nimit, Y., Napaswat, C., Kasisit, J., Yoosook, C., 2003. Anti-HIV-1 Limonoid: First Isolation from *Clausena excavata*. Phyther. Res. 17, 1101–1103.
- Suwito, H., Hardiyanti, H. D., Ul Haq, K., Kristanti, A. N., Khasanah, M., 2018. (E)-3-[3-(4-morpholinophenyl) acryloyl]-2H-chromen-2-one. Molbank 2018, 1–5.
- Thant, T. M., Aminah, N. S., Kristanti, A. N., Ramadhan, R., Phuwapraisirisan, P., Takaya, Y., 2019. A new pyrano coumarin from *Clausena excavata* roots displaying dual inhibition against α -glucosidase and free radical. Nat. Prod. Res. 0, 1–6.
- Waziri, P. M., Abdullah, R., Yeap, S. K., Omar, A. R., Abdul, A. B., Kassim, N. K., Malami, I., Karunakaran, T., Imam, M. U., 2016a. Clausenidin from *Clausena excavata* induces apoptosis in hepG2 cells via the mitochondrial pathway. J. Ethnopharmacol. 194, 549–558.
- Waziri, P. M., Abdullah, R., Yeap, S. K., Omar, A. R., Kassim, N. K., Malami, I., How, C. W., Etti, I. C., Abu, M. L., 2016b. Clausenidin induces caspase-dependent apoptosis in colon cancer. BMC Complement. Altern. Med. 16, 1–12.

[HOME](#) [LOG IN](#) [ARCHIVE](#) [ANNOUNCEMENTS](#) [CALL FOR PAPER](#)

REGISTRATION FEE
[Home](#) > International Conference on Chemical Science and Technology Innovation (ICOCTI) >
[International Conference on Chemical Science and Technology Innovation \(ICOCTI 2019\)](#)

International Conference on Chemical Science and Technology Innovation (ICOCTI 2019)

The 1st International Conference on Chemical Science and Technology Innovation (ICOCTI) is an annual conference organized by Department Of Chemistry Universitas Sumatera Utara- Indonesia. The theme of ICOCTI 2019 is "*A Chemical Breakthrough for Scoience Technology Innovation*". The event is designed to emphasize advances and new findings in chemical science & technology innovation and their impacts on technology development. The conference will bring together scholars, leading researchers, and experts from diverse backgrounds and applications areas in Science. This Conference is organized by Department of Chemistry Universitas Sumatera Utara. The conference would offer a large number of invited lectures from renowned speakers all over the country. The Best paper awards will be given for the papers judged to make the most significant contribution to the conference.

Conference Theme "A Chemical Breakthrough for Scoience Technology Innovation"

CONFERENCE OBJECTIVES:

- To disseminate and develop the most recent innovations, trends, and concerns as well as practical challenges encountered and solutions in

<https://ocs.usu.ac.id/ICOCTI/ICOCTI2019?y=2019>

VENUE



Hotel Grandhika
Setiabudi

Jl. Setiabudi No.169
Medan

IMPORTANT DATE

Abstract Submission
deadline

30 Maret 2019

30 April 2019

27 May 2019

10 June 2019

Notification of Abstract

- To promote and facilitate the growth of scientific and technical development in the field of chemical advance material development in Indonesia and the Asia region.

12 April 2019

03 May 2019

31 May 2019

14 June 2019

Full Paper submission deadline

11 Mei 2019

28 June 2019

05 July 2019

16 July 2019

Notification of Accepted paper

8 Juni 2019

17 July 2019


Conference Date

18 - 19 July 2019

KEYNOTE/INVITED SPEAKER



Prof. Dr. Markus Fischer
University of Hamburg
Hamburg School of Food Sciences



Prof. Dr. Lee Hian Kee
Department of Chemistry
National University Singapore



Prof. Dr. Rizalman Mamat
Universiti Pahang
Malaysia



Dr. Saharman Gea
Universitas Sumatera Utara
Indonesia



Dr. Kamyar Shameli
Universiti Teknologi Malaysia
MUTT
Malaysia

Honorary Board

- Prof. Dr. Runtung Sitepu. Sh. M.Hum
- Drs. Mahyuddin K. M. Nasution. MIT, Ph.D

Editorial Board

- Prof. Dr. Basuki Wirjosentono. MS (USU-Indonesia)
- Prof. Dr. Harlem Marpaung (USU-Indonesia)
- Prof. Dr. Seribima Sembiring. M. Sc (USU-Indonesia)
- Prof. Dr. Zul Alfian. M.Sc (USU-Indonesia)
- Prof. Dr. Rizalman Mamat (UPM-Malaysia)
- Prof. Dr. Zuriati Zakaria (UTM-Malaysia)
- Dr. Kamyar Shameli
- Dr. Irwana Nainggolan
- Dr. Saharman Gea (USU- Indonesia)

Organizing Committee

- | | | |
|-----------|---|--------------------------------|
| Chairman | : | Prof. Dr. Harry Agusnar. M. Sc |
| Co-Chair | : | Dr. Cut Fatimah Zuhra |
| Secretary | : | Dr. M. Taufik. M. Si |
| Treasurer | : | Dr. Sovia Lenny. M. Si |

PUBLISHED BY



INDEXED BY



USER

Username _____

Password _____

Remember me[Log In](#)

CONTACT INFO

Organizing Committee
The 1st ICOCTI 2019

Email : icocsti@usu.ac.id

Secretariat

Department of
Chemistry Faculty of
Matematic and Natural
Science, Universitas
Sumatera Utara Jl.
Bioteknologi. No.1
Kampus USU Medan
20155- INDONESIA